

Supplementary Data

SUPPLEMENTARY TABLE S1. THE CHEMICAL PROFILE OF CI_{SCFE}

<i>Number</i>	<i>Components</i>	<i>Percentage</i>
1	Camphene	0.475 ^a
2	β -Cymene	0.998 ^a
3	Eucalyptol	3.091 ^a
4	Linalool oxide	0.521 ^a
5	α -Thujone	2.186 ^a
6	β -Thujone	2.169 ^a
7	Isothujol	1.094 ^a
8	L-Pinocarveol	0.765 ^a
9	d-Camphor	8.582 ^a
10	Cis-Verbenol	4.720 ^a
11	Endo-Borneol	7.845 ^a
12	L-4-Terpineol	1.634 ^a
13	α -Terpineol	1.022 ^a
14	Myrtenol	1.054 ^a
15	Cumaldehyde	0.486 ^a
16	Bornyl acetate	2.948 ^a
17	Thymol	3.071 ^a
18	β -Caryophyllene	3.336 ^a
19	Cis- β -Farnesene	2.270 ^a
20	α -Curcumene	5.932 ^a
21	σ -Cadinene	1.815 ^a
22	Spathulenol	1.362 ^a
23	Caryophyllene oxide	8.460 ^a
24	γ -Eudesmol	1.568 ^a
25	T-Muurolol	1.487 ^a
26	α -Gurjunene	2.161 ^a
27	Aromadendrene	2.280 ^a
28	α -Bisabolol	2.289 ^a
29	Cubenol	1.742 ^a
30	Longifolenaldehyde	2.572 ^a
31	α -Bisabolol oxide	2.600 ^a
32	Hexahydrofarnesyl acetone	1.212 ^a
33	Ethyl hexadecanoate	1.362 ^a
34	α -Linolenic acid	2.130 ^a
35	Ethyl octadec-9,12-dienoate	2.470 ^a
36	Chlorogenic acid	2.110 ^b
37	Luteolin-7-glucoside	2.800 ^b
38	Linarin	4.830 ^b
39	Luteolin	1.140 ^b

^aDetermined by GC-MS analysis, and the relative percentage calculated by integrated peak area in Agilent MSD Chemstation data analysis program.

^bAnalyzed and quantified by HPLC-PAD analysis, and the relative percentage was represented by the content quantitatively analyzed with peak areas under the standard curves.

CI_{SCFE}, *C. indicum* super-critical carbon dioxide fluid extract; GC-MS, gas chromatography-mass spectrometry; HPLC-PAD, high-performance liquid chromatography photodiode array detector.