

Online supplement

ETHNIC DIFFERENCES IN ASSOCIATIONS BETWEEN BLOOD PRESSURE AND STROKE IN SOUTH ASIAN AND EUROPEAN MEN

Running title: BP and stroke in South Asians and Europeans

Authors: Sophie V Eastwood MSc¹, Therese Tillin MSc¹, Nish Chaturvedi MD¹, Alun D Hughes PhD¹

Affiliations:¹UCL Institute of Cardiovascular Science, University College London, Gower Street, London, United Kingdom, WC1E 6BT.

Corresponding author: Dr Sophie V Eastwood, UCL Institute of Cardiovascular Science, UCL, Gower Street, London, United Kingdom, WC1E 6BT. Email: sophie.eastwood@ucl.ac.uk, Tel: +44 (0)207 679 9434.

Table S1. Hypertension control at follow-up clinic (2008-2011) in European and South Asian men. SBP=systolic blood pressure, DBP= diastolic blood pressure. Data are n(%) or mean±SD. *Diabetes vs. no diabetes p<0·05, †p value for ethnic difference in all participants, ‡p value for ethnic difference in those without diabetes, §p value for ethnic difference in those with diabetes.

BP treatment/ measure	Europeans			South Asians			p†	p‡	p§
	All	No diabetes	Diabetes	All	No diabetes	Diabetes			
On anti-hypertensive medication									
hypertensive	299(56)	212(50)	87(84)*	347(78)	174(70)	173(89)*	<0·001	<0·001	0·30
+									
BP≤140/90 mmHg	156(52)	113(53)	43(49)	161(47)	77(45)	84(49)	0·15	0·09	0·90
SBP, mmHg	140 ±17	140±17	140±17	143±18	142±17	144±20	0·05	0·30	0·11
DBP, mmHg	76 ±10	77±10	76±10	75±10	75±10	74±10	0·04	0·18	0·20

Table S2. Associations between blood pressure measures and incident stroke in men without baseline diabetes. Participants are men not in receipt of baseline anti-hypertensive medication. Data are odds ratios (OR) for a 1SD increase in BP measure unless otherwise stated, * $p<0.05$, † $p<0.01$, ‡ $p<0.001$, § p for ethnicity x BP interaction. ||Cardiovascular disease (CVD) risk factors comprise: smoking, WHR, total/ HDL- cholesterol ratio, diabetes, fasting glucose, physical activity and heart rate. SBP=systolic blood pressure, DBP=diastolic blood pressure, MAP= mean arterial blood pressure PP=pulse pressure.

BP measure	Model factors	Europeans		South Asians		p\$
		OR	95% CI	OR	95% CI	
SBP	Age	1.25	1.01,1.53*	1.72	1.34,2.20‡	0.08
	Age + CVD risk factors	1.28	1.03,1.61*	1.66	1.25,2.19‡	0.13
DBP	Age	0.91	0.73,1.13	1.95	1.50,2.54‡	<0.001
	Age + CVD risk factors	0.90	0.71,1.15	1.85	1.37,2.50‡	<0.001
MAP	Age	1.05	0.85,1.30	1.89	1.47,2.43‡	0.001
	Age + CVD risk factors	1.07	0.84,1.35	1.81	1.36,2.42‡	0.001
PP	Age	1.47	1.20,1.81‡	1.28	0.97,1.70	0.35
	Age + CVD risk factors	1.49	1.20,1.85‡	1.27	0.94,1.73	0.28

Table S3. Associations between blood pressure and incident stroke, accounting for time-varying effects of diabetes diagnosis, by ethnicity. Participants are men not in receipt of baseline anti-hypertensive medication. Data are odds ratios (OR) for a 1SD increase in BP measure, * $p<0\cdot05$, † $p<0\cdot01$, ‡ $p<0\cdot001$, § p for ethnicity x BP interaction. The time-varying covariate for diabetes diagnosis was categorised as follows: no diabetes ever, diabetes diagnosed after 2009, diabetes diagnosed between 1999 and 2008, diabetes diagnosed between 1988 and 1998, baseline diabetes. SBP=systolic blood pressure, DBP=diastolic blood pressure, MAP= mean arterial blood pressure PP=pulse pressure.

BP measure	Model factors	Europeans		South Asians		$p\$$
		OR	95% CI	OR	95% CI	
SBP	Age	1·21	1·00,1·47*	1·63	1·34,1·97‡	0·04
	Age + diabetes (time-varying)	1·22	0·98,1·52	1·46	1·16,1·85†	0·10
DBP	Age	0·91	0·74, 1·13	1·64	1·33,2·03‡	<0·001
	Age + diabetes (time-varying)	0·95	0·76, 1·19	1·57	1·21,2·03‡	0·004
MAP	Age	1·04	0·85, 1·28	1·69	1·38,2·08‡	0·001
	Age + diabetes (time-varying)	1·07	0·86,1·34	1·59	1·38,2·03‡	0·009
PP	Age	1·39	1·15, 1·68†	1·35	1·10, 1·64†	0·73
	Age + diabetes (time-varying)	1·37	1·11,1·68†	1·18	0·94,1·48	0·92

Table S4. Associations between blood pressure and incident stroke, by age group.**Excludes participants with baseline stroke or receipt of anti-hypertensive medication.**Data are odds ratios (OR) for a 1SD increase in BP measure unless stated, * $p<0.05$, † $p<0.01$,‡ $p<0.001$, § p for age group x BP interaction, || p for ethnicity x BP interaction. ¶CVD risk

factors comprise: smoking, WHR, total/ HDL- cholesterol ratio, diabetes, fasting glucose,

physical activity, heart rate. SBP=systolic blood pressure, DBP=diastolic blood pressure,

MAP= mean arterial blood pressure PP=pulse pressure.

BP measure	Model factors	Europeans			South Asians		
		OR	95% CI	p§	OR	95% CI	p
a) <55 year, events/N: Europeans 46/823, South Asians 59/768							
SBP	Age	1.02	0.72,1.43	0.25	1.74	1.34,2.27‡	0.02
	+ CVD risk factors¶	1.04	0.71,1.53	0.54	1.58	1.16,2.16†	0.03
DBP	Age	0.99	0.73,1.34	0.42	1.74	1.33,2.28‡	0.006
	+ CVD risk factors¶	1.05	0.75,1.47	0.19	1.60	1.16,2.21†	0.02
MAP	Age	1.00	0.73,1.36	0.80	1.79	1.38,2.33‡	0.005
	+ CVD risk factors¶	1.05	0.74,1.49	0.73	1.65	1.21,2.27†	0.01
PP	Age	1.04	0.71,1.52	0.10	1.37	1.04,1.80*	0.27
	+ CVD risk factors¶	1.00	0.66,1.52	0.12	1.27	0.93,1.73	0.21
SBP	Age	1.07	0.63,1.78	0.08	1.40	0.94,2.08	0.44
DBP	Age	0.95	0.61,1.51	0.008	1.35	0.91,2.01	0.24
SBP	+ CVD risk factors¶	1.00	0.56,1.77	0.15	1.32	0.86,2.03	0.34
DBP	+ CVD risk factors¶	1.05	0.63,1.73	0.09	1.31	0.84,2.04	0.42
MAP	Age	0.99	0.71,1.37	0.65	1.76	1.31,2.36*	0.01
PP	Age	1.05	0.71,1.37	0.49	1.05	0.76,1.45	0.96
MAP	+ CVD risk factors¶	1.05	0.73,1.53	0.34	1.63	1.15,2.31†	0.03
PP	+ CVD risk factors¶	0.98	0.64,1.52	0.47	1.03	0.73,1.46	0.83
b) ≥55 years, events /N: Europeans 58/553, South Asians 43/306							
SBP	Age	1.34	1.05,1.70*	-	1.48	1.10,1.98†	0.62
	+ CVD risk factors¶	1.33	1.02,1.74*	-	1.56	1.11,2.21†	0.40
DBP	Age	0.84	0.63,1.12	-	1.44	1.02,2.04*	0.02
	+ CVD risk factors¶	0.78	0.56,1.08	-	1.87	1.22,2.87†	0.004
MAP	Age	1.07	0.82,1.40	-	1.52	1.10,2.10*	0.10
	+ CVD risk factors¶	1.13	0.58,2.23	-	1.81	1.22,2.68†	0.03

PP	Age	1·58	1·26,1·99‡	-	1·33	1·00,1·77*	0·36	-
	+ CVD risk factors¶	1·63	1·27,2·11‡	-	1·24	0·89,1·73	0·42	-
SBP		1·94	1·41,2·67‡	-	1·39	0·94,2·07	0·23	-
DBP	Age	0·51	0·36,0·75†	-	1·11	0·69,1·77	0·01	-
SBP		2·03	1·42,2·93‡	-	1·21	0·77,1·92	0·24	-
DBP	+ CVD risk factors¶	0·47	0·31,0·72‡	-	1·61	0·92,2·81	0·005	-
MAP		0·77	0·56, 1·80	-	1·41	0·96,2·07	0·02	-
PP	Age	1·76	1·35,2·31‡	-	1·13	0·81,1·58	0·05	-
MAP		0·75	0·51,1·03	-	1·92	1·20,3·07†	0·005	-
PP	+ CVD risk factors¶	1·87	1·39,2·54‡	-	0·92	0·62,1·36	0·03	-