Web Appendix A. Country-specific stroke (IS and HS) burden counts in 1990 and 2013 by sex.

							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Afghanistan	Deaths	Male	1990	2,357	(1,593 to 3,334)	217.4	(152.9 to 292.1)
			2013	8,261	(5,476 to 11,620)	231.1	(162.3 to 307.3)
		Female	1990	4,560	(2,980 to 7,091)	300	(209.2 to 432.6)
			2013	15,016	(9,911 to 23,549)	329.9	(229.5 to 480.5)
		Both	1990	6,918	(5,172 to 9,714)	262.8	(207.7 to 341.9)
			2013	23,277	(16,937 to 31,926)	284.3	(218.6 to 368.2)
	DALYs	Male	1990	65,016	(42,544 to 93,712)	3945.4	(2701.3 to 5513.6)
			2013	210,046	(135,499 to 310,295)	4152.9	(2773.5 to 5781.6)
		Female	1990	132,316	(82,722 to 217,528)	6102.8	(4030.8 to 9384.4)
			2013	402,767	(248,174 to 672,352)	6621.6	(4348.2 to 10451.6)
		Both	1990	197,332	(143,135 to 289,787)	5091	(3856.3 to 6997.3)
			2013	612,813	(427,918 to 879,037)	5432	(3961.3 to 7437.3)
	YLDs	Male	1990	212	(137 to 301)	10.4	(6.7 to 15.1)
			2013	651	(431 to 915)	10.8	(7.3 to 15.2)
		Female	1990	188	(118 to 272)	8.7	(5.7 to 12.3)
			2013	553	(363 to 781)	8.8	(5.8 to 12.3)
		Both	1990	401	(264 to 553)	9.5	(6.4 to 13.0)
			2013	1,205	(830 to 1,658)	9.8	(6.8 to 13.3)
Albania	Deaths	Male	1990	1,422	(1,261 to 1,591)	194.1	(172.4 to 217.1)
			2013	3,213	(2,499 to 4,025)	216.2	(171.2 to 270.1)
		Female	1990	1,611	(1,462 to 1,845)	178.7	(161.7 to 204.1)
			2013	3,056	(2,478 to 3,780)	169.4	(136.9 to 209.0)
		Both	1990	3,033	(2,833 to 3,321)	186.1	(173.4 to 205.3)

1			2013	6,269	(5,345 to 7,314)	191.6	(164.1 to 223.1)
	DALYs	Male	1990	26,809	(23,645 to 30,171)	2843.1	(2516.9 to 3173.7)
_			2013	51,152	(38,591 to 65,968)	3154.6	(2415.8 to 4027.2)
_		Female	1990	25,161	(22,668 to 29,126)	2444.9	(2189.4 to 2784.9)
			2013	38,531	(30,131 to 49,810)	2139.2	(1668.9 to 2745.8)
_		Both	1990	51,970	(48,314 to 57,103)	2635.4	(2452.1 to 2869.9)
			2013	89,683	(74,215 to 107,491)	2625.2	(2179.7 to 3130.6)
	YLDs	Male	1990	305	(195 to 437)	27.4	(17.7 to 39.1)
			2013	526	(348 to 763)	30.7	(20.6 to 44.1)
		Female	1990	208	(139 to 293)	18.7	(12.4 to 26.2)
			2013	331	(219 to 479)	18.3	(12.2 to 26.6)
_		Both	1990	514	(353 to 700)	23	(15.7 to 31.4)
			2013	857	(579 to 1,196)	24.3	(16.5 to 33.9)
Algeria	Deaths	Male	1990	5,681	(4,576 to 7,010)	150.2	(121.8 to 184.1)
	_				, , ,		•
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
	measure	sex	year 2013	number 10,152		Rate 115.7	
	measure	sex Female			Uncertainty Interval		Interval
	measure		2013	10,152	Uncertainty Interval (8,189 to 12,346)	115.7	Interval (93.6 to 138.6)
	measure		2013 1990	10,152 7,268	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838)	115.7 154.4	Interval (93.6 to 138.6) (129.7 to 185.9)
	measure	Female	2013 1990 2013	10,152 7,268 12,294	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116)	115.7 154.4 126.1	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9)
	measure DALYs	Female	2013 1990 2013 1990	10,152 7,268 12,294 12,949	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079)	115.7 154.4 126.1 152.4	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0)
		Female Both	2013 1990 2013 1990 2013	10,152 7,268 12,294 12,949 22,446	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850)	115.7 154.4 126.1 152.4 121.2	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4)
		Female Both	2013 1990 2013 1990 2013 1990	10,152 7,268 12,294 12,949 22,446 141,644	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850) (112,242 to 177,400)	115.7 154.4 126.1 152.4 121.2 2418.6	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4) (1926.6 to 2992.8)
		Female Both Male	2013 1990 2013 1990 2013 1990 2013	10,152 7,268 12,294 12,949 22,446 141,644 197,976	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850) (112,242 to 177,400) (157,421 to 244,575)	115.7 154.4 126.1 152.4 121.2 2418.6 1706.5	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4) (1926.6 to 2992.8) (1374.5 to 2087.2)
		Female Both Male	2013 1990 2013 1990 2013 1990 2013 1990	10,152 7,268 12,294 12,949 22,446 141,644 197,976 171,808	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850) (112,242 to 177,400) (157,421 to 244,575) (137,176 to 220,419)	115.7 154.4 126.1 152.4 121.2 2418.6 1706.5 2533.2	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4) (1926.6 to 2992.8) (1374.5 to 2087.2) (2081.5 to 3120.0)
		Female Both Male Female	2013 1990 2013 1990 2013 1990 2013 1990 2013	10,152 7,268 12,294 12,949 22,446 141,644 197,976 171,808 225,830	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850) (112,242 to 177,400) (157,421 to 244,575) (137,176 to 220,419) (186,222 to 278,661)	115.7 154.4 126.1 152.4 121.2 2418.6 1706.5 2533.2 1845.6	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4) (1926.6 to 2992.8) (1374.5 to 2087.2) (2081.5 to 3120.0) (1528.7 to 2285.2)
		Female Both Male Female	2013 1990 2013 1990 2013 1990 2013 1990 2013	10,152 7,268 12,294 12,949 22,446 141,644 197,976 171,808 225,830 313,452	Uncertainty Interval (8,189 to 12,346) (6,053 to 8,838) (10,131 to 15,116) (11,199 to 15,079) (19,563 to 25,850) (112,242 to 177,400) (157,421 to 244,575) (137,176 to 220,419) (186,222 to 278,661) (264,396 to 374,085)	115.7 154.4 126.1 152.4 121.2 2418.6 1706.5 2533.2 1845.6 2474.7	(93.6 to 138.6) (129.7 to 185.9) (103.8 to 154.9) (133.1 to 175.0) (106.3 to 139.4) (1926.6 to 2992.8) (1374.5 to 2087.2) (2081.5 to 3120.0) (1528.7 to 2285.2) (2118.3 to 2900.0)

		Female	1990	581	(391 to 812)	9.3	(6.2 to 13.0)
			2013	1,205	(789 to 1,678)	8.9	(5.9 to 12.3)
		Both	1990	1,195	(820 to 1,620)	9.9	(6.8 to 13.3)
			2013	2,755	(1,851 to 3,754)	10.2	(6.8 to 13.9)
Andorra	Deaths	Male	1990	19	(13 to 26)	73.3	(52.5 to 102.4)
			2013	30	(21 to 41)	47.2	(33.3 to 65.6)
		Female	1990	31	(20 to 42)	62.7	(39.5 to 85.1)
			2013	44	(27 to 65)	40.2	(24.9 to 59.9)
		Both	1990	50	(38 to 64)	67.6	(51.4 to 85.6)
			2013	74	(55 to 99)	43.6	(32.6 to 57.7)
	DALYs	Male	1990	283	(204 to 391)	1039.1	(749.0 to 1435.8)
			2013	377	(283 to 517)	628.3	(469.2 to 863.5)
		Female	1990	306	(199 to 410)	690.9	(448.7 to 939.6)
			2013	366	(251 to 527)	406.4	(272.0 to 581.4)
		Both	1990	589	(461 to 741)	841.4	(654.1 to 1072.1)
			2013	744	(589 to 952)	507.5	(398.2 to 651.0)
	YLDs	Male	1990	30	(20 to 41)	110.7	(73.4 to 149.1)
			2013	59	(39 to 83)	102.9	(67.7 to 143.1)
		Female	1990	30	(20 to 43)	77.1	(50.5 to 110.2)
			2013	53	(35 to 73)	72.9	(48.3 to 99.5)
		Both	1990	61	(42 to 82)	91.2	(62.8 to 122.4)
			2013	113	(78 to 150)	86.5	(60.0 to 115.4)
Angola	Deaths	Male	1990	3,088	(1,918 to 4,280)	214.5	(138.0 to 288.9)
			2013	5,238	(3,910 to 7,752)	197.6	(152.1 to 279.1)
		Female	1990	3,514	(2,253 to 5,115)	224.3	(147.6 to 314.1)
			2013	6,180	(4,259 to 9,303)	210.9	(147.7 to 306.8)
		Both	1990	6,603	(4,288 to 9,283)	221.6	(148.7 to 302.5)
			2013	11,419	(8,494 to 16,751)	205.9	(156.8 to 290.7)

	DALYs	Male	1990	90,454	(57,576 to 126,396)	4282.6	(2660.4 to 5942.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	141,298	(102,913 to 210,035)	3614	(2703.9 to 5369.8)
		Female	1990	89,356	(55,613 to 131,073)	3983.8	(2543.3 to 5850.1)
			2013	142,582	(95,861 to 222,576)	3487.1	(2383.6 to 5273.3)
		Both	1990	179,810	(119,880 to 251,680)	4142	(2662.8 to 5843.9)
			2013	283,880	(205,860 to 419,748)	3560.2	(2630.1 to 5252.4)
	YLDs	Male	1990	364	(237 to 528)	17.6	(11.4 to 24.9)
			2013	755	(496 to 1,072)	17.8	(11.7 to 24.8)
		Female	1990	308	(203 to 431)	13.1	(8.5 to 18.4)
			2013	676	(444 to 949)	14.6	(9.5 to 20.7)
		Both	1990	673	(454 to 933)	15.2	(10.2 to 21.0)
			2013	1,432	(979 to 1,932)	16.1	(11.0 to 21.8)
Antigua and							
Barbuda	Deaths	Male	1990	27	(23 to 33)	145.3	(120.8 to 173.4)
			2013	33	(25 to 43)	114.1	(85.8 to 145.5)
		Female	1990	41	(32 to 46)	144.5	(111.4 to 163.5)
			2013	38	(28 to 52)	93.6	(69.7 to 129.8)
		Both	1990	69	(58 to 77)	147	(124.0 to 163.5)
			2013	71	(58 to 86)	103.1	(84.9 to 124.6)
	DALYs	Male	1990	548	(455 to 655)	2603	(2183.2 to 3097.7)
			2013	662	(490 to 880)	1960.2	(1467.2 to 2557.3)
		Female	1990	619	(481 to 708)	2232	(1734.0 to 2560.5)
			2013	575	(411 to 810)	1385.1	(989.4 to 1952.1)
		Both	1990	1,167	(1,007 to 1,304)	2414.8	(2085.0 to 2702.7)
			2013	1,237	(1,006 to 1,529)	1650	(1345.4 to 2026.6)
	YLDs	Male	1990	12	(8 to 17)	60	(40.1 to 83.2)
			2013	18	(12 to 26)	57.5	(37.3 to 80.1)

		ı	1				Ī
		Female	1990	11	(7 to 16)	43.1	(28.3 to 59.5)
			2013	17	(11 to 24)	42.1	(27.6 to 60.9)
		Both	1990	24	(16 to 32)	50.4	(34.8 to 67.6)
			2013	36	(24 to 49)	49	(33.3 to 68.0)
Argentina	Deaths	Male	1990	13,294	(11,290 to 16,032)	114.8	(97.0 to 138.9)
			2013	15,178	(12,482 to 17,925)	85	(69.6 to 100.6)
		Female	1990	15,033	(11,528 to 16,631)	94	(72.2 to 104.0)
			2013	15,222	(12,368 to 22,018)	51.6	(42.5 to 74.2)
		Both	1990	28,327	(24,488 to 31,225)	103.6	(89.5 to 114.3)
			2013	30,401	(26,259 to 37,641)	65.4	(56.7 to 80.4)
	DALYs	Male	1990	278,682	(239,551 to 334,408)	2120	(1815.0 to 2542.9)
			2013	278,262	(232,039 to 324,882)	1478.9	(1232.0 to 1722.0)
		Female	1990	245,505	(193,404 to 268,482)	1515.2	(1193.5 to 1657.2)
			2013	200,612	(170,726 to 282,563)	789.3	(674.7 to 1102.3)
		Both	1990	524,187	(464,591 to 582,552)	1791.6	(1580.9 to 1988.6)
			2013	478,875	(424,934 to 571,826)	1091.7	(967.9 to 1302.1)
	YLDs	Male	1990	6,857	(4,687 to 9,040)	51.7	(35.6 to 68.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	8,059	(5,519 to 11,104)	42.4	(29.1 to 58.3)
		Female	1990	4,809	(3,236 to 6,547)	29.3	(19.9 to 39.8)
			2013	6,119	(4,044 to 8,794)	25.1	(16.7 to 36.0)
		Both	1990	11,666	(8,000 to 15,285)	39.4	(27.0 to 51.6)
			2013	14,178	(9,649 to 19,601)	32.7	(22.2 to 45.1)
Armenia	Deaths	Male	1990	1,311	(1,123 to 1,552)	153.9	(131.1 to 183.4)
			2013	1,751	(1,377 to 2,104)	125.9	(99.0 to 151.1)
		Female	1990	1,726	(1,518 to 2,033)	131.6	(115.6 to 155.6)
			2013	2,082	(1,614 to 2,488)	102.2	(80.1 to 121.9)
		Both	1990	3,038	(2,769 to 3,426)	140.6	(128.0 to 158.8)

			2013	3,834	(3,263 to 4,322)	112.8	(96.0 to 127.3)
	DALYs	Male	1990	26,018	(22,193 to 30,846)	2480.3	(2124.4 to 2933.2)
			2013	28,155	(22,348 to 33,981)	1954.8	(1552.3 to 2360.5)
		Female	1990	27,200	(23,908 to 32,590)	1906.7	(1685.0 to 2269.4)
			2013	26,275	(21,288 to 31,724)	1392.9	(1137.9 to 1689.2)
		Both	1990	53,218	(48,508 to 60,378)	2149.6	(1961.8 to 2432.7)
			2013	54,431	(47,660 to 61,887)	1647.8	(1446.7 to 1875.6)
	YLDs	Male	1990	198	(130 to 282)	15.6	(10.2 to 22.1)
			2013	246	(156 to 343)	16.4	(10.5 to 22.8)
		Female	1990	195	(123 to 282)	12.6	(8.0 to 18.2)
			2013	226	(148 to 313)	12.8	(8.4 to 17.8)
		Both	1990	394	(268 to 530)	14	(9.6 to 19.1)
			2013	473	(324 to 628)	14.5	(10.1 to 19.4)
Australia	Deaths	Male	1990	5,001	(4,111 to 6,289)	72.1	(59.0 to 90.4)
			2013	5,977	(4,565 to 7,429)	37.5	(28.7 to 46.4)
		Female	1990	7,229	(5,913 to 8,659)	66.7	(54.6 to 79.9)
			2013	8,655	(7,042 to 11,336)	38.9	(31.9 to 51.0)
		Both	1990	12,230	(10,934 to 14,104)	69.6	(62.1 to 80.7)
			2013	14,633	(12,318 to 17,590)	38.6	(32.3 to 45.8)
	DALYs	Male	1990	81,775	(68,896 to 102,216)	1054	(886.6 to 1308.9)
			2013	77,850	(60,288 to 93,064)	513.7	(396.3 to 613.8)
		Female	1990	85,491	(71,541 to 101,916)	829.9	(693.7 to 994.2)
			2013	80,773	(67,749 to 104,098)	431.1	(363.1 to 553.5)
		Both	1990	167,267	(150,956 to 193,230)	931.9	(841.0 to 1073.6)
			2013	158,623	(134,312 to 184,098)	471.5	(394.7 to 546.8)
	YLDs	Male	1990	6,314	(4,324 to 8,831)	77.4	(53.3 to 107.7)
			2013	11,371	(7,796 to 15,335)	76.7	(52.3 to 103.3)
		Female	1990	5,086	(3,539 to 7,100)	51.7	(35.9 to 71.7)

			2013	8,302	(5,533 to 11,250)	50.6	(33.8 to 68.3)
		Both	1990	11,401	(7,969 to 15,361)	63.3	(44.2 to 85.1)
			2013	19,673	(13,810 to 25,840)	62.9	(44.0 to 82.6)
Austria	Deaths	Male	1990	3,676	(3,087 to 4,433)	96	(80.3 to 115.7)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
			2013	3,010	(2,357 to 3,690)	45.4	(35.8 to 55.6)
		Female	1990	7,061	(5,247 to 7,805)	86.7	(64.7 to 95.3)
			2013	4,612	(3,605 to 7,206)	40.2	(31.9 to 62.0)
		Both	1990	10,738	(9,003 to 11,821)	90.8	(76.6 to 99.6)
			2013	7,623	(6,435 to 10,063)	42.9	(36.4 to 56.0)
	DALYs	Male	1990	57,137	(48,456 to 68,205)	1447.7	(1230.6 to 1726.5)
			2013	44,374	(35,804 to 52,887)	692.5	(560.0 to 824.7)
		Female	1990	80,435	(62,272 to 87,673)	1126.5	(882.3 to 1230.1)
			2013	46,113	(37,971 to 66,542)	496.1	(411.5 to 710.8)
		Both	1990	137,572	(119,338 to 151,005)	1258.2	(1097.3 to 1383.0)
			2013	90,487	(78,622 to 111,453)	586.7	(511.6 to 706.6)
	YLDs	Male	1990	4,252	(2,870 to 5,798)	106.3	(71.8 to 144.2)
			2013	7,453	(4,869 to 10,369)	117.3	(77.1 to 162.2)
		Female	1990	4,914	(3,328 to 6,765)	75.8	(51.3 to 104.2)
			2013	6,121	(3,989 to 8,593)	74.6	(49.6 to 104.2)
		Both	1990	9,166	(6,336 to 12,288)	87.5	(60.8 to 116.8)
			2013	13,574	(9,257 to 18,405)	93.7	(64.5 to 126.1)
Azerbaijan	Deaths	Male	1990	2,296	(1,982 to 2,753)	182.3	(158.1 to 218.4)
			2013	3,894	(3,030 to 4,776)	153.7	(121.1 to 186.4)
		Female	1990	2,955	(2,660 to 3,512)	151.7	(135.7 to 182.1)
			2013	4,707	(3,379 to 5,690)	136.6	(98.0 to 163.8)
		Both	1990	5,251	(4,832 to 5,983)	163.8	(151.0 to 187.7)
			2013	8,601	(6,896 to 9,927)	144.4	(115.8 to 165.9)

		,			T		
	DALYs	Male	1990	55,783	(47,691 to 67,173)	3196.8	(2763.1 to 3819.3)
			2013	78,133	(60,250 to 98,032)	2502	(1942.0 to 3095.4)
		Female	1990	57,353	(51,107 to 68,723)	2422.3	(2167.8 to 2893.1)
			2013	76,480	(55,139 to 94,166)	1990.8	(1431.3 to 2436.1)
		Both	1990	113,136	(103,317 to 128,605)	2743.4	(2521.0 to 3127.2)
			2013	154,613	(125,326 to 180,427)	2220.6	(1783.2 to 2574.1)
	YLDs	Male	1990	334	(225 to 470)	15.9	(10.7 to 22.1)
			2013	622	(414 to 863)	16.7	(11.1 to 22.9)
		Female	1990	314	(204 to 443)	12.2	(7.9 to 17.1)
			2013	525	(351 to 728)	12.4	(8.3 to 17.3)
		Both	1990	648	(450 to 868)	13.9	(9.7 to 18.6)
			2013	1,148	(789 to 1,553)	14.4	(9.9 to 19.6)
Bahrain	Deaths	Male	1990	76	(60 to 107)	100.6	(78.3 to 137.7)
			2013	92	(64 to 125)	48.3	(34.5 to 64.5)
		Female	1990	71	(59 to 95)	121.8	(100.4 to 165.1)
			2013	99	(74 to 126)	65.4	(48.6 to 83.0)
		Both	1990	148	(127 to 189)	109.8	(94.4 to 139.1)
			2013	192	(153 to 231)	57	(46.3 to 68.2)
	DALYs	Male	1990	2,104	(1,664 to 3,075)	1702.8	(1333.8 to 2378.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	2,356	(1,613 to 3,215)	683.2	(486.1 to 928.2)
		Female	1990	1,924	(1,548 to 2,449)	2119.6	(1763.4 to 2790.3)
			2013	1,975	(1,439 to 2,616)	878.8	(647.6 to 1129.6)
		Both	1990	4,029	(3,457 to 5,100)	1881.4	(1615.0 to 2385.8)
			2013	4,332	(3,386 to 5,326)	777.1	(624.1 to 937.5)
	YLDs	Male	1990	15	(9 to 21)	10.8	(7.3 to 15.2)
			2013	54	(34 to 79)	10.2	(6.6 to 15.1)
		Female	1990	7	(4 to 10)	8.3	(5.3 to 12.0)

			2013	24	(15 to 34)	8	(5.2 to 11.4)
		Both	1990	22	(15 to 31)	9.8	(6.6 to 13.3)
			2013	78	(51 to 112)	9.3	(6.3 to 13.3)
Bangladesh	Deaths	Male	1990	29,398	(22,826 to 35,832)	134.8	(104.9 to 161.9)
			2013	86,545	(66,234 to 111,732)	224.7	(173.7 to 283.4)
		Female	1990	28,363	(21,980 to 38,673)	154.5	(122.3 to 208.0)
			2013	91,808	(47,559 to 122,186)	258.2	(133.3 to 340.9)
		Both	1990	57,761	(49,563 to 67,049)	143.8	(123.6 to 165.8)
			2013	178,354	(127,309 to 214,741)	240.8	(172.9 to 287.6)
	DALYs	Male	1990	628,762	(480,241 to 780,285)	2347.4	(1804.6 to 2881.5)
			2013	1,636,288	(1,211,022 to 2,188,158)	3542.4	(2672.0 to 4625.6)
		Female	1990	590,934	(453,641 to 816,830)	2660	(2063.6 to 3630.8)
			2013	1,667,167	(830,758 to 2,289,281)	4012.8	(2043.8 to 5414.1)
		Both	1990	1,219,697	(1,041,632 to 1,440,296)	2490.7	(2141.2 to 2896.0)
			2013	3,303,456	(2,311,184 to 4,064,255)	3765.3	(2654.5 to 4584.0)
	YLDs	Male	1990	12,520	(8,124 to 17,354)	42.8	(27.7 to 59.6)
			2013	23,040	(15,275 to 32,256)	43.9	(29.1 to 61.9)
		Female	1990	8,886	(6,057 to 12,175)	34.8	(23.5 to 48.1)
			2013	18,071	(12,021 to 24,717)	36.7	(24.8 to 51.0)
		Both	1990	21,406	(14,668 to 28,265)	39.1	(26.7 to 51.5)
			2013	41,112	(27,926 to 54,844)	40.5	(27.8 to 54.2)
Barbados	Deaths	Male	1990	120	(103 to 145)	125.2	(107.5 to 150.8)
			2013	124	(92 to 157)	94.7	(71.3 to 120.1)
		Female	1990	205	(163 to 242)	127.4	(101.1 to 149.5)
			2013	170	(130 to 236)	89.4	(68.1 to 122.9)
		Both	1990	326	(284 to 369)	127.8	(111.6 to 144.7)

			2013	294	(242 to 363)	92.6	(76.0 to 114.3)
	DALYs	Male	1990	2,123	(1,839 to 2,519)	2142	(1860.0 to 2548.0)
			2013	2,264	(1,668 to 2,856)	1541.1	(1145.8 to 1934.6)
		Female	1990	2,701	(2,184 to 3,152)	1823.5	(1475.9 to 2108.8)
							(950.4 to 1685.1)
			2013	2,323	(1,745 to 3,084)	1268	
							I I manufactura
Country	maacura	COV	vear	number	Uncertainty Interval	Rate	Uncertainty Interval
Country	measure	sex	year		•		
		Both	1990	4,824	(4,314 to 5,367)	1970.8	(1765.0 to 2180.6)
			2013	4,587	(3,795 to 5,529)	1401.5	(1163.9 to 1686.6)
	YLDs	Male	1990	58	(39 to 79)	59.4	(40.3 to 81.0)
			2013	81	(54 to 110)	55.8	(37.7 to 75.6)
		Female	1990	62	(40 to 88)	42.3	(27.4 to 59.1)
			2013	82	(56 to 111)	44.6	(30.5 to 60.0)
		Both	1990	120	(82 to 159)	49.3	(33.5 to 65.5)
			2013	163	(114 to 214)	49.6	(34.7 to 65.0)
Belarus	Deaths	Male	1990	6,347	(5,372 to 7,403)	172.5	(146.4 to 201.7)
			2013	8,576	(7,112 to 10,406)	181.5	(151.2 to 220.1)
		Female	1990	10,792	(9,194 to 12,152)	141.3	(120.2 to 158.9)
			2013	13,698	(11,389 to 16,015)	135.7	(114.8 to 158.7)
		Both	1990	17,139	(15,306 to 18,819)	153.5	(136.8 to 168.6)
			2013	22,275	(19,715 to 25,496)	155.4	(137.7 to 177.6)
	DALYs	Male	1990	131,809	(112,051 to 152,475)	3119.3	(2649.6 to 3612.2)
			2013	177,261	(145,868 to 219,277)	3464.2	(2857.2 to 4255.9)
		Female	1990	161,905	(137,524 to 181,286)	2217.7	(1877.7 to 2482.4)
			2013	183,803	(157,809 to 219,781)	2068.8	(1776.0 to 2484.8)
		Both	1990	293,715	(264,955 to 320,665)	2583.1	(2327.0 to 2827.4)
			2013	361,064	(321,250 to 420,045)	2655.6	(2363.1 to 3118.7)

	YLDs	Male	1990	1,810	(1,145 to 2,621)	40	(25.7 to 57.1)
			2013	2,229	(1,473 to 3,140)	43.2	(28.8 to 60.1)
		Female	1990	1,790	(1,165 to 2,570)	26.2	(17.0 to 37.5)
			2013	1,968	(1,225 to 2,799)	25.2	(16.2 to 35.6)
		Both	1990	3,600	(2,421 to 4,974)	31.8	(21.3 to 43.8)
			2013	4,197	(2,794 to 5,775)	32.6	(21.6 to 44.7)
Belgium	Deaths	Male	1990	4,326	(3,577 to 5,261)	81	(66.7 to 98.9)
			2013	4,321	(3,399 to 5,278)	48.1	(37.9 to 58.7)
		Female	1990	7,215	(5,531 to 8,241)	73.3	(56.5 to 83.3)
			2013	6,076	(4,852 to 8,600)	40.9	(33.1 to 57.9)
		Both	1990	11,542	(9,743 to 12,887)	77	(65.1 to 86.1)
			2013	10,398	(8,916 to 13,020)	44.2	(38.2 to 54.4)
	DALYs	Male	1990	69,540	(59,178 to 82,829)	1245.4	(1056.7 to 1481.0)
			2013	60,601	(48,832 to 72,117)	724.7	(586.5 to 861.9)
		Female	1990	84,041	(66,805 to 95,120)	988	(791.4 to 1127.9)
			2013	61,961	(51,249 to 84,824)	533.7	(442.9 to 728.3)
		Both	1990	153,581	(135,080 to 170,423)	1098.5	(966.6 to 1221.9)
			2013	122,562	(107,473 to 145,551)	620.2	(542.7 to 729.5)
	YLDs	Male	1990	5,864	(3,987 to 8,039)	102.8	(70.0 to 139.9)
			2013	8,322	(5,676 to 11,456)	102.5	(69.5 to 140.9)
		Female	1990	5,710	(3,902 to 7,921)	72.4	(49.2 to 99.9)
			2013	7,105	(4,783 to 9,927)	68.9	(47.1 to 93.8)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	11,575	(8,140 to 15,579)	84.9	(59.6 to 114.6)
			2013	15,428	(10,749 to 20,470)	84.3	(58.9 to 111.1)
Belize	Deaths	Male	1990	43	(35 to 51)	111.6	(91.3 to 132.9)
			2013	82	(63 to 102)	120.7	(93.8 to 149.4)
		Female	1990	46	(38 to 54)	107.6	(89.8 to 127.6)

			2013	99	(77 to 128)	125.6	(97.6 to 160.4)
		Both	1990	89	(78 to 101)	109.7	(95.7 to 124.5)
			2013	182	(154 to 216)	123.7	(105.5 to 145.5)
	DALYs	Male	1990	957	(810 to 1,126)	1995	(1660.6 to 2347.9)
			2013	1,803	(1,371 to 2,282)	2173.1	(1668.8 to 2720.5)
		Female	1990	897	(736 to 1,037)	1804.7	(1491.4 to 2107.0)
			2013	1,880	(1,416 to 2,538)	2091	(1582.6 to 2773.8)
		Both	1990	1,854	(1,624 to 2,073)	1899.2	(1658.1 to 2135.7)
			2013	3,683	(3,041 to 4,463)	2134	(1782.2 to 2554.2)
	YLDs	Male	1990	26	(17 to 36)	63.4	(42.9 to 88.6)
			2013	46	(30 to 62)	59	(39.4 to 80.0)
		Female	1990	18	(11 to 26)	41.2	(27.0 to 60.3)
			2013	36	(24 to 50)	42.1	(28.0 to 58.3)
		Both	1990	44	(30 to 60)	51.9	(36.0 to 71.7)
			2013	82	(56 to 112)	50.1	(34.7 to 67.5)
Benin	Deaths	Male	1990	951	(791 to 1,134)	127.5	(104.7 to 152.6)
			2013	2,198	(1,692 to 2,899)	160.6	(126.1 to 208.3)
		Female	1990	1,193	(966 to 1,488)	125.4	(100.7 to 156.8)
			2013	2,258	(1,658 to 3,043)	141.5	(104.7 to 189.9)
		Both	1990	2,145	(1,863 to 2,462)	126.8	(109.4 to 147.5)
			2013	4,457	(3,619 to 5,464)	150.6	(123.9 to 182.9)
	DALYs	Male	1990	27,446	(22,180 to 33,233)	2407.6	(1996.9 to 2864.3)
			2013	56,284	(42,573 to 74,675)	2867.3	(2219.2 to 3766.7)
		Female	1990	28,135	(21,958 to 34,735)	2102.6	(1706.1 to 2613.8)
			2013	47,171	(33,707 to 64,582)	2239.9	(1636.6 to 3026.6)
		Both	1990	55,582	(46,706 to 64,108)	2242.7	(1944.3 to 2569.4)
			2013	103,456	(84,480 to 126,846)	2536.4	(2063.0 to 3109.7)
	YLDs	Male	1990	205	(130 to 290)	19.9	(12.5 to 28.4)

			2013	445	(288 to 627)	19.5	(12.7 to 27.5)
		Female	1990	183	(118 to 263)	13.9	(8.8 to 20.1)
			2013	373	(238 to 527)	14.7	(9.4 to 20.6)
		Both	1990	388	(265 to 525)	16.4	(11.2 to 22.3)
			2013	818	(549 to 1,137)	16.9	(11.2 to 23.2)
Bhutan	Deaths	Male	1990	112	(69 to 169)	124.5	(79.3 to 176.6)
			2013	290	(187 to 415)	143.4	(95.5 to 199.3)
		Female	1990	109	(73 to 165)	137.4	(98.0 to 198.3)
			2013	252	(129 to 387)	156.3	(80.5 to 237.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	222	(148 to 330)	130.8	(92.6 to 183.2)
			2013	543	(334 to 797)	149.2	(93.3 to 213.5)
	DALYs	Male	1990	2,951	(1,817 to 4,582)	2225.2	(1398.3 to 3324.6)
			2013	5,738	(3,469 to 8,771)	2340.7	(1468.9 to 3462.0)
		Female	1990	2,679	(1,718 to 4,114)	2377.7	(1598.2 to 3601.7)
			2013	4,531	(2,234 to 7,498)	2422.5	(1208.1 to 3926.1)
		Both	1990	5,630	(3,666 to 8,562)	2298.5	(1546.0 to 3404.9)
			2013	10,269	(5,998 to 16,008)	2376.3	(1427.2 to 3611.5)
	YLDs	Male	1990	60	(39 to 85)	41.7	(26.7 to 59.1)
			2013	117	(76 to 165)	43.6	(28.3 to 61.5)
		Female	1990	45	(30 to 63)	37.3	(24.8 to 51.2)
			2013	76	(51 to 106)	36	(24.0 to 50.6)
		Both	1990	106	(71 to 143)	39.7	(26.9 to 53.1)
			2013	194	(131 to 259)	40.2	(27.2 to 54.4)
Bolivia	Deaths	Male	1990	1,106	(901 to 1,316)	82.9	(66.9 to 99.9)
			2013	1,617	(1,202 to 2,158)	68.4	(52.4 to 90.2)
		Female	1990	1,361	(1,101 to 1,676)	88	(70.4 to 108.7)
			2013	2,111	(1,497 to 2,921)	72.7	(51.5 to 99.6)

		Both	1990	2,467	(2,144 to 2,858)	85.8	(73.5 to 98.7)
			2013	3,728	(2,898 to 4,738)	70.9	(55.6 to 89.1)
	DALYs	Male	1990	36,062	(28,642 to 44,335)	1713.6	(1392.0 to 2039.3)
			2013	39,039	(28,290 to 52,831)	1224.5	(895.4 to 1652.8)
		Female	1990	40,425	(30,878 to 50,965)	1797.4	(1444.4 to 2231.6)
			2013	44,075	(30,170 to 61,486)	1247.4	(858.5 to 1746.9)
		Both	1990	76,487	(64,476 to 90,161)	1757.8	(1527.9 to 2038.7)
			2013	83,115	(64,107 to 106,070)	1238.8	(958.9 to 1576.6)
	YLDs	Male	1990	388	(260 to 547)	24.4	(16.5 to 34.3)
			2013	677	(455 to 924)	21.9	(14.8 to 30.0)
		Female	1990	220	(150 to 314)	11.9	(8.2 to 16.9)
			2013	407	(265 to 566)	11.4	(7.4 to 15.9)
		Both	1990	609	(425 to 837)	17.6	(12.2 to 24.3)
			2013	1,085	(740 to 1,447)	16.2	(11.2 to 22.0)
Bosnia and							
Herzegovina	Deaths	Male	1990	2,036	(1,752 to 2,454)	173.3	(148.7 to 208.0)
			2013	3,083	(2,353 to 3,734)	139	(106.7 to 169.4)
		Female	1990	2,441	(2,082 to 2,816)	142.4	(120.9 to 164.7)
			2013	3,476	(2,659 to 4,115)	111.3	(85.4 to 132.2)
		Both	1990	4,477	(4,103 to 5,087)	154.7	(141.4 to 175.4)
			2013	6,559	(5,582 to 7,500)	123.2	(104.4 to 141.4)
	DALYs	Male	1990	39,920	(34,432 to 48,328)	2717.8	(2350.2 to 3283.3)
			2013	47,315	(36,276 to 56,775)	1980.3	(1514.2 to 2378.3)
		Female	1990	39,719	(34,134 to 45,383)	2038.5	(1752.9 to 2338.4)
			2013	43,750	(33,630 to 50,855)	1394.7	(1076.1 to 1627.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	79,640	(73,188 to 90,444)	2322.2	(2133.6 to 2640.8)
			2013	91,065	(77,389 to 103,574)	1656.1	(1413.4 to 1880.8)

	1	1	ı ı				
	YLDs	Male	1990	462	(297 to 667)	25.4	(16.4 to 36.0)
			2013	742	(495 to 1,057)	30.7	(20.6 to 43.6)
		Female	1990	379	(255 to 530)	17.5	(11.8 to 24.3)
			2013	514	(334 to 741)	18	(11.8 to 25.7)
		Both	1990	841	(574 to 1,146)	21	(14.5 to 28.6)
			2013	1,256	(848 to 1,724)	23.9	(16.2 to 32.9)
Botswana	Deaths	Male	1990	146	(96 to 221)	99.3	(68.7 to 143.2)
			2013	264	(176 to 400)	86.5	(59.9 to 128.1)
		Female	1990	215	(141 to 321)	107.9	(73.7 to 152.5)
			2013	203	(109 to 451)	48.3	(26.6 to 105.5)
		Both	1990	361	(259 to 479)	105.2	(78.4 to 136.5)
			2013	468	(331 to 737)	63.4	(45.4 to 100.3)
	DALYs	Male	1990	3,576	(2,317 to 5,444)	1659.3	(1103.6 to 2510.4)
			2013	6,216	(4,100 to 9,345)	1416.2	(942.1 to 2171.7)
		Female	1990	4,211	(2,653 to 6,633)	1599.9	(1037.3 to 2412.7)
			2013	3,716	(2,017 to 7,907)	613.3	(324.0 to 1402.8)
		Both	1990	7,788	(5,334 to 10,669)	1634	(1174.1 to 2181.0)
			2013	9,933	(7,085 to 14,606)	946.3	(669.0 to 1482.9)
	YLDs	Male	1990	64	(42 to 88)	25	(16.8 to 34.7)
			2013	122	(78 to 170)	24.7	(15.9 to 34.5)
		Female	1990	71	(45 to 100)	22.2	(14.5 to 31.2)
			2013	131	(88 to 181)	22.3	(15.1 to 30.3)
		Both	1990	135	(90 to 182)	23.4	(16.0 to 31.5)
			2013	254	(172 to 343)	23.4	(16.0 to 31.6)
Brazil	Deaths	Male	1990	47,548	(40,965 to 55,332)	148.3	(127.3 to 173.0)
			2013	72,206	(57,918 to 83,513)	104.2	(83.3 to 121.2)
		Female	1990	47,039	(36,977 to 51,000)	127.1	(101.1 to 139.6)
			2013	71,564	(62,021 to 90,258)	79.5	(68.8 to 100.1)

		Both	1990	94,588	(82,676 to 102,963)	137.1	(119.5 to 149.7)
			2013	143,771	(129,078 to 162,465)	90.5	(81.0 to 102.6)
	DALYs	Male	1990	1,091,967	(942,110 to 1,273,336)	2718.9	(2355.6 to 3152.6)
					(1,131,935 to		
_			2013	1,410,563	1,651,312)	1786.8	(1445.1 to 2081.7)
		Female	1990	980,544		2167.8	(1686.5 to 2328.2)
					(1,014,386 to		
			2013	1,174,102	1,500,203)	1239.9	(1072.0 to 1580.2)
		5	1000	2 072 544	(1,815,328 to	2.426.2	(2422.4) 2642.0)
		Both	1990	2,072,511	· · · · · · · · · · · · · · · · · · ·	2426.3	(2132.4 to 2642.0)
			2013	2 504 666	(2,287,301 to	1488.5	/1227 0 to 1602 E\
	VIDa	Mala		2,584,666			(1327.8 to 1683.5)
	YLDs	Male	1990	16,997	(10,784 to 24,536)	42.6	(27.0 to 61.5) Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Courtery	measare	JEX	2013	32,349	(20,843 to 45,917)	40.1	(25.6 to 56.6)
		Female	1990	12,553	(8,400 to 17,384)	27.9	(18.6 to 39.2)
			2013	25,107	(16,578 to 34,604)	26.3	(17.4 to 36.5)
		_					
L		Both	1990	29,551	(19,595 to 40,263)	34.7	(22.9 to 47.5)
		Both	1990 2013	29,551 57,457	(19,595 to 40,263) (39,551 to 76,339)	34.7 32.6	(22.9 to 47.5) (22.4 to 43.3)
Brunei	Deaths	Both Male					•
Brunei	Deaths		2013	57,457	(39,551 to 76,339)	32.6	(22.4 to 43.3)
Brunei	Deaths		2013 1990	57,457 60	(39,551 to 76,339) (50 to 70)	32.6 149.2	(22.4 to 43.3) (126.3 to 177.2)
Brunei	Deaths	Male	2013 1990 2013	57,457 60 84	(39,551 to 76,339) (50 to 70) (62 to 108)	32.6 149.2 90.5	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7)
Brunei	Deaths	Male	2013 1990 2013 1990	57,457 60 84 57	(39,551 to 76,339) (50 to 70) (62 to 108) (43 to 67)	32.6 149.2 90.5 158.3	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7) (121.0 to 183.6)
Brunei	Deaths	Male Female	2013 1990 2013 1990 2013	57,457 60 84 57 87	(39,551 to 76,339) (50 to 70) (62 to 108) (43 to 67) (67 to 110)	32.6 149.2 90.5 158.3 89.8	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7) (121.0 to 183.6) (70.9 to 114.9)
Brunei	Deaths	Male Female	2013 1990 2013 1990 2013 1990	57,457 60 84 57 87 118	(39,551 to 76,339) (50 to 70) (62 to 108) (43 to 67) (67 to 110) (102 to 130)	32.6 149.2 90.5 158.3 89.8 153.8	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7) (121.0 to 183.6) (70.9 to 114.9) (133.6 to 171.5)
Brunei		Male Female Both	2013 1990 2013 1990 2013 1990 2013	57,457 60 84 57 87 118 171	(39,551 to 76,339) (50 to 70) (62 to 108) (43 to 67) (67 to 110) (102 to 130) (143 to 204)	32.6 149.2 90.5 158.3 89.8 153.8 90.5	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7) (121.0 to 183.6) (70.9 to 114.9) (133.6 to 171.5) (76.8 to 108.4)
Brunei		Male Female Both	2013 1990 2013 1990 2013 1990 2013 1990	57,457 60 84 57 87 118 171 1,385	(39,551 to 76,339) (50 to 70) (62 to 108) (43 to 67) (67 to 110) (102 to 130) (143 to 204) (1,169 to 1,606)	32.6 149.2 90.5 158.3 89.8 153.8 90.5 2608.7	(22.4 to 43.3) (126.3 to 177.2) (69.3 to 115.7) (121.0 to 183.6) (70.9 to 114.9) (133.6 to 171.5) (76.8 to 108.4) (2218.4 to 3039.4)

		Both	1990	2,587	(2,233 to 2,902)	2570.8	(2249.5 to 2857.7)
			2013	3,686	(3,049 to 4,407)	1368.1	(1149.4 to 1624.0)
	YLDs	Male	1990	65	(43 to 91)	132	(87.8 to 187.1)
			2013	165	(110 to 239)	123.7	(81.3 to 177.1)
		Female	1990	35	(23 to 48)	77.7	(52.9 to 106.1)
			2013	107	(73 to 148)	81.6	(54.5 to 111.9)
		Both	1990	100	(69 to 137)	106	(73.5 to 145.8)
			2013	272	(186 to 371)	102.4	(70.3 to 140.9)
Bulgaria	Deaths	Male	1990	12,461	(10,853 to 14,204)	271.7	(236.4 to 310.2)
			2013	12,232	(10,598 to 13,906)	224.7	(194.1 to 256.0)
		Female	1990	14,106	(12,007 to 15,283)	237.2	(202.3 to 258.5)
			2013	14,014	(12,356 to 16,739)	162.3	(143.9 to 195.1)
		Both	1990	26,567	(24,012 to 28,557)	253.3	(228.7 to 272.9)
			2013	26,247	(23,810 to 29,709)	189.2	(171.7 to 214.4)
	DALYs	Male	1990	215,900	(188,289 to 246,174)	4333.6	(3788.2 to 4953.7)
			2013	187,795	(162,812 to 214,727)	3474.6	(3005.5 to 3974.8)
		Female	1990	205,673	(172,014 to 220,136)	3388.5	(2825.6 to 3628.6)
			2013	164,752	(147,021 to 200,607)	2098	(1871.7 to 2570.3)
		Both	1990	421,574	(380,573 to 453,732)	3834.1	(3466.0 to 4124.0)
			2013	352,548	(319,254 to 400,227)	2716.3	(2454.1 to 3097.1)
	YLDs	Male	1990	1,423	(915 to 2,015)	27.8	(18.2 to 39.5)
			2013	1,570	(1,045 to 2,249)	30.4	(20.4 to 43.8)
		Female	1990	1,072	(709 to 1,486)	18.3	(12.2 to 25.3)
			2013	1,176	(778 to 1,683)	18.3	(12.3 to 26.0)
		Both	1990	2,496	(1,706 to 3,449)	22.8	(15.6 to 31.2)
			2013	2,747	(1,881 to 3,776)	23.9	(16.3 to 32.7)
Burkina Faso	Deaths	Male	1990	791	(587 to 971)	59.3	(43.6 to 73.5)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval

			2013	1,861	(1,406 to 2,336)	103.5	(77.8 to 128.4)
		Female	1990	966	(734 to 1,305)	56	(42.1 to 76.3)
			2013	2,154	(1,539 to 2,859)	87.2	(60.2 to 116.6)
_		Both	1990	1,757	(1,477 to 2,085)	57.4	(48.0 to 68.6)
			2013	4,015	(3,162 to 4,834)	93.9	(73.5 to 113.3)
	DALYs	Male	1990	21,359	(15,935 to 26,294)	1204.5	(903.0 to 1470.1)
			2013	46,403	(34,809 to 59,272)	1854.6	(1408.1 to 2322.0)
		Female	1990	24,435	(18,485 to 31,999)	1093.4	(837.9 to 1461.6)
			2013	47,660	(34,260 to 64,676)	1459.3	(1041.8 to 1936.0)
		Both	1990	45,794	(38,313 to 53,865)	1140.8	(954.5 to 1348.4)
			2013	94,063	(73,432 to 113,989)	1623.6	(1279.7 to 1949.5)
	YLDs	Male	1990	351	(235 to 492)	20.1	(13.3 to 27.9)
			2013	595	(396 to 846)	18.8	(12.3 to 26.8)
		Female	1990	321	(207 to 458)	14.2	(9.2 to 20.4)
			2013	568	(376 to 816)	14.5	(9.6 to 21.2)
		Both	1990	673	(454 to 928)	16.8	(11.5 to 22.7)
			2013	1,164	(800 to 1,612)	16.3	(11.3 to 22.3)
Burundi	Deaths	Male	1990	1,874	(1,260 to 2,615)	239.4	(162.6 to 321.6)
			2013	2,253	(1,653 to 2,862)	172.2	(128.0 to 217.6)
		Female	1990	2,442	(1,559 to 3,567)	249.8	(162.4 to 364.7)
			2013	2,614	(1,963 to 3,636)	186.2	(138.9 to 254.4)
		Both	1990	4,317	(3,274 to 5,780)	245.7	(185.6 to 323.3)
			2013	4,868	(4,070 to 5,982)	180.3	(150.4 to 220.3)
	DALYs	Male	1990	45,641	(30,977 to 64,638)	4542.6	(3048.8 to 6342.4)
			2013	53,044	(38,141 to 68,470)	3032.2	(2227.3 to 3855.8)
		Female	1990	54,105	(32,737 to 80,640)	4568.8	(2873.6 to 6730.4)
			2013	53,271	(39,958 to 75,617)	3073.6	(2318.9 to 4280.7)
		Both	1990	99,747	(74,374 to 134,733)	4565.1	(3443.5 to 6101.4)

			2013	106,315	(87,306 to 131,464)	3063.9	(2552.1 to 3769.5)
	YLDs	Male	1990	163	(107 to 240)	13.6	(9.0 to 19.9)
			2013	297	(188 to 433)	13.2	(8.3 to 19.0)
		Female	1990	123	(80 to 178)	8.9	(5.8 to 12.9)
			2013	221	(142 to 317)	9.4	(6.1 to 13.2)
		Both	1990	287	(196 to 395)	11	(7.6 to 15.4)
			2013	518	(348 to 714)	11.2	(7.5 to 15.3)
Cambodia	Deaths	Male	1990	2,961	(2,416 to 3,505)	213.6	(174.8 to 252.2)
			2013	7,153	(5,757 to 8,890)	188.9	(152.6 to 234.8)
		Female	1990	3,137	(2,311 to 3,919)	168.7	(128.7 to 212.2)
			2013	9,442	(7,449 to 11,656)	183	(144.7 to 226.0)
		Both	1990	6,098	(5,113 to 7,167)	187.8	(158.5 to 221.5)
			2013	16,596	(14,179 to 19,460)	186.1	(158.5 to 217.6)
	DALYs	Male	1990	90,165	(72,812 to 109,847)	4279.3	(3489.3 to 5052.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	156,913	(121,054 to 197,403)	3544.5	(2795.1 to 4439.2)
		Female	1990		(50 400 + 00 405)		
			1990	74,497	(50,492 to 93,485)	3070.9	(2208.5 to 3850.2)
			2013	74,497 165,311	· · · · · · · · · · · · · · · · · · ·	3070.9 3005.3	(2208.5 to 3850.2) (2320.4 to 3792.6)
		Both			· · · · · · · · · · · · · · · · · · ·		· `
		Both	2013	165,311	(126,631 to 207,488) (131,078 to 194,954)	3005.3	(2320.4 to 3792.6)
	YLDs	Both Male	2013 1990	165,311 164,663	(126,631 to 207,488) (131,078 to 194,954)	3005.3 3614.5	(2320.4 to 3792.6) (3031.2 to 4243.3)
	YLDs		2013 1990 2013	165,311 164,663 322,225	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018)	3005.3 3614.5 3250.8	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4)
	YLDs		2013 1990 2013 1990	165,311 164,663 322,225 772	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018) (524 to 1,059)	3005.3 3614.5 3250.8 47.8	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4) (32.2 to 65.8)
	YLDs	Male	2013 1990 2013 1990 2013	165,311 164,663 322,225 772 2,226	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018) (524 to 1,059) (1,492 to 3,061)	3005.3 3614.5 3250.8 47.8 55.3	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4) (32.2 to 65.8) (37.3 to 77.2)
	YLDs	Male	2013 1990 2013 1990 2013 1990	165,311 164,663 322,225 772 2,226 840	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018) (524 to 1,059) (1,492 to 3,061) (558 to 1,174)	3005.3 3614.5 3250.8 47.8 55.3 39.9	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4) (32.2 to 65.8) (37.3 to 77.2) (25.8 to 56.0)
	YLDs	Male Female	2013 1990 2013 1990 2013 1990 2013	165,311 164,663 322,225 772 2,226 840 2,377	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018) (524 to 1,059) (1,492 to 3,061) (558 to 1,174) (1,569 to 3,289)	3005.3 3614.5 3250.8 47.8 55.3 39.9 44.2	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4) (32.2 to 65.8) (37.3 to 77.2) (25.8 to 56.0) (28.8 to 61.5)
Cameroon	YLDs	Male Female	2013 1990 2013 1990 2013 1990 2013 1990	165,311 164,663 322,225 772 2,226 840 2,377 1,613	(126,631 to 207,488) (131,078 to 194,954) (271,887 to 385,018) (524 to 1,059) (1,492 to 3,061) (558 to 1,174) (1,569 to 3,289) (1,141 to 2,159)	3005.3 3614.5 3250.8 47.8 55.3 39.9 44.2 43.1	(2320.4 to 3792.6) (3031.2 to 4243.3) (2752.9 to 3854.4) (32.2 to 65.8) (37.3 to 77.2) (25.8 to 56.0) (28.8 to 61.5) (30.3 to 57.9)

		Female	1990	3,738	(2,982 to 4,898)	161	(126.3 to 212.0)
			2013	5,870	(4,114 to 8,601)	153.4	(108.9 to 219.0)
		Both	1990	6,471	(5,505 to 7,652)	149.5	(126.7 to 177.8)
			2013	12,032	(9,982 to 15,108)	164.7	(136.6 to 206.7)
	DALYs	Male	1990	71,063	(58,208 to 84,950)	2456.1	(2009.7 to 2935.4)
			2013	153,178	(117,089 to 190,089)	3209.2	(2499.1 to 4005.7)
		Female	1990	85,019	(68,436 to 112,997)	2780.1	(2220.5 to 3645.4)
			2013	123,381	(83,026 to 193,776)	2493.8	(1723.3 to 3674.5)
		Both	1990	156,083	(133,227 to 184,027)	2636.4	(2250.8 to 3116.8)
			2013	276,560	(226,438 to 356,603)	2839.3	(2340.8 to 3571.8)
	YLDs	Male	1990	549	(365 to 769)	19.6	(13.1 to 27.5)
			2013	988	(663 to 1,350)	20	(13.7 to 27.2)
		Female	1990	436	(285 to 608)	14	(9.1 to 19.7)
			2013	766	(498 to 1,066)	14.2	(9.3 to 19.9)
		Both	1990	985	(670 to 1,335)	16.6	(11.4 to 22.6)
			2013	1,755	(1,200 to 2,367)	16.9	(11.6 to 22.6)
Canada	Deaths	Male	1990	6,364	(5,180 to 7,843)	54.7	(44.3 to 67.7)
			2013	6,720	(4,714 to 8,383)	27.2	(19.1 to 33.9)
		Female	1990	8,588	(6,980 to 10,634)	47.6	(38.7 to 58.9)
			2013	10,807	(8,165 to 13,992)	29.7	(22.5 to 38.3)
		Both	1990	14,952	(13,190 to 17,597)	50.8	(44.7 to 59.7)
			2013	17,528	(14,128 to 20,918)	28.9	(23.3 to 34.1)
	DALYs	Male	1990	115,732	(97,264 to 138,906)	907.6	(761.3 to 1094.0)
			2013	112,987	(87,754 to 135,297)	471.6	(368.2 to 565.0)
		Female	1990	117,395	(102,853 to 143,563)	694	(607.5 to 846.3)
			2013	129,962	(108,137 to 158,635)	433.5	(357.9 to 526.9)
		Both	1990	233,128	(208,580 to 267,210)	786.5	(703.5 to 899.9)
			2013	242,949	(203,364 to 280,805)	451.9	(378.5 to 525.2)

		_					
	YLDs	Male	1990	19,094	(12,581 to 26,386)	148.1	(97.1 to 203.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	33,135	(22,215 to 46,569)	138.5	(92.8 to 194.4)
		Female	1990	18,757	(11,836 to 27,364)	112.6	(70.9 to 165.3)
			2013	33,604	(21,717 to 47,935)	118.6	(75.6 to 168.3)
		Both	1990	37,851	(26,649 to 53,016)	127.7	(90.0 to 178.4)
			2013	66,740	(45,097 to 91,295)	127.5	(86.0 to 174.0)
Cape Verde	Deaths	Male	1990	161	(141 to 185)	216	(189.1 to 247.6)
			2013	252	(177 to 349)	206.1	(145.9 to 282.1)
		Female	1990	175	(152 to 208)	186.7	(162.7 to 219.6)
			2013	172	(97 to 362)	89.1	(49.7 to 189.1)
		Both	1990	336	(306 to 379)	199.5	(181.8 to 223.9)
			2013	425	(298 to 673)	136.7	(95.3 to 216.6)
	DALYs	Male	1990	3,592	(3,084 to 4,241)	3985.2	(3481.5 to 4624.4)
			2013	5,014	(3,381 to 7,132)	3561.7	(2400.1 to 5100.6)
		Female	1990	2,780	(2,403 to 3,514)	2563.1	(2238.1 to 3190.3)
			2013	1,986	(1,059 to 4,560)	1075.2	(558.1 to 2496.0)
		Both	1990	6,373	(5,704 to 7,438)	3168.8	(2866.5 to 3661.0)
			2013	7,001	(4,684 to 10,986)	2147.1	(1427.8 to 3399.5)
	YLDs	Male	1990	16	(11 to 23)	19.9	(13.3 to 28.5)
			2013	28	(18 to 40)	19.2	(12.5 to 27.1)
		Female	1990	15	(10 to 21)	14.2	(9.5 to 19.6)
			2013	27	(17 to 38)	14.7	(9.5 to 20.4)
		Both	1990	32	(21 to 43)	16.7	(11.4 to 22.7)
			2013	55	(37 to 76)	16.6	(11.2 to 22.7)
Central African				_			
Republic	Deaths	Male	1990	970	(713 to 1,292)	171.4	(128.0 to 224.1)
			2013	787	(531 to 1,109)	99.5	(66.5 to 138.8)

		Female	1990	1,519	(1,005 to 1,890)	233.6	(155.2 to 292.4)
			2013	1,997	(961 to 2,885)	201.9	(98.1 to 286.4)
		Both	1990	2,490	(1,866 to 3,067)	207.3	(155.8 to 255.1)
			2013	2,785	(1,671 to 3,706)	158.8	(94.7 to 210.2)
	DALYs	Male	1990	24,912	(18,022 to 33,885)	3417.6	(2498.9 to 4603.3)
			2013	18,718	(12,674 to 26,324)	1822.5	(1231.1 to 2588.0)
		Female	1990	33,877	(22,102 to 42,530)	4191.8	(2751.6 to 5238.0)
			2013	41,917	(19,999 to 62,737)	3460.6	(1658.0 to 5075.0)
		Both	1990	58,789	(43,217 to 73,623)	3854.8	(2859.8 to 4791.6)
			2013	60,635	(36,749 to 81,849)	2740.1	(1646.5 to 3655.0)
	YLDs	Male	1990	136	(93 to 188)	18.5	(12.6 to 25.5)
			2013	198	(124 to 277)	17.6	(11.2 to 24.2)
		Female	1990	115	(75 to 160)	13.5	(8.9 to 18.9)
			2013	179	(116 to 250)	13.9	(8.9 to 19.6)
		Both	1990	251	(175 to 342)	15.8	(11.1 to 21.3)
			2013	377	(255 to 513)	15.6	(10.5 to 21.3)
Chad	Deaths	Male	1990	748	(572 to 944)	80	(60.4 to 101.4)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	1,394	(983 to 1,929)	87.9	(61.7 to 119.7)
		Female	1990	1,093	(763 to 1,358)	102.9	(72.8 to 129.6)
			2013	1,946	(1,251 to 2,794)	112	(72.7 to 158.3)
		Both	1990	1,841	(1,456 to 2,221)	93.1	(73.0 to 112.8)
			2013	3,341	(2,480 to 4,340)	101.5	(74.9 to 131.8)
	DALYs	Male	1990	21,223	(16,307 to 27,010)	1481.7	(1134.8 to 1864.4)
			2013	37,478	(26,421 to 52,347)	1577	(1113.8 to 2189.1)
		Female	1990	26,763	(18,583 to 33,655)	1769.1	(1226.6 to 2201.7)
			2013	45,448	(29,800 to 66,017)	1846	(1170.8 to 2664.6)

			2013	82,927	(62,172 to 108,320)	1725.3	(1278.9 to 2259.7)
	YLDs	Male	1990	251	(161 to 351)	19.3	(12.5 to 27.0)
			2013	464	(304 to 648)	19.1	(12.4 to 27.5)
		Female	1990	198	(128 to 281)	13.7	(8.8 to 19.7)
			2013	356	(240 to 488)	13.3	(8.9 to 18.2)
		Both	1990	450	(307 to 619)	16.3	(11.1 to 22.4)
			2013	821	(563 to 1,113)	16	(11.1 to 21.9)
Chile	Deaths	Male	1990	3,879	(3,344 to 4,739)	118.7	(101.5 to 146.6)
			2013	5,454	(4,444 to 6,415)	72.9	(59.0 to 85.9)
		Female	1990	4,296	(3,566 to 5,164)	91.1	(75.4 to 109.6)
			2013	5,778	(4,809 to 7,635)	53.4	(44.6 to 70.7)
		Both	1990	8,175	(7,390 to 9,362)	102.7	(92.7 to 118.0)
			2013	11,233	(9,745 to 13,062)	62	(53.7 to 72.1)
	DALYs	Male	1990	75,278	(65,031 to 90,302)	1965.2	(1706.3 to 2385.3)
			2013	92,460	(76,394 to 107,372)	1125.5	(931.2 to 1306.6)
		Female	1990	68,995	(57,838 to 81,785)	1384.7	(1157.9 to 1638.1)
			2013	74,367	(62,664 to 96,860)	725.7	(613.5 to 944.2)
		Both	1990	144,274	(132,281 to 164,509)	1634.5	(1496.3 to 1864.8)
			2013	166,827	(146,818 to 190,513)	906.6	(797.2 to 1033.2)
	YLDs	Male	1990	1,879	(1,273 to 2,513)	45.5	(31.2 to 60.7)
			2013	3,715	(2,479 to 4,999)	43.5	(29.1 to 59.2)
		Female	1990	1,315	(869 to 1,843)	25.6	(16.9 to 35.9)
			2013	2,473	(1,604 to 3,470)	24.7	(16.1 to 34.5)
		Both	1990	3,195	(2,173 to 4,247)	34.3	(23.3 to 46.0)
			2013	6,189	(4,227 to 8,161)	33.3	(22.7 to 44.1)
China	Deaths	Male	1990	702,451	(601,130 to 780,076)	233.5	(201.4 to 258.2)
			2013	1,110,238	(928,780 to 1,356,982)	188.9	(159.3 to 231.0)
		Female	1990	603,603	(533,293 to 728,820)	171.6	(151.8 to 207.0)

	1	1			<u></u>	1	
			2013	810,449	(692,117 to 924,961)	127.5	(109.3 to 145.9)
					(1,191,710 to		
		Both	1990	1,306,054	1,436,929)	199.7	(182.7 to 219.6)
					(1,715,665 to		
			2013	1,920,688	2,196,206)	157.3	(141.9 to 180.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
					(13,579,956 to		
	DALYs	Male	1990	15,810,086	17,802,632)	4149	(3572.2 to 4615.2)
					(17,184,678 to		
			2013	20,778,030	25,034,999)	3032.6	(2520.0 to 3665.7)
					(10,042,789 to		
		Female	1990	11,510,587	14,058,890)	2913.5	(2561.1 to 3527.9)
					(10,010,331 to		
			2013	11,917,010	13,848,970)	1725	(1457.2 to 1990.0)
					(24,692,698 to		(
		Both	1990	27,320,673	30,335,280)	3508.5	(3191.6 to 3863.1)
			2010		(28,591,270 to	2071 -	(2006 4) 2744 6)
			2013	32,695,040	37,613,155)	2371.7	(2086.1 to 2711.6)
	YLDs	Male	1990	219,111	(154,730 to 284,153)	62.5	(44.2 to 81.2)
			2013	429,633	(304,101 to 554,874)	63.7	(45.3 to 82.1)
		Female	1990	154,071	(110,120 to 199,469)	39.9	(28.6 to 51.5)
			2013	277,731	(195,371 to 361,615)	39.5	(27.9 to 51.3)
		Both	1990	373,182	(265,035 to 482,793)	50.2	(35.8 to 64.9)
			2013	707,364	(500,459 to 917,801)	51.2	(36.2 to 66.3)
Colombia	Deaths	Male	1990	6,174	(5,249 to 7,343)	97.1	(82.2 to 115.8)
			2013	9,759	(7,653 to 12,326)	74.1	(58.4 to 93.6)
		Female	1990	7,203	(5,965 to 8,350)	91.1	(75.0 to 105.3)
			2013	11,013	(8,829 to 14,113)	63.9	(51.0 to 81.5)
		Both	1990	13,378	(12,045 to 14,943)	93.9	(83.9 to 105.2)
			2013	20,772	(17,642 to 24,710)	68.5	(58.4 to 81.4)

	DALYs	Male	1990	141,161	(120,617 to 168,007)	1719.5	(1467.9 to 2053.7)
			2013	189,739	(147,680 to 241,985)	1179.4	(927.8 to 1494.7)
		Female	1990	149,913	(123,858 to 172,483)	1558.4	(1288.0 to 1795.7)
			2013	180,681	(142,046 to 235,113)	932.3	(736.2 to 1201.9)
		Both	1990	291,075	(261,360 to 325,824)	1630.5	(1473.1 to 1824.1)
			2013	370,420	(310,561 to 441,831)	1043.5	(882.4 to 1245.6)
	YLDs	Male	1990	2,148	(1,458 to 2,918)	26.2	(17.9 to 35.8)
			2013	4,726	(3,156 to 6,646)	27.6	(18.7 to 38.5)
		Female	1990	1,726	(1,178 to 2,331)	18.3	(12.5 to 24.7)
			2013	3,744	(2,434 to 5,188)	18.4	(12.0 to 25.6)
		Both	1990	3,874	(2,661 to 5,093)	21.9	(15.1 to 28.9)
			2013	8,471	(5,832 to 11,629)	22.5	(15.5 to 30.5)
Comoros	Deaths	Male	1990	154	(99 to 230)	251.9	(171.2 to 362.2)
			2013	248	(162 to 345)	230.5	(160.5 to 304.7)
		Female	1990	187	(114 to 273)	259.7	(165.5 to 365.7)
			2013	248	(166 to 354)	210.3	(142.7 to 294.3)
		Both	1990	342	(240 to 457)	257.3	(187.5 to 335.2)
			2013	496	(371 to 635)	220.4	(168.4 to 277.2)
	DALYs	Male	1990	3,783	(2,416 to 5,768)	4637.6	(2989.8 to 6936.4)
			2013	5,978	(3,716 to 8,640)	4136.1	(2705.4 to 5747.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	3,990	(2,351 to 6,040)	4472.9	(2710.0 to 6620.8)
			2013	5,020	(3,161 to 7,438)	3428.8	(2261.5 to 4925.4)
		Both	1990	7,774	(5,310 to 10,586)	4567.3	(3190.2 to 6149.8)
			2013	10,999	(7,953 to 14,486)	3776.6	(2808.5 to 4864.2)
	YLDs	Male	1990	12	(7 to 18)	12.8	(8.0 to 18.5)
			2013	25	(15 to 36)	13.8	(9.2 to 20.1)
		Female	1990	9	(5 to 13)	8.8	(5.7 to 12.7)

	DALYs	Male	1990	7,344	(6,264 to 8,926)	877.7	(741.7 to 1068.0)
			2013	1,483	(1,286 to 1,744)	40.9	(35.4 to 48.2)
		Both	1990	897	(795 to 1,022)	60.2	(53.1 to 69.2)
			2013	776	(622 to 1,009)	39.1	(31.2 to 50.7)
		Female	1990	483	(395 to 575)	61	(49.8 to 73.1)
			2013	707	(556 to 858)	42.8	(33.3 to 52.0)
Costa Rica	Deaths	Male	1990	413	(343 to 504)	59.1	(48.8 to 72.6)
			2013	344	(238 to 472)	15.4	(10.6 to 21.1)
		Both	1990	194	(133 to 260)	16.1	(11.0 to 21.7)
			2013	156	(104 to 223)	13.6	(8.9 to 19.4)
		Female	1990	87	(57 to 123)	13.9	(9.1 to 19.3)
			2013	188	(127 to 269)	17.5	(11.6 to 24.9)
	YLDs	Male	1990	106	(69 to 146)	18.5	(12.1 to 25.8)
			2013	85,577	(68,931 to 103,445)	4396.6	(3583.1 to 5283.2)
		Both	1990	61,441	(51,814 to 71,813)	5559.3	(4763.7 to 6446.6)
			2013	38,462	(28,014 to 54,307)	3955.8	(2922.8 to 5417.7)
		Female	1990	30,557	(22,408 to 38,071)	5370.7	(4120.5 to 6630.9)
			2013	47,115	(34,543 to 60,346)	4843.1	(3602.4 to 6081.5)
	DALYs	Male	1990	30,883	(25,113 to 37,473)	5731.3	(4695.5 to 6880.7)
			2013	3,927	(3,209 to 4,735)	254.2	(207.9 to 304.6)
		Both	1990	2,751	(2,367 to 3,163)	306	(263.9 to 349.8)
			2013	1,962	(1,464 to 2,641)	245.5	(184.4 to 325.2)
		Female	1990	1,467	(1,150 to 1,785)	310	(248.9 to 376.0)
			2013	1,964	(1,473 to 2,457)	261.6	(200.8 to 322.3)
Congo	Deaths	Male	1990	1,284	(1,063 to 1,527)	298.6	(247.1 to 349.2)
			2013	43	(27 to 61)	11.5	(7.6 to 16.4)
		Both	1990	21	(14 to 29)	10.7	(7.2 to 14.7)
			2013	18	(11 to 27)	9.5	(6.1 to 14.0)

			2013	11,449	(9,267 to 13,693)	605.3	(487.4 to 721.1)
		Female	1990	7,599	(6,120 to 8,860)	857.9	(694.7 to 1003.9)
			2013	10,158	(8,443 to 12,864)	491.7	(408.0 to 621.3)
		Both	1990	14,943	(13,232 to 16,900)	867.8	(773.7 to 982.0)
			2013	21,608	(18,925 to 24,939)	546.3	(478.7 to 630.3)
	YLDs	Male	1990	240	(162 to 330)	27.4	(18.1 to 37.7)
			2013	565	(375 to 788)	27.8	(18.4 to 38.4)
Country					Unanatainto latemal	Data	Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	167	(111 to 230)	18.5	(12.3 to 25.3)
			2013	395	(259 to 555)	18.6	(12.1 to 26.5)
		Both	1990	408	(279 to 549)	22.8	(15.5 to 30.7)
			2013	960	(667 to 1,265)	23.1	(16.0 to 30.3)
Cote d'Ivoire	Deaths	Male	1990	3,636	(3,055 to 4,313)	169.9	(142.6 to 198.6)
			2013	6,660	(5,321 to 8,549)	181.7	(147.0 to 226.1)
		Female	1990	2,672	(1,874 to 3,501)	158.1	(112.6 to 209.9)
			2013	4,986	(3,720 to 6,950)	177.3	(135.7 to 245.9)
		Both	1990	6,308	(5,385 to 7,450)	165	(141.5 to 194.3)
			2013	11,646	(9,791 to 14,274)	179.9	(152.6 to 217.8)
	DALYs	Male	1990	104,862	(86,709 to 125,638)	3290.7	(2763.4 to 3908.2)
			2013	171,309	(135,603 to 223,340)	3354.6	(2690.8 to 4306.8)
		Female	1990	68,800	(45,589 to 90,167)	2721.1	(1918.7 to 3566.1)
			2013	118,157	(86,757 to 169,363)	2918.7	(2181.4 to 4062.9)
		Both	1990	173,662	(146,841 to 205,047)	3037	(2599.5 to 3573.9)
			2013	289,466	(242,633 to 355,879)	3153.2	(2650.7 to 3862.8)
	YLDs	Male	1990	617	(402 to 861)	19.6	(13.1 to 27.5)
			2013	1,010	(647 to 1,470)	18.6	(11.7 to 27.5)
		Female	1990	377	(247 to 519)	14.5	(9.8 to 19.7)
			2013	680	(445 to 958)	14.3	(9.4 to 20.3)

		Both	1990	995	(665 to 1,355)	17.3	(11.8 to 23.3)
			2013	1,690	(1,135 to 2,336)	16.6	(11.0 to 23.0)
Croatia	Deaths	Male	1990	4,105	(3,605 to 4,740)	206.8	(181.0 to 239.2)
			2013	3,584	(3,022 to 4,129)	119.9	(100.6 to 138.7)
		Female	1990	5,488	(4,533 to 6,261)	160.3	(132.7 to 183.2)
			2013	4,874	(4,177 to 5,898)	90.6	(78.1 to 110.0)
		Both	1990	9,594	(8,569 to 10,554)	178.5	(159.5 to 197.0)
			2013	8,458	(7,592 to 9,481)	103	(92.8 to 115.5)
	DALYs	Male	1990	73,704	(64,134 to 85,154)	3281.8	(2869.7 to 3787.2)
			2013	52,990	(44,466 to 61,104)	1743.6	(1465.5 to 2011.0)
		Female	1990	76,323	(63,254 to 85,234)	2253.9	(1871.4 to 2525.8)
			2013	51,196	(44,775 to 62,399)	1092.8	(957.5 to 1348.8)
		Both	1990	150,027	(135,092 to 163,488)	2682.6	(2416.1 to 2923.8)
			2013	104,186	(94,655 to 116,843)	1376.6	(1253.0 to 1546.6)
	YLDs	Male	1990	756	(498 to 1,106)	30.7	(20.3 to 44.4)
			2013	921	(625 to 1,314)	31.4	(21.4 to 44.6)
		Female	1990	635	(407 to 904)	19.6	(12.8 to 27.5)
			2013	733	(476 to 1,047)	19.3	(12.4 to 27.3)
		Both	1990	1,392	(939 to 1,922)	24.4	(16.5 to 33.7)
			2013	1,655	(1,135 to 2,262)	24.9	(17.1 to 33.8)
Cuba	Deaths	Male	1990	4,288	(3,601 to 5,207)	96.6	(81.1 to 117.1)
			2013	5,383	(4,320 to 6,535)	77.5	(62.1 to 94.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	4,810	(4,164 to 5,946)	96.4	(83.4 to 119.1)
			2013	6,029	(4,363 to 7,160)	69.9	(50.6 to 82.8)
		Both	1990	9,098	(8,191 to 10,628)	96.5	(86.9 to 112.8)
			2013	11,413	(9,407 to 13,036)	73.8	(61.0 to 84.2)
	DALYs	Male	1990	73,700	(62,947 to 89,232)	1636.6	(1398.0 to 1976.2)

			2013	86,053	(70,665 to 103,639)	1214.8	(997.9 to 1461.9)
		Female	1990	74,808	(64,821 to 90,810)	1541.6	(1335.0 to 1868.3)
			2013	76,606	· · · · · · · · · · · · · · · · · · ·	951.3	(711.6 to 1112.4)
		Both	1990	148,509	(135,422 to 170,210)	1587.5	(1449.0 to 1821.2)
			2013	162,660	(138,275 to 184,124)	1080.2	(919.1 to 1226.6)
	YLDs	Male	1990	2,647	(1,709 to 3,712)	59.2	(38.0 to 83.1)
			2013	3,960	(2,669 to 5,574)	56.4	(38.1 to 79.3)
		Female	1990	2,028	(1,350 to 2,816)	41.5	(27.8 to 57.4)
			2013	3,417	(2,262 to 4,792)	43.1	(28.8 to 60.1)
		Both	1990	4,675	(3,167 to 6,295)	50	(33.9 to 67.4)
			2013	7,377	(5,060 to 10,082)	49.5	(34.0 to 67.6)
Cyprus	Deaths	Male	1990	351	(245 to 430)	114.8	(80.2 to 140.4)
			2013	240	(193 to 299)	43.8	(35.4 to 54.5)
		Female	1990	420	(252 to 508)	109	(66.1 to 131.3)
			2013	320	(265 to 432)	41.8	(35.0 to 55.9)
		Both	1990	772	(538 to 903)	112.1	(78.7 to 130.8)
			2013	560	(486 to 690)	43	(37.4 to 52.7)
	DALYs	Male	1990	5,373	(3,917 to 6,352)	1748	(1275.8 to 2065.4)
			2013	3,569	(2,987 to 4,338)	676.5	(566.6 to 818.7)
		Female	1990	5,146	(3,440 to 5,976)	1434.3	(968.8 to 1658.4)
			2013	3,386	(2,871 to 4,354)	518.6	(440.0 to 669.6)
		Both	1990	10,519	(7,797 to 11,947)	1583	(1176.3 to 1794.7)
			2013	6,955	(6,136 to 8,110)	593.4	(524.6 to 690.4)
	YLDs	Male	1990	331	(224 to 451)	108.1	(73.2 to 147.4)
			2013	575	(374 to 802)	106.8	(69.5 to 148.9)
		Female	1990	248	(164 to 340)	70.7	(47.2 to 96.5)
			2013	461	(304 to 663)	73	(48.0 to 104.8)
		Both	1990	580	(402 to 769)	88.1	(61.2 to 116.8)

			2013	1,036	(722 to 1,397)	88.8	(62.1 to 119.1)
Czech Republic	Deaths	Male	1990	11,062	(9,724 to 12,982)	255.4	(224.9 to 296.2)
			2013	7,443	(6,370 to 8,728)	112.3	(96.3 to 131.7)
		Female	1990	15,755	(12,751 to 17,085)	196.9	(158.6 to 213.1)
			2013	10,038	(8,622 to 13,775)	86	(74.2 to 117.7)
		Both	1990	26,818	(23,824 to 28,977)	219.6	(195.5 to 237.0)
			2013	17,481	(15,530 to 21,570)	97	(86.3 to 119.2)
	DALYs	Male	1990	176,886	(153,703 to 210,305)	3714	(3267.5 to 4372.5)
			2013	101,687	(85,889 to 118,132)	1462.4	(1235.5 to 1695.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	195,251	(154,449 to 209,743)	2540.1	(1991.3 to 2725.0)
			2013	99,531	(86,158 to 135,408)	946.4	(825.9 to 1280.2)
		Both	1990	372,138	(330,730 to 404,343)	3022.5	(2692.0 to 3284.6)
			2013	201,219	(179,845 to 240,077)	1172	(1049.3 to 1390.3)
	YLDs	Male	1990	1,454	(949 to 2,058)	28	(18.3 to 39.6)
			2013	2,202	(1,448 to 3,049)	31	(20.3 to 42.9)
		Female	1990	1,300	(829 to 1,849)	19.1	(12.2 to 27.0)
			2013	1,571	(1,042 to 2,200)	18.6	(12.1 to 25.8)
		Both	1990	2,755	(1,862 to 3,794)	23	(15.5 to 31.5)
			2013	3,774	(2,578 to 5,137)	24.3	(16.7 to 32.9)
Democratic							
Republic of					/·		
the Congo	Deaths	Male	1990	7,704	(5,991 to 9,793)	153.6	(119.3 to 196.1)
			2013	17,909	(13,028 to 23,154)	186.6	(137.7 to 237.3)
		Female	1990	13,105	(10,038 to 16,685)	217	(167.3 to 270.9)
			2013	21,377	(15,287 to 28,073)	199.8	(146.0 to 261.1)
		Both	1990	20,809	(17,049 to 24,526)	191.8	(159.3 to 225.5)
			2013	39,287	(31,578 to 47,382)	195.2	(158.6 to 232.8)

	DALYs	Male	1990	225,844	(176,320 to 287,309)	2913.3	(2254.1 to 3714.7)
			2013	485,999	(352,969 to 634,975)	3522.1	(2554.9 to 4575.3)
		Female	1990	316,654	(233,321 to 411,657)	3791.4	(2900.6 to 4848.8)
			2013	490,625	(349,471 to 667,171)	3392.1	(2409.0 to 4474.6)
		Both	1990	542,499	(439,545 to 647,560)	3429.8	(2812.9 to 4040.5)
			2013	976,625	(775,563 to 1,188,990)	3471.8	(2782.5 to 4201.2)
	YLDs	Male	1990	1,340	(848 to 1,911)	17.8	(11.5 to 25.7)
			2013	2,593	(1,713 to 3,608)	17.7	(11.7 to 24.5)
		Female	1990	1,145	(734 to 1,614)	13.1	(8.5 to 18.5)
			2013	2,151	(1,411 to 3,074)	13.5	(8.8 to 19.6)
		Both	1990	2,486	(1,643 to 3,406)	15.2	(10.0 to 20.7)
			2013	4,745	(3,219 to 6,421)	15.5	(10.5 to 21.2)
Denmark	Deaths	Male	1990	2,594	(2,132 to 3,349)	84.1	(68.8 to 108.4)
			2013	2,286	(1,557 to 2,789)	55.9	(38.1 to 68.1)
		Female	1990	3,790	(3,008 to 4,367)	75.1	(60.3 to 87.8)
			2013	2,758	(2,193 to 3,551)	46.5	(37.0 to 59.8)
		Both	1990	6,385	(5,522 to 7,384)	79.3	(68.9 to 92.1)
			2013	5,045	(4,137 to 5,894)	50.9	(41.6 to 59.4)
	DALYs	Male	1990	42,131	(35,332 to 54,068)	1358.1	(1139.7 to 1745.9)
			2013	35,322	(25,136 to 42,053)	856.8	(609.0 to 1017.5)
		Female	1990	46,829	(38,421 to 55,090)	1088.8	(904.3 to 1309.7)
			2013	31,436	(26,223 to 39,275)	613.8	(508.2 to 757.9)
		Both	1990	88,961	(79,703 to 103,627)	1207.2	(1082.5 to 1420.6)
			2013	66,758	(55,470 to 75,896)	727.2	(598.3 to 827.5)
	YLDs	Male	1990	3,755	(2,580 to 5,061)	120.6	(83.1 to 161.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	5,073	(3,319 to 7,271)	122.4	(80.1 to 174.4)
		Female	1990	3,435	(2,324 to 4,792)	83.6	(56.9 to 116.6)

		1				1	
			2013	4,014	(2,582 to 5,698)	83.3	(54.3 to 116.6)
		Both	1990	7,190	(5,027 to 9,691)	99.6	(69.5 to 133.5)
			2013	9,087	(6,275 to 12,118)	101.8	(70.5 to 135.6)
Djibouti	Deaths	Male	1990	147	(100 to 212)	197.8	(142.4 to 273.2)
			2013	251	(180 to 377)	156.8	(116.6 to 227.9)
		Female	1990	220	(127 to 329)	256.3	(156.1 to 363.5)
			2013	359	(227 to 542)	201.3	(132.5 to 294.7)
		Both	1990	367	(242 to 527)	230.5	(160.4 to 312.6)
			2013	610	(428 to 903)	181.2	(132.0 to 258.3)
	DALYs	Male	1990	3,726	(2,480 to 5,546)	3444.2	(2364.0 to 4963.5)
			2013	5,526	(3,781 to 8,577)	2651.2	(1892.5 to 3997.1)
		Female	1990	5,029	(2,783 to 7,864)	4391.1	(2525.8 to 6610.1)
			2013	7,051	(4,261 to 11,061)	3252.7	(2026.8 to 4983.6)
		Both	1990	8,756	(5,610 to 12,994)	3961.8	(2596.6 to 5694.6)
			2013	12,578	(8,486 to 19,430)	2976.1	(2061.8 to 4499.5)
	YLDs	Male	1990	17	(11 to 25)	13.1	(8.7 to 19.3)
			2013	34	(21 to 52)	13.3	(8.4 to 20.3)
		Female	1990	12	(7 to 17)	8.6	(5.4 to 12.4)
			2013	26	(17 to 38)	9.7	(6.3 to 14.0)
		Both	1990	29	(19 to 41)	10.7	(7.1 to 14.8)
			2013	60	(39 to 86)	11.4	(7.4 to 16.3)
Dominica	Deaths	Male	1990	25	(21 to 30)	124.6	(102.9 to 151.7)
			2013	30	(21 to 39)	105.4	(77.2 to 138.9)
		Female	1990	35	(29 to 42)	150.7	(127.3 to 182.1)
			2013	31	(23 to 41)	90.5	(66.1 to 118.1)
		Both	1990	60	(53 to 68)	138.9	(122.9 to 158.1)
			2013	62	(50 to 75)	98.2	(80.6 to 120.1)
	DALYs	Male	1990	510	(426 to 611)	2237.2	(1873.7 to 2690.2)

			2013	573	(419 to 769)	1882.7	(1383.5 to 2521.0)
		Female	1990	610	(516 to 729)	2482.4	(2092.5 to 2971.8)
			2013	439	(321 to 586)	1270.1	(928.9 to 1696.6)
		Both	1990	1,120	(993 to 1,259)	2371.1	(2093.5 to 2679.7)
			2013	1,012	(816 to 1,246)	1566.9	(1265.6 to 1927.0)
	YLDs	Male	1990	12	(7 to 17)	55.7	(35.3 to 79.8)
			2013	16	(11 to 23)	56.7	(38.0 to 79.2)
		Female	1990	10	(6 to 14)	42.2	(27.6 to 59.5)
			2013	14	(10 to 19)	42.4	(29.5 to 57.7)
		Both	1990	22	(15 to 30)	48.7	(33.2 to 66.1)
			2013	31	(22 to 42)	49.1	(34.6 to 66.4)
Dominican							
Republic	Deaths	Male	1990	1,503	(1,260 to 1,738)	103.8	(86.1 to 120.2)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
Country	IIICasarc	JCA	2013	4,043	(3,137 to 5,461)	118.4	(92.1 to 158.9)
		Female	1990	1,328	(1,145 to 1,621)	95.4	(81.4 to 117.1)
		remaie	2013	3,834	(2,893 to 5,045)	103	(77.8 to 135.7)
		Both	1990	2,831	(2,550 to 3,166)	99.8	(89.6 to 111.9)
		200	2013	7,878	(6,526 to 9,768)	110.7	(91.8 to 137.5)
	DALYs	Male	1990	34,073	(28,737 to 39,375)	1816	(1528.8 to 2095.9)
			2013	75,891	(57,673 to 103,470)	2049.6	(1565.5 to 2785.2)
		Female	1990	28,873	(24,722 to 34,723)	1588.9	(1376.0 to 1902.5)
			2013	61,627	(46,858 to 82,575)	1585.5	(1202.5 to 2129.0)
		Both	1990	62,947	(56,796 to 70,497)	1706.4	(1539.6 to 1897.5)
			2013	137,519	(111,432 to 174,083)	1813.5	(1476.0 to 2294.6)
							_ · /
	YLDs	Male	1990	1,117	(749 to 1,506)	65.2	(43.8 to 87.4)
	YLDs	Male			(749 to 1,506) (1,454 to 3,142)	65.2 60.9	(43.8 to 87.4) (40.3 to 88.6)
	YLDs	Male Female	1990	1,117		1	,

			2013	1,758	(1,203 to 2,414)	46.1	(31.3 to 63.3)
		Both	1990	1,836	(1,275 to 2,451)	55.1	(38.3 to 74.1)
			2013	3,951	(2,699 to 5,446)	53.3	(36.2 to 73.5)
Ecuador	Deaths	Male	1990	1,513	(1,266 to 1,784)	70	(58.3 to 83.0)
			2013	3,145	(2,460 to 3,980)	60.9	(47.9 to 76.5)
		Female	1990	1,496	(1,157 to 1,733)	61	(47.3 to 71.0)
			2013	2,869	(2,281 to 3,854)	48.3	(38.4 to 65.0)
		Both	1990	3,009	(2,631 to 3,352)	65.2	(56.7 to 73.1)
			2013	6,015	(5,042 to 7,262)	54.3	(45.6 to 65.4)
	DALYs	Male	1990	35,711	(29,709 to 41,592)	1280.9	(1075.2 to 1499.8)
			2013	64,618	(49,243 to 83,136)	1093.7	(842.2 to 1395.1)
		Female	1990	31,936	(25,092 to 36,623)	1068.7	(830.7 to 1219.1)
			2013	50,002	(38,074 to 69,622)	783.9	(598.9 to 1078.8)
		Both	1990	67,647	(59,069 to 74,968)	1169.6	(1024.5 to 1295.9)
			2013	114,620	(94,260 to 141,051)	933	(772.6 to 1140.7)
	YLDs	Male	1990	575	(383 to 798)	21.4	(14.3 to 29.8)
			2013	1,318	(870 to 1,846)	22.9	(15.2 to 32.4)
		Female	1990	331	(222 to 456)	11.5	(7.7 to 15.9)
			2013	739	(497 to 1,025)	11.8	(7.9 to 16.4)
		Both	1990	906	(620 to 1,233)	16.3	(11.1 to 22.0)
			2013	2,057	(1,434 to 2,834)	17.1	(11.9 to 23.4)
Egypt	Deaths	Male	1990	30,633	(25,205 to 35,742)	253.9	(206.1 to 295.7)
			2013	42,850	(37,052 to 50,865)	214.3	(186.7 to 252.8)
		Female	1990	32,611	(26,879 to 36,977)	210.5	(173.3 to 240.3)
			2013	43,095	(36,664 to 51,938)	168.3	(144.2 to 201.2)
		Both	1990	63,244	(54,331 to 70,577)	228.9	(194.5 to 255.3)
			2013	85,945	(77,590 to 98,122)	188.4	(170.6 to 214.6)
	DALYs	Male	1990	963,660	(782,908 to 1,129,509)	4916.4	(4058.7 to 5750.0)

Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
,			2013	1,020,448	(868,191 to 1,216,766)	3762.3	(3235.7 to 4478.4)
		Female	1990	792,416	(607,456 to 942,084)	3764.6	(3097.0 to 4308.7)
			2013	774,206	(653,763 to 928,879)	2612.6	(2207.1 to 3141.0)
					(1,450,099 to		
		Both	1990	1,756,077		4298.4	(3715.7 to 4817.4)
			2010	. =0.4.6==	(1,600,665 to	24.42 =	(2027 0 : 0700 0)
			2013	1,794,655	2,054,314)	3149.7	(2827.8 to 3590.8)
	YLDs	Male	1990	1,702	(1,109 to 2,384)	10.7	(7.0 to 15.1)
			2013	3,051	(2,044 to 4,322)	10.9	(7.2 to 15.4)
		Female	1990	1,646	(1,089 to 2,314)	9	(5.9 to 12.6)
			2013	2,673	(1,696 to 3,784)	8.6	(5.4 to 12.3)
		Both	1990	3,348	(2,265 to 4,564)	9.8	(6.7 to 13.2)
			2013	5,724	(3,822 to 7,910)	9.7	(6.5 to 13.3)
El Salvador	Deaths	Male	1990	686	(567 to 833)	58.3	(47.6 to 71.1)
			2013	894	(705 to 1,153)	44.3	(34.9 to 57.0)
		Female	1990	777	(582 to 895)	54.1	(40.4 to 62.7)
			2013	1,106	(848 to 1,496)	40.2	(30.9 to 54.3)
		Both	1990	1,464	(1,212 to 1,637)	56	(46.5 to 63.0)
			2013	2,000	(1,694 to 2,491)	42	(35.5 to 52.4)
	DALYs	Male	1990	16,978	(14,108 to 20,413)	1099.5	(917.1 to 1318.9)
			2013	16,937	(13,317 to 21,966)	797.4	(625.6 to 1039.9)
		Female	1990	17,085	(12,646 to 19,117)	971.1	(723.1 to 1080.9)
			2013	17,824	(13,754 to 24,611)	648.2	(500.0 to 896.8)
		Both	1990	34,064	(28,395 to 37,957)	1029.8	(861.2 to 1149.0)
			2013	34,761	(29,442 to 43,926)	713.4	(603.4 to 900.5)
	YLDs	Male	1990	345	(233 to 475)	25.9	(17.4 to 35.8)
			2013	544	(354 to 755)	26.8	(17.2 to 37.2)

	Female	1990	289	(197 to 385)	18.7	(12.7 to 24.8)
		2013	486	(316 to 699)	18.2	(11.7 to 26.4)
	Both	1990	635	(433 to 844)	22	(15.1 to 29.3)
		2013	1,031	(697 to 1,402)	21.9	(14.9 to 29.8)
Deaths	Male	1990	203	(140 to 276)	256.9	(178.7 to 342.1)
		2013	264	(185 to 404)	185	(138.0 to 265.0)
	Female	1990	215	(143 to 300)	252	(170.1 to 346.2)
		2013	221	(152 to 363)	179.2	(126.1 to 284.2)
	Both	1990	419	(291 to 562)	256.3	(180.7 to 339.1)
		2013	486	(349 to 739)	183.6	(139.3 to 268.4)
DALYs	Male	1990	5,221	(3,548 to 7,137)	5204.8	(3533.8 to 7091.1)
		2013	7,076	(4,707 to 11,058)	3409.4	(2390.1 to 5153.9)
	Female	1990	4,607	(2,961 to 6,651)	4506.5	(2942.7 to 6445.4)
		2013	5,055	(3,181 to 8,452)	2911.4	(1966.7 to 4816.0)
	Both	1990	9,829	(6,734 to 13,387)	4878.2	(3340.0 to 6653.5)
						Uncertainty
measure	sex	year	number	Uncertainty Interval	Rate	Interval
		2013	12,132	(8,112 to 18,736)	3186.4	(2259.7 to 4864.1)
YLDs	Male	1990	18	(12 to 26)	18	(12.2 to 25.4)
		2013	40	(26 to 60)	18.5	(12.2 to 27.3)
			10	1==/		, ,
	Female	1990	13	(9 to 19)	13.3	(8.8 to 18.4)
	Female			·		
	Female Both	1990	13	(9 to 19)	13.3	(8.8 to 18.4)
		1990 2013	13 27	(9 to 19) (17 to 38)	13.3 13.8	(8.8 to 18.4) (8.9 to 19.3)
Deaths		1990 2013 1990	13 27 32	(9 to 19) (17 to 38) (22 to 44)	13.3 13.8 15.6	(8.8 to 18.4) (8.9 to 19.3) (10.9 to 21.3)
Deaths	Both	1990 2013 1990 2013	13 27 32 67	(9 to 19) (17 to 38) (22 to 44) (45 to 95)	13.3 13.8 15.6 16.2	(8.8 to 18.4) (8.9 to 19.3) (10.9 to 21.3) (10.9 to 22.8)
Deaths	Both	1990 2013 1990 2013 1990	13 27 32 67 430	(9 to 19) (17 to 38) (22 to 44) (45 to 95) (306 to 576)	13.3 13.8 15.6 16.2 152.2	(8.8 to 18.4) (8.9 to 19.3) (10.9 to 21.3) (10.9 to 22.8) (108.8 to 199.5)
	DALYs	Both Deaths Male Female Both DALYS Male Female Both Both Female Both	2013 Both 1990 2013	Both 1990 635 2013 1,031 Deaths Male 1990 203 2013 264 Female 1990 215 2013 221 Both 1990 419 DALYs Male 1990 5,221 DALYs Male 1990 4,607 Female 1990 4,607 Both 1990 9,829 measure sex year number 2013 12,132 YLDs Male 1990 18	Both 1990 635 (433 to 844) 2013 1,031 (697 to 1,402)	Both 1990 635 (433 to 844) 22 Deaths Male 1990 203 (140 to 276) 256.9 Deaths Male 1990 203 (140 to 276) 256.9 Female 1990 215 (143 to 300) 252 Both 1990 215 (143 to 300) 252 Both 1990 419 (291 to 562) 256.3 Both 1990 419 (291 to 562) 256.3 DALYS Male 1990 5,221 (3,548 to 7,137) 5204.8 DALYS Male 1990 5,221 (3,548 to 7,137) 5204.8 Female 1990 4,607 (2,961 to 6,651) 4506.5 Both 1990 9,829 (6,734 to 13,387) 4878.2 measure sex year number Uncertainty Interval Rate measure sex year number Uncertainty Interval Rate

		Both	1990	1,162	(699 to 1,489)	182.3	(107.3 to 235.0)
			2013	2,523	(1,810 to 3,164)	173.6	(123.0 to 217.4)
	DALYs	Male	1990	12,554	(8,939 to 16,636)	2823.4	(2029.9 to 3745.6)
			2013	24,190	(16,596 to 32,867)	2594.7	(1812.2 to 3447.4)
		Female	1990	18,134	(7,813 to 25,241)	3447.7	(1596.9 to 4681.7)
			2013	33,805	(19,739 to 48,606)	3081.3	(1867.0 to 4219.8)
		Both	1990	30,688	(18,443 to 39,428)	3223.7	(1937.8 to 4133.4)
			2013	57,995	(40,690 to 74,628)	2910.2	(2092.4 to 3641.0)
	YLDs	Male	1990	83	(53 to 123)	13.1	(8.8 to 18.7)
			2013	179	(118 to 249)	13.9	(9.0 to 19.3)
		Female	1990	59	(37 to 85)	8.3	(5.3 to 12.2)
			2013	133	(86 to 191)	8.6	(5.5 to 12.2)
		Both	1990	143	(95 to 199)	10.4	(7.1 to 14.1)
			2013	313	(211 to 421)	10.8	(7.4 to 14.7)
Estonia	Deaths	Male	1990	997	(844 to 1,204)	175	(148.1 to 211.9)
			2013	892	(701 to 1,067)	115.7	(90.9 to 138.7)
		Female	1990	2,150	(1,688 to 2,339)	164.4	(128.6 to 178.7)
			2013	1,157	(863 to 1,833)	63	(47.3 to 99.6)
		Both	1990	3,147	(2,645 to 3,427)	170.4	(143.2 to 185.6)
			2013	2,049	(1,666 to 2,747)	81.9	(67.0 to 108.4)
	DALYs	Male	1990	18,934	(16,008 to 23,141)	2915.6	(2477.3 to 3563.6)
			2013	14,633	(11,677 to 17,308)	1871.8	(1489.4 to 2206.1)
		Female	1990	28,555	(21,985 to 31,029)	2328.7	(1777.7 to 2534.5)
			2013	12,389	(9,437 to 19,207)	807.7	(623.2 to 1237.1)
		Both	1990	47,489	(40,625 to 52,074)	2577.1	(2200.8 to 2826.1)
			2013	27,023	(22,504 to 34,268)	1223.5	(1028.8 to 1537.0)
	YLDs	Male	1990	352	(226 to 498)	49.1	(31.7 to 69.4)
			2013	373	(230 to 543)	48.8	(30.4 to 70.8)

		Female	1990	373	(252 to 522)	33.7	(22.6 to 47.0)
			2013	385	(252 to 536)	32	(21.1 to 44.5)
		Both	1990	725	(492 to 988)	40	(27.2 to 54.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	759	(513 to 1,038)	39.1	(26.3 to 54.0)
Ethiopia	Deaths	Male	1990	14,814	(11,169 to 17,965)	187.1	(141.0 to 226.5)
			2013	23,674	(19,160 to 28,487)	160.4	(130.6 to 191.0)
		Female	1990	15,246	(11,971 to 19,249)	195.1	(154.0 to 245.2)
			2013	30,102	(22,393 to 36,271)	188.2	(141.1 to 225.4)
		Both	1990	30,060	(25,366 to 34,459)	191.2	(163.1 to 219.4)
			2013	53,777	(43,469 to 61,946)	175.2	(141.6 to 200.9)
	DALYs	Male	1990	344,053	(258,291 to 424,488)	3475.1	(2628.5 to 4217.0)
			2013	473,723	(376,293 to 578,826)	2639.7	(2126.0 to 3209.9)
		Female	1990	336,654	(262,743 to 425,062)	3431.7	(2681.3 to 4322.0)
			2013	553,015	(408,475 to 677,748)	2945.9	(2193.5 to 3571.0)
		Both	1990	680,708	(564,005 to 781,995)	3452.1	(2898.6 to 3959.7)
			2013	1,026,739	(827,994 to 1,188,943)	2803.3	(2263.3 to 3221.0)
	YLDs	Male	1990	790	(535 to 1,085)	6.9	(4.6 to 9.7)
			2013	1,781	(1,184 to 2,454)	8	(5.4 to 11.0)
		Female	1990	661	(446 to 921)	5.7	(3.8 to 7.7)
			2013	1,289	(873 to 1,787)	5.4	(3.7 to 7.5)
		Both	1990	1,451	(990 to 1,952)	6.3	(4.3 to 8.4)
			2013	3,071	(2,102 to 4,196)	6.6	(4.6 to 9.1)
Federated States of							
Micronesia	Deaths	Male	1990	39	(24 to 55)	223.7	(139.7 to 300.1)
			2013	47	(31 to 69)	201.7	(139.5 to 281.5)
		Female	1990	24	(16 to 36)	136	(91.0 to 206.9)

			2013	34	(21 to 51)	135.7	(87.6 to 197.0)
		Both	1990	64	(42 to 90)	178.5	(124.2 to 242.0)
			2013	81	(54 to 118)	166.3	(116.6 to 231.0)
	DALYs	Male	1990	1,079	(661 to 1,589)	4699	(2876.0 to 6739.4)
			2013	1,218	(776 to 1,890)	4101.4	(2714.3 to 6134.2)
		Female	1990	578	(367 to 883)	2579.8	(1663.9 to 3932.5)
			2013	718	(429 to 1,135)	2426.2	(1490.7 to 3722.6)
		Both	1990	1,658	(1,065 to 2,439)	3622.8	(2353.3 to 5215.9)
			2013	1,937	(1,230 to 2,964)	3230.7	(2108.8 to 4810.1)
	YLDs	Male	1990	10	(7 to 14)	54.4	(36.7 to 77.4)
			2013	14	(9 to 20)	56	(38.3 to 78.9)
		Female	1990	7	(5 to 10)	39	(26.4 to 54.1)
			2013	12	(8 to 17)	43.5	(29.6 to 58.7)
		Both	1990	18	(12 to 25)	46.5	(31.9 to 63.2)
			2013	26	(18 to 36)	49.3	(34.0 to 66.2)
Fiji	Deaths	Male	1990	247	(187 to 327)	187.5	(146.6 to 239.6)
			2013	324	(245 to 424)	139.1	(108.5 to 175.9)
		Female	1990	306	(181 to 410)	240	(145.1 to 315.0)
			2013	330	(253 to 478)	132.6	(104.3 to 184.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	553	(415 to 675)	214.6	(162.4 to 258.0)
			2013	654	(537 to 816)	136.2	(114.0 to 168.0)
	DALYs	Male	1990	7,541	(5,541 to 10,324)	3809	(2918.8 to 5048.0)
			2013	8,907	(6,669 to 11,925)	2760.4	(2100.5 to 3591.6)
		Female	1990	8,056	(4,683 to 10,924)	4592.4	(2719.4 to 6160.4)
			2013	7,748	(5,792 to 11,344)	2399.2	(1826.5 to 3487.2)
		Both	1990	15,597	(11,719 to 19,341)	4205.7	(3166.6 to 5141.2)
			2013	16,655	(13,384 to 21,144)	2581.9	(2117.5 to 3224.1)

		Both	1990	52,384	(45,342 to 58,515)	61	(53.2 to 68.1)
			2013	49,081	(42,019 to 59,935)	34.1	(29.5 to 40.7)
	DALYs	Male	1990	328,019	(282,094 to 387,969)	1081.6	(930.6 to 1280.2)
			2013	279,127	(227,254 to 332,158)	597	(485.6 to 711.9)
		Female	1990	337,966	(272,205 to 386,595)	713.8	(579.7 to 817.4)
			2013	264,242	(224,407 to 341,093)	382.8	(326.1 to 487.7)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	665,985	(597,548 to 739,733)	875.8	(785.7 to 976.6)
			2013	543,369	(471,393 to 626,910)	479.2	(414.0 to 548.0)
	YLDs	Male	1990	28,221	(18,209 to 40,206)	92.3	(59.6 to 130.8)
			2013	41,260	(27,277 to 57,971)	90.3	(59.7 to 127.5)
		Female	1990	26,510	(17,395 to 37,606)	61.3	(40.5 to 85.7)
			2013	37,935	(23,724 to 54,159)	62	(40.1 to 87.0)
		Both	1990	54,732	(38,251 to 74,357)	74.2	(52.2 to 99.6)
			2013	79,196	(53,660 to 105,965)	74.9	(50.5 to 100.5)
Gabon	Deaths	Male	1990	695	(564 to 835)	287.5	(233.4 to 342.3)
			2013	1,179	(920 to 1,462)	296.1	(230.9 to 361.6)
		Female	1990	671	(542 to 832)	229.9	(186.6 to 286.4)
			2013	1,059	(763 to 1,469)	224.7	(162.7 to 310.8)
		Both	1990	1,366	(1,186 to 1,573)	255.9	(222.3 to 295.6)
			2013	2,239	(1,851 to 2,755)	259	(214.3 to 319.2)
	DALYs	Male	1990	15,416	(12,407 to 18,659)	5476.9	(4431.4 to 6609.7)
			2013	26,738	(20,403 to 34,202)	5724.3	(4398.5 to 7200.6)
		Female	1990	12,455	(9,877 to 15,388)	3730.6	(2973.5 to 4595.9)
			2013	17,975	(12,529 to 25,805)	3593.8	(2506.2 to 5123.5)
		Both	1990	27,872	(23,886 to 32,178)	4525.2	(3898.2 to 5207.0)
			2013	44,714	(36,828 to 55,010)	4630.4	(3806.8 to 5705.1)
	YLDs	Male	1990	53	(35 to 75)	18.7	(12.3 to 26.3)

			2013	85	(55 to 119)	17.6	(11.1 to 24.6)
		Female	1990	48	(31 to 67)	13.9	(9.0 to 19.3)
			2013	73	(46 to 102)	13.8	(8.7 to 19.7)
		Both	1990	102	(67 to 137)	16	(10.6 to 21.5)
			2013	158	(104 to 215)	15.6	(10.2 to 21.2)
Georgia	Deaths	Male	1990	4,939	(4,307 to 5,822)	264.5	(230.8 to 309.7)
			2013	6,300	(5,252 to 7,361)	263	(220.0 to 307.7)
		Female	1990	6,433	(5,773 to 7,469)	194.1	(174.2 to 225.4)
			2013	6,520	(5,089 to 7,533)	155.8	(121.4 to 180.5)
		Both	1990	11,373	(10,438 to 12,897)	220.8	(202.2 to 250.2)
			2013	12,820	(11,038 to 14,140)	198.7	(171.4 to 219.4)
	DALYs	Male	1990	101,038	(88,129 to 120,312)	4598.9	(4007.4 to 5458.1)
			2013	109,010	(90,759 to 127,760)	4476	(3737.2 to 5251.8)
		Female	1990	98,732	(88,489 to 114,768)	2940	(2634.2 to 3408.9)
			2013	80,730	(63,271 to 94,606)	2163.6	(1695.4 to 2546.4)
		Both	1990	199,771	(182,938 to 223,971)	3616.3	(3322.8 to 4050.5)
			2013	189,741	(165,386 to 211,508)	3134.1	(2734.6 to 3505.7)
	YLDs	Male	1990	396	(266 to 569)	16.5	(11.1 to 23.7)
			2013	400	(254 to 578)	16.6	(10.5 to 23.7)
		Female	1990	429	(273 to 612)	12.9	(8.3 to 18.3)
			2013	431	(282 to 603)	12.8	(8.4 to 17.7)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	825	(575 to 1,135)	14.5	(10.2 to 19.8)
			2013	831	(559 to 1,127)	14.5	(9.8 to 19.6)
Germany	Deaths	Male	1990	41,458	(34,260 to 49,896)	103.8	(85.9 to 124.8)
			2013	34,818	(27,812 to 42,325)	47.2	(37.7 to 57.5)
		Female	1990	80,678	(60,405 to 88,570)	91.8	(69.7 to 100.5)
			2013	47,165	(39,344 to 66,591)	38.9	(32.8 to 54.3)

			, ,				
		Both	1990	122,137	(100,823 to 134,575)	96.7	(80.0 to 106.4)
			2013	81,984	(71,442 to 101,935)	42.8	(37.4 to 52.9)
	DALYs	Male	1990	645,789	(544,031 to 769,060)	1547.9	(1309.9 to 1845.8)
			2013	522,272	(433,255 to 611,429)	734.8	(607.8 to 862.4)
		Female	1990	912,115	(716,315 to 991,367)	1184.2	(935.1 to 1288.9)
			2013	509,638	(439,241 to 670,431)	517.4	(449.0 to 673.6)
					(1,341,820 to		
		Both	1990	1,557,905	1,708,268)	1330.6	(1157.1 to 1457.3)
			2013	1,031,910	(910,697 to 1,200,845)	617.9	(547.4 to 708.9)
	YLDs	Male	1990	49,459	(34,066 to 68,172)	116.1	(80.5 to 159.5)
			2013	96,102	(66,372 to 128,196)	137.6	(95.5 to 183.4)
		Female	1990	58,705	(39,911 to 81,330)	83.9	(57.0 to 115.7)
			2013	94,618	(66,067 to 124,828)	107.4	(74.6 to 142.1)
		Both	1990	108,165	(76,467 to 143,205)	96.4	(68.2 to 128.1)
			2013	190,721	(136,118 to 249,622)	121.3	(85.6 to 159.2)
Ghana	Deaths	Male	1990	1,916	(1,515 to 2,359)	91	(71.9 to 110.8)
			2013	7,058	(5,731 to 8,416)	164.4	(135.4 to 195.9)
		Female	1990	1,793	(1,081 to 3,848)	83.9	(49.7 to 178.9)
			2013	8,775	(4,862 to 11,727)	170.8	(94.5 to 226.6)
		Both	1990	3,710	(2,784 to 5,747)	87.4	(65.1 to 138.2)
			2013	15,834	(11,727 to 19,027)	169.8	(125.7 to 203.6)
	DALYs	Male	1990	53,957	(42,692 to 67,360)	1677.1	(1328.3 to 2056.8)
			2013	176,841	(139,055 to 215,653)	2991.3	(2412.1 to 3573.6)
		Female	1990	45,604	(28,589 to 94,717)	1434.9	(868.1 to 3072.0)
			2013	175,476	(96,395 to 237,735)	2743.3	(1512.6 to 3704.3)
		Both	1990	99,561	(76,884 to 148,872)	1554.7	(1171.5 to 2406.1)
			2013	352,318	(261,390 to 425,448)	2877.2	(2137.8 to 3442.1)
	YLDs	Male	1990	638	(412 to 918)	19.5	(12.6 to 27.9)
			2013	1,236	(788 to 1,758)	19.5	(12.5 to 28.1)
					-		

		Female	1990	490	(308 to 701)	14.5	(9.1 to 20.9)
			2013	1,079	(699 to 1,553)	14.9	(9.4 to 21.9)
		Both	1990	1,129	(747 to 1,576)	16.9	(11.2 to 23.7)
			2013	2,316	(1,542 to 3,134)	17	(11.2 to 23.2)
Greece	Deaths	Male	1990	7,881	(6,762 to 9,097)	130.7	(111.7 to 151.0)
			2013	8,826	(7,354 to 10,479)	83.7	(69.9 to 98.7)
		Female	1990	11,380	(9,117 to 12,375)	141.1	(112.8 to 153.3)
			2013	13,384	(11,289 to 17,756)	86	(72.6 to 113.8)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	19,261	(16,901 to 20,810)	137.2	(120.4 to 148.3)
			2013	22,211	(19,589 to 27,057)	85.7	(75.9 to 104.0)
	DALYs	Male	1990	112,257	(97,944 to 130,183)	1842.6	(1606.8 to 2137.4)
			2013	108,186	(91,709 to 126,047)	1146.2	(972.0 to 1336.3)
		Female	1990	132,142	(107,550 to 142,386)	1710.7	(1397.2 to 1842.1)
			2013	127,432	(110,060 to 166,426)	950.5	(825.2 to 1246.8)
		Both	1990	244,400	(218,383 to 264,915)	1780.1	(1592.6 to 1929.5)
			2013	235,619	(210,478 to 279,112)	1047.8	(936.7 to 1240.8)
	YLDs	Male	1990	6,582	(4,371 to 9,217)	107.2	(71.2 to 149.8)
			2013	9,629	(6,272 to 13,693)	106.2	(69.9 to 151.1)
		Female	1990	5,193	(3,529 to 7,090)	69.5	(47.0 to 94.9)
			2013	8,010	(5,114 to 11,386)	69.5	(45.0 to 97.9)
		Both	1990	11,776	(8,176 to 15,619)	86.5	(59.8 to 114.7)
			2013	17,639	(12,019 to 24,729)	86.3	(59.3 to 119.1)
Grenada	Deaths	Male	1990	41	(35 to 49)	151.7	(128.1 to 179.8)
			2013	45	(37 to 54)	138.7	(114.1 to 164.2)
		Female	1990	67	(55 to 75)	168.1	(138.2 to 191.1)
			2013	64	(52 to 81)	130.3	(107.5 to 165.3)
		Both	1990	108	(96 to 120)	162.1	(143.6 to 179.7)

			2013	110	(96 to 129)	135.2	(117.8 to 158.5)
	DALYs	Male	1990	844	(725 to 995)	2730.1	(2331.1 to 3220.4)
			2013	903	(732 to 1,093)	2484.5	(2023.5 to 2986.3)
		Female	1990	1,147	(930 to 1,292)	2806.8	(2276.5 to 3172.4)
			2013	956	(779 to 1,214)	2074	(1696.2 to 2639.3)
		Both	1990	1,992	(1,773 to 2,194)	2783.8	(2473.1 to 3062.1)
			2013	1,859	(1,619 to 2,151)	2275.7	(1981.5 to 2632.6)
	YLDs	Male	1990	18	(12 to 24)	62.6	(42.4 to 84.8)
			2013	21	(13 to 29)	59.7	(39.5 to 83.0)
		Female	1990	16	(10 to 23)	40.8	(26.3 to 58.4)
			2013	19	(12 to 26)	42	(27.5 to 57.8)
		Both	1990	34	(24 to 46)	49.9	(35.1 to 67.5)
			2013	40	(27 to 54)	49.7	(34.2 to 67.5)
Guatemala	Deaths	Male	1990	896	(761 to 1,078)	58.8	(48.7 to 71.6)
			2013	1,459	(1,183 to 1,821)	43.1	(34.7 to 54.2)
		Female	1990	847	(741 to 1,266)	52.8	(45.8 to 78.5)
			2013	1,995	(1,311 to 2,354)	49.5	(32.7 to 58.4)
		Both	1990	1,744	(1,547 to 2,207)	55.7	(48.9 to 70.9)
			2013	3,455	(2,704 to 3,975)	46.6	(36.6 to 54.0)
	DALYs	Male	1990	22,899	(19,458 to 27,374)	1053.4	(900.2 to 1256.8)
			2013	31,051	(25,546 to 37,802)	753.3	(617.2 to 919.5)
		Female	1990	21,077	(18,311 to 31,983)	940.9	(826.2 to 1399.9)
			2013	39,538	(26,247 to 46,531)	836.3	(566.7 to 980.9)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	43,976	(38,983 to 55,908)	996.2	(888.4 to 1258.6)
			2013	70,590	(55,886 to 80,391)	798.4	(638.9 to 908.4)
	YLDs	Male	1990	470	(323 to 640)	22.4	(15.5 to 30.3)
			2013	1,013	(665 to 1,411)	26.6	(17.4 to 37.6)

		Female	1990	362	(244 to 498)	17.7	(12.0 to 24.4)
			2013	776	(515 to 1,110)	17.5	(11.4 to 25.3)
		Both	1990	832	(580 to 1,095)	20	(14.1 to 26.3)
			2013	1,790	(1,212 to 2,436)	21.7	(14.6 to 29.6)
Guinea	Deaths	Male	1990	1,116	(873 to 1,350)	108.2	(85.4 to 130.9)
			2013	2,897	(2,216 to 3,613)	159.4	(123.7 to 194.2)
		Female	1990	1,558	(804 to 2,099)	136.5	(74.5 to 183.7)
			2013	3,363	(2,233 to 4,867)	172.2	(115.8 to 242.5)
		Both	1990	2,674	(1,838 to 3,237)	124	(86.1 to 150.9)
			2013	6,260	(4,933 to 7,789)	166.7	(132.7 to 206.5)
	DALYs	Male	1990	34,020	(26,010 to 42,357)	2041	(1608.4 to 2472.5)
			2013	73,489	(54,999 to 93,642)	2843.7	(2176.3 to 3550.2)
		Female	1990	40,723	(19,337 to 56,451)	2436.4	(1258.4 to 3278.4)
			2013	75,540	(49,101 to 111,740)	2882	(1911.8 to 4164.5)
		Both	1990	74,743	(48,295 to 91,566)	2260	(1558.4 to 2732.6)
			2013	149,030	(116,667 to 186,520)	2871.3	(2267.9 to 3571.6)
	YLDs	Male	1990	283	(176 to 412)	19.3	(11.8 to 28.0)
			2013	539	(350 to 748)	19.4	(12.7 to 27.0)
		Female	1990	230	(147 to 328)	14	(8.9 to 19.7)
			2013	429	(277 to 624)	14.2	(9.2 to 21.0)
		Both	1990	514	(334 to 706)	16.4	(10.9 to 22.7)
			2013	969	(658 to 1,324)	16.6	(11.3 to 22.8)
Guinea-Bissau	Deaths	Male	1990	195	(128 to 307)	116.1	(78.5 to 176.5)
			2013	546	(367 to 819)	196.4	(135.0 to 286.5)
		Female	1990	326	(200 to 499)	172.2	(109.4 to 257.5)
			2013	636	(401 to 1,026)	218	(140.2 to 337.4)
		Both	1990	521	(354 to 783)	147.8	(102.4 to 216.3)
			2013	1,182	(816 to 1,741)	208.6	(146.1 to 296.4)

	DALYs	Male	1990	5,664	(3,741 to 8,638)	2237.2	(1474.9 to 3478.7)
			2013	15,032	(10,039 to 22,560)	3767.9	(2538.3 to 5641.1)
		Female	1990	8,287	(4,958 to 12,673)	3128.9	(1902.8 to 4801.0)
			2013	15,703	(9,793 to 26,052)	3897.4	(2461.1 to 6250.4)
		Both	1990	13,951	(9,665 to 20,512)	2727.1	(1842.2 to 4102.7)
			2013	30,736	(21,258 to 45,584)	3847.1	(2670.6 to 5651.9)
	YLDs	Male	1990	46	(30 to 66)	19.8	(12.7 to 28.1)
			2013	79	(51 to 112)	19.3	(12.8 to 27.5)
		Female	1990	37	(25 to 51)	14.2	(9.5 to 19.7)
			2013	63	(39 to 88)	14.4	(9.3 to 20.0)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
,		Both	1990	83	(57 to 114)	16.8	(11.7 to 22.7)
			2013	142	(95 to 194)	16.7	(11.5 to 22.7)
Guyana	Deaths	Male	1990	460	(385 to 531)	266.3	(222.4 to 307.3)
			2013	301	(218 to 395)	232	(176.5 to 290.0)
		Female	1990	460	(373 to 511)	233.7	(189.6 to 259.3)
			2013	393	(289 to 506)	208.6	(155.7 to 264.5)
		Both	1990	920	(813 to 1,007)	249.5	(220.6 to 272.7)
			2013	695	(560 to 840)	220.8	(180.3 to 263.5)
	DALYs	Male	1990	10,112	(8,558 to 11,646)	5257	(4428.0 to 6043.0)
			2013	8,368	(6,111 to 11,193)	4514.2	(3343.8 to 5775.6)
		Female	1990	8,936	(7,106 to 9,943)	4292.2	(3442.5 to 4763.3)
			2013	8,320	(5,978 to 11,192)	3675.3	(2669.1 to 4825.9)
		Both	1990	19,049	(16,768 to 20,926)	4759	(4196.4 to 5235.6)
			2013	16,688	(13,314 to 20,542)	4066.7	(3285.2 to 4920.9)
	YLDs	Male	1990	113	(73 to 157)	62.2	(39.7 to 86.7)
			2013	95	(63 to 134)	62.1	(40.7 to 87.6)
		Female	1990	89	(59 to 127)	43.7	(29.1 to 62.6)

			2013	93	(63 to 127)	43.8	(29.5 to 60.1)
		Both	1990	202	(139 to 275)	52.4	(36.1 to 71.5)
			2013	189	(130 to 258)	50.6	(35.2 to 68.5)
Haiti	Deaths	Male	1990	2,615	(2,204 to 3,026)	174.3	(146.4 to 201.1)
			2013	3,654	(3,005 to 4,389)	165.3	(138.1 to 198.3)
		Female	1990	4,353	(3,622 to 5,246)	264.8	(220.8 to 316.2)
			2013	5,963	(4,574 to 7,387)	227.7	(177.2 to 280.6)
		Both	1990	6,968	(6,109 to 7,874)	223.2	(196.5 to 251.7)
			2013	9,617	(8,060 to 11,152)	199.9	(169.0 to 230.9)
	DALYs	Male	1990	82,444	(67,914 to 98,114)	3677.9	(3088.3 to 4265.7)
			2013	91,690	(73,850 to 111,470)	3107	(2548.9 to 3737.8)
		Female	1990	119,703	(98,192 to 147,585)	5350.8	(4435.8 to 6484.3)
			2013	133,229	(100,950 to 166,068)	4154.4	(3154.6 to 5166.2)
		Both	1990	202,148	(174,633 to 231,506)	4570.8	(3989.5 to 5174.4)
			2013	224,920	(187,759 to 262,554)	3673.7	(3073.4 to 4264.1)
	YLDs	Male	1990	931	(610 to 1,312)	60.9	(39.2 to 86.9)
			2013	1,489	(990 to 2,055)	60	(39.7 to 82.8)
		Female	1990	718	(493 to 977)	42.8	(29.4 to 58.6)
			2013	1,233	(820 to 1,675)	43	(27.9 to 59.3)
		Both	1990	1,649	(1,153 to 2,246)	51.2	(35.7 to 70.6)
			2013	2,723	(1,895 to 3,654)	50.7	(35.1 to 68.4)
Honduras	Deaths	Male	1990	743	(575 to 1,143)	85.4	(67.2 to 129.3)
			2013	1,635	(1,110 to 2,579)	89.1	(61.3 to 138.0)
		Female	1990	883	(674 to 1,104)	90.6	(70.2 to 113.3)
			2013	1,966	(1,265 to 2,717)	92.1	(58.4 to 126.4)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	1,627	(1,382 to 2,123)	88.4	(76.0 to 113.4)
			2013	3,602	(2,659 to 4,824)	91.1	(67.6 to 120.1)

			Г				
	DALYs	Male	1990	21,418	(16,045 to 34,184)	1613.8	(1244.7 to 2478.9)
			2013	34,894	(23,052 to 57,264)	1557.5	(1036.6 to 2483.7)
		Female	1990	23,329	(16,849 to 30,061)	1677.6	(1280.1 to 2102.1)
			2013	36,134	(22,407 to 52,237)	1495.9	(944.7 to 2125.9)
		Both	1990	44,747	(36,460 to 60,592)	1647.2	(1380.8 to 2179.8)
			2013	71,028	(50,178 to 99,533)	1529.3	(1106.7 to 2087.8)
	YLDs	Male	1990	277	(188 to 386)	26.4	(17.9 to 37.6)
			2013	618	(416 to 832)	28.1	(19.1 to 38.7)
		Female	1990	201	(132 to 280)	18.5	(11.8 to 25.9)
			2013	449	(303 to 630)	19.1	(12.8 to 26.9)
		Both	1990	478	(325 to 646)	22.4	(15.1 to 30.2)
			2013	1,068	(750 to 1,437)	23.4	(16.4 to 31.5)
Hungary	Deaths	Male	1990	10,240	(8,852 to 11,767)	206.4	(177.7 to 238.0)
			2013	7,969	(6,849 to 9,402)	128.7	(110.3 to 151.5)
		Female	1990	13,126	(10,381 to 14,586)	158.6	(124.9 to 176.1)
			2013	9,983	(8,489 to 14,154)	82.3	(70.5 to 116.9)
		Both	1990	23,366	(20,688 to 25,405)	178.5	(158.2 to 194.9)
			2013	17,953	(15,984 to 22,316)	100.4	(89.5 to 124.0)
	DALYs	Male	1990	186,914	(161,717 to 217,696)	3485.4	(3014.2 to 4039.8)
			2013	126,316	(107,811 to 147,785)	2002.8	(1714.4 to 2337.2)
		Female	1990	184,373	(145,937 to 200,881)	2336.5	(1844.2 to 2542.5)
			2013	110,630	(95,596 to 157,263)	1063.5	(919.6 to 1518.1)
		Both	1990	371,288	(328,060 to 405,337)	2831.3	(2504.9 to 3089.9)
			2013	236,947	(212,934 to 288,385)	1454.9	(1306.4 to 1756.0)
	YLDs	Male	1990	1,702	(1,105 to 2,360)	30.9	(20.1 to 43.0)
			2013	2,098	(1,390 to 2,931)	33.6	(22.4 to 46.5)
		Female	1990	1,675	(1,096 to 2,344)	22.8	(14.8 to 32.1)
			2013	1,857	(1,225 to 2,600)	21.2	(13.9 to 29.8)

		Both	1990	3,378	(2,260 to 4,457)	26.4	(17.7 to 34.8)
			2013	3,955	(2,731 to 5,395)	26.6	(18.4 to 36.1)
Iceland	Deaths	Male	1990	74	(60 to 93)	64	(52.2 to 80.1)
			2013	79	(58 to 100)	39.1	(28.8 to 49.7)
		Female	1990	99	(80 to 121)	61.8	(50.1 to 75.7)
			2013	95	(73 to 121)	33.6	(25.8 to 42.4)
		Both	1990	173	(152 to 202)	63	(55.4 to 73.3)
			2013	175	(142 to 209)	36.3	(29.5 to 43.0)
	DALYs	Male	1990	1,149	(968 to 1,399)	967.8	(814.6 to 1178.7)
			2013	1,118	(854 to 1,386)	574.2	(437.6 to 714.5)
		Female	1990	1,242	(1,014 to 1,481)	854.8	(696.3 to 1021.9)
			2013	974	(771 to 1,188)	408.7	(325.1 to 499.8)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	2,391	(2,139 to 2,715)	906.9	(809.5 to 1026.6)
			2013	2,092	(1,739 to 2,437)	487.7	(404.0 to 570.7)
	YLDs	Male	1990	120	(78 to 170)	101.4	(65.9 to 143.0)
			2013	205	(141 to 288)	106.4	(73.2 to 148.7)
		Female	1990	108	(72 to 148)	77.7	(51.9 to 105.7)
			2013	155	(101 to 225)	71.3	(47.3 to 103.4)
		Both	1990	229	(157 to 307)	88.6	(60.7 to 119.1)
			2013	360	(245 to 489)	88.1	(60.2 to 119.1)
India	Deaths	Male	1990	205,140	(164,823 to 249,155)	109.5	(87.7 to 132.6)
			2013	365,297	(286,511 to 465,499)	109.8	(87.0 to 139.2)
		Female	1990	208,439	(153,149 to 243,557)	116.7	(85.9 to 135.4)
			2013	348,918	(249,980 to 433,870)	99.1	(71.9 to 121.3)
		Both	1990	413,579	(346,014 to 473,116)	113.3	(94.2 to 129.0)
			2013	714,215	(590,159 to 841,167)	104.5	(86.2 to 121.8)
	DALYs	Male	1990	5,303,689	(4,295,763 to	2129	(1716.9 to 2576.1)

					6,434,369)		
					(6,563,543 to		
			2013	8,523,060	10,980,735)	2023.3	(1579.1 to 2564.7)
					(3,696,535 to		,
		Female	1990	5,027,199	5,925,911)	2129.8	(1569.3 to 2492.5)
					(4,763,513 to		
			2013	6,880,827	8,768,688)	1639.1	(1159.5 to 2057.4)
					(8,597,565 to		
		Both	1990	10,330,888	11,898,336)	2131.4	(1784.0 to 2443.4)
					(12,643,000 to		
			2013	15,403,888	18,446,373)	1831.4	(1511.3 to 2161.1)
	YLDs	Male	1990	110,023	(76,279 to 149,594)	42.3	(29.5 to 57.7)
			2013	198,460	(133,355 to 271,598)	43.3	(29.4 to 60.0)
		Female	1990	86,931	(58,259 to 117,698)	34.3	(23.1 to 46.8)
			2013	166,710	(107,989 to 234,701)	35.6	(23.3 to 49.7)
		Both	1990	196,955	(136,268 to 261,352)	38.3	(26.8 to 50.5)
			2013	365,171	(250,006 to 490,584)	39.3	(26.7 to 52.9)
Indonesia	Deaths	Male	1990	62,470	(54,333 to 71,376)	194.3	(168.5 to 220.4)
			2013	134,060	(110,863 to 157,090)	214	(178.9 to 247.9)
		Female	1990	66,082	(56,778 to 79,265)	177	(148.0 to 209.3)
			2013	155,856	(125,126 to 179,636)	216.6	(174.7 to 249.1)
		Both	1990	128,553	(116,549 to 141,419)	184.9	(165.0 to 204.6)
			2013	289,917	(256,355 to 321,575)	216.6	(191.5 to 238.8)
					(1,294,430 to		
	DALYs	Male	1990	1,499,375	1,733,856)	3443.9	(2992.0 to 3936.1)
					(2,468,232 to		
			2013	3,059,116	3,632,739)	3764.6	(3098.6 to 4409.0)
			4005		(1,247,026 to	00	/ac+= =
		Female	1990	1,429,799	1,778,326)	3009	(2617.5 to 3650.4)

			2013	3,617	(2,476 to 4,971)	12.6	(8.7 to 17.2)
		Both	1990	2,756	(1,862 to 3,820)	9.6	(6.5 to 13.3)
			2013	7,782	(5,395 to 10,550)	13.6	(9.5 to 18.5)
Iraq	Deaths	Male	1990	4,864	(3,680 to 6,202)	155.1	(117.8 to 196.0)
			2013	7,092	(5,056 to 9,869)	139.8	(101.2 to 190.7)
		Female	1990	5,975	(4,635 to 7,837)	170.3	(132.4 to 219.9)
			2013	10,305	(7,552 to 13,399)	163.9	(121.8 to 209.5)
		Both	1990	10,840	(8,906 to 13,054)	163.6	(135.7 to 194.6)
			2013	17,398	(13,940 to 21,085)	154.1	(124.3 to 184.2)
	DALYs	Male	1990	118,200	(87,913 to 151,522)	2653.9	(1962.4 to 3460.9)
			2013	168,561	(118,571 to 237,119)	2342	(1656.6 to 3291.0)
		Female	1990	120,172	(90,149 to 163,716)	2814.1	(2128.5 to 3791.1)
			2013	202,443	(143,339 to 275,450)	2624.1	(1875.2 to 3469.4)
		Both	1990	238,373	(193,895 to 292,556)	2742.2	(2235.0 to 3357.7)
			2013	371,005	(289,210 to 460,878)	2505.7	(1969.5 to 3077.9)
	YLDs	Male	1990	422	(269 to 621)	10.4	(6.5 to 15.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Country	measure	sex	year 2013	number 821	Uncertainty Interval (556 to 1,157)	Rate 10.4	
Country	measure	sex Female	· ·		•		Interval
Country	measure		2013	821	(556 to 1,157)	10.4	Interval (7.0 to 14.5)
Country	measure		2013 1990	821 354	(556 to 1,157) (229 to 489)	10.4 8.3	(7.0 to 14.5) (5.3 to 11.5)
Country	measure	Female	2013 1990 2013	821 354 727	(556 to 1,157) (229 to 489) (487 to 1,013)	10.4 8.3 8.4	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5)
Country	measure	Female	2013 1990 2013 1990	821 354 727 777	(556 to 1,157) (229 to 489) (487 to 1,013) (515 to 1,062)	10.4 8.3 8.4 9.3	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5) (6.2 to 12.6)
		Female Both	2013 1990 2013 1990 2013	821 354 727 777 1,549	(556 to 1,157) (229 to 489) (487 to 1,013) (515 to 1,062) (1,059 to 2,107)	10.4 8.3 8.4 9.3 9.3	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5) (6.2 to 12.6) (6.5 to 12.5)
		Female Both	2013 1990 2013 1990 2013 1990	821 354 727 777 1,549 1,274	(556 to 1,157) (229 to 489) (487 to 1,013) (515 to 1,062) (1,059 to 2,107) (1,054 to 1,579)	10.4 8.3 8.4 9.3 9.3 82.5	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5) (6.2 to 12.6) (6.5 to 12.5) (68.1 to 102.3)
		Female Both Male	2013 1990 2013 1990 2013 1990 2013	821 354 727 777 1,549 1,274 1,061	(556 to 1,157) (229 to 489) (487 to 1,013) (515 to 1,062) (1,059 to 2,107) (1,054 to 1,579) (820 to 1,339)	10.4 8.3 8.4 9.3 9.3 82.5 43.7	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5) (6.2 to 12.6) (6.5 to 12.5) (68.1 to 102.3) (33.8 to 55.2)
		Female Both Male	2013 1990 2013 1990 2013 1990 2013 1990	821 354 727 777 1,549 1,274 1,061 1,828	(556 to 1,157) (229 to 489) (487 to 1,013) (515 to 1,062) (1,059 to 2,107) (1,054 to 1,579) (820 to 1,339) (1,415 to 2,111)	10.4 8.3 8.4 9.3 9.3 82.5 43.7 80.9	(7.0 to 14.5) (5.3 to 11.5) (5.6 to 11.5) (6.2 to 12.6) (6.5 to 12.5) (68.1 to 102.3) (33.8 to 55.2) (62.9 to 94.1)

	DALYs	Male	1990	21,293	(17,940 to 25,639)	1299.7	(1097.1 to 1572.2)
	DALLIS	iviaic	2013	16,537		656.8	(521.8 to 807.0)
		Female	1990	24,134		1140	(905.3 to 1328.7)
		Temate	2013	16,660	(13,404 to 21,466)	535.6	(431.7 to 696.5)
		Both	1990	45,428	(40,165 to 51,060)	1213.5	(1075.7 to 1364.2)
		20011	2013	33,197	-	593.3	(505.9 to 696.1)
	YLDs	Male	1990	1,805	(1,214 to 2,477)	108.6	(73.0 to 149.0)
			2013	2,718	(1,776 to 3,826)	105.8	(69.0 to 149.3)
		Female	1990	1,579	(1,055 to 2,242)	76.9	(51.4 to 108.1)
			2013	2,178	(1,472 to 3,017)	73.7	(50.1 to 102.1)
		Both	1990	3,384	(2,320 to 4,524)	91	(62.8 to 121.4)
			2013	4,896	(3,321 to 6,656)	88.8	(60.3 to 120.6)
Israel	Deaths	Male	1990	1,238	(1,018 to 1,637)	71.3	(58.4 to 94.4)
			2013	1,008	(670 to 1,282)	26.4	(17.6 to 33.5)
		Female	1990	1,464	(1,189 to 1,820)	69.6	(56.4 to 86.5)
			2013	1,672	(1,022 to 2,228)	29.7	(18.4 to 39.7)
		Both	1990	2,702	(2,395 to 3,225)	70.5	(62.5 to 84.2)
			2013	2,680	(1,867 to 3,240)	28.6	(19.9 to 34.4)
	DALYs	Male	1990	19,937	(16,699 to 26,322)	1125.3	(941.3 to 1479.3)
			2013	15,170	(10,577 to 18,473)	408.3	(284.9 to 497.6)
		Female	1990	20,762	(17,332 to 25,533)	987.8	(824.6 to 1210.8)
			2013	17,359	(11,609 to 22,727)	352.5	(238.8 to 462.4)
		Both	1990	40,700	(36,252 to 48,557)	1052.6	(937.5 to 1257.2)
			2013	32,530	(24,575 to 38,059)	380.7	(288.0 to 443.8)
	YLDs	Male	1990	1,248	(852 to 1,662)	69.9	(47.6 to 93.2)
			2013	2,719	(1,800 to 3,772)	73.4	(49.0 to 101.6)
		Female	1990	1,266	(829 to 1,755)	60.8	(39.8 to 83.9)
			2013	2,462	(1,692 to 3,365)	53.9	(37.0 to 72.9)

		Both	1990	2,515	(1,711 to 3,341)	65	(44.0 to 86.4)
			2013	5,182	(3,612 to 6,926)	62.8	(44.0 to 83.7)
Italy	Deaths	Male	1990	31,410	(26,611 to 36,525)	99.4	(84.1 to 115.7)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	27,869	(22,097 to 33,813)	47.1	(37.4 to 57.1)
		Female	1990	44,686	(35,462 to 49,382)	83.2	(66.0 to 91.5)
			2013	42,687	(34,491 to 55,687)	42.1	(34.3 to 55.2)
		Both	1990	76,097	(66,326 to 83,467)	90.2	(78.7 to 99.0)
			2013	70,556	(60,359 to 82,030)	44.6	(38.2 to 51.5)
	DALYs	Male	1990	480,657	(413,775 to 553,322)	1461.8	(1261.8 to 1681.4)
			2013	344,645	(282,741 to 410,270)	644.3	(525.8 to 766.3)
		Female	1990	507,137	(411,188 to 553,272)	1047.3	(849.5 to 1150.4)
			2013	365,507	(304,813 to 469,890)	462.8	(387.2 to 598.1)
		Both	1990	987,795	(883,151 to 1,079,234)	1226.4	(1098.1 to 1340.0)
			2013	710,153	(618,687 to 807,266)	545	(475.4 to 619.9)
	YLDs	Male	1990	29,795	(20,325 to 41,669)	88.4	(60.9 to 123.4)
			2013	37,836	(24,962 to 51,521)	74.8	(49.4 to 101.6)
		Female	1990	23,579	(16,229 to 32,260)	51.2	(35.6 to 69.3)
			2013	28,329	(17,988 to 40,393)	44.3	(29.1 to 61.7)
		Both	1990	53,374	(36,938 to 70,987)	67	(46.3 to 89.6)
			2013	66,166	(45,170 to 88,779)	58	(39.7 to 76.8)
Jamaica	Deaths	Male	1990	881	(761 to 1,008)	117.1	(100.9 to 134.3)
			2013	1,093	(833 to 1,378)	105.4	(80.5 to 133.0)
		Female	1990	1,433	(1,213 to 1,746)	146.2	(123.7 to 177.7)
			2013	1,681	(1,250 to 2,120)	120.3	(89.6 to 152.7)
		Both	1990	2,315	(2,064 to 2,644)	133.9	(119.3 to 153.0)
			2013	2,774	(2,260 to 3,291)	114.8	(93.2 to 136.4)
	DALYs	Male	1990	15,368	(13,447 to 17,610)	1938.7	(1695.1 to 2218.0)

			2013	18,492	(14,008 to 23,576)	1670.2	(1264.3 to 2128.8)
		Female	1990	20,695	(17,414 to 24,168)	2227.5	(1878.0 to 2604.8)
			2013	22,069	(16,505 to 28,875)	1689.3	(1246.9 to 2219.0)
		Both	1990	36,064	(32,380 to 40,260)	2101.5	(1885.5 to 2342.9)
			2013	40,562	(33,261 to 48,457)	1691.2	(1386.3 to 2021.5)
	YLDs	Male	1990	479	(327 to 653)	62.9	(42.8 to 85.7)
			2013	660	(446 to 959)	60.3	(40.7 to 86.9)
		Female	1990	373	(238 to 525)	41	(26.3 to 57.6)
			2013	526	(338 to 744)	40.6	(26.4 to 57.4)
		Both	1990	852	(594 to 1,129)	51.1	(35.6 to 67.7)
			2013	1,187	(812 to 1,608)	49.6	(34.1 to 67.5)
Japan	Deaths	Male	1990	60,064	(53,539 to 67,711)	106	(93.9 to 119.8)
			2013	85,537	(71,303 to 99,514)	62.8	(52.3 to 73.0)
		Female	1990	71,040	(58,347 to 80,114)	82.8	(68.0 to 93.4)
			2013	100,039	(81,366 to 141,583)	42.8	(35.3 to 60.2)
		Both	1990	131,104	(118,833 to 142,701)	92.7	(83.8 to 101.2)
			2013	185,577	(163,108 to 228,573)	51.8	(45.8 to 62.9)
	DALYs	Male	1990	1,051,122	(936,413 to 1,173,900)	1657	(1477.7 to 1851.7)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2012	4 407 226	(1,011,607 to	4004.5	(054.2 : 4450.4)
			2013	1,187,336	1,355,752)	1004.5	(854.3 to 1150.4)
		Female	1990	902,537	(740,983 to 1,000,591)	1069.2	(878.0 to 1189.2)
			2013	897,168	(760,465 to 1,220,113)	533.4	(453.2 to 720.3)
			4000		(1,785,194 to	1000	/4244 Q : 442 7 Q
		Both	1990	1,953,659	2,107,863)	1329.3	(1214.9 to 1435.0)
			2013	2,084,505	(1,868,717 to 2,421,503)	751.7	(670.3 to 860.7)
	YLDs	Male	1990		(52,495 to 105,531)	120.2	(82.9 to 164.8)
	TLUS	iviale	1990	76,635	(32,493 (0 103,331)	120.2	(02.9 (0 104.8)

			2012	120 705	(02 E69 to 100 1E0)	116.3	/77 0 to 150 6\
		F	2013	139,785	(92,568 to 190,150)	116.3	(77.8 to 158.6)
		Female	1990	61,646	(41,888 to 84,156)	73.1	(49.7 to 99.7)
			2013	117,166		74.7	(50.6 to 103.3)
		Both	1990	138,281	(95,988 to 184,549)	93.4	(64.9 to 124.4)
			2013	256,951	(178,079 to 343,784)	93.7	(65.6 to 125.3)
Jordan	Deaths	Male	1990	931	(740 to 1,196)	162.7	(130.5 to 208.7)
			2013	973	(767 to 1,232)	91.7	(73.4 to 113.9)
		Female	1990	829	(662 to 1,016)	155.2	(124.2 to 190.6)
			2013	998	(800 to 1,197)	94.5	(77.2 to 112.5)
		Both	1990	1,760	(1,499 to 2,063)	159	(135.7 to 186.9)
			2013	1,972	(1,689 to 2,269)	93.5	(80.7 to 106.7)
	DALYs	Male	1990	17,082	(13,297 to 22,150)	2487.8	(1938.9 to 3236.6)
			2013	17,945	(13,681 to 23,332)	1225.4	(959.6 to 1560.7)
		Female	1990	14,767	(11,703 to 18,072)	2284	(1815.2 to 2834.1)
			2013	16,112	(12,443 to 19,931)	1141.6	(906.5 to 1367.7)
		Both	1990	31,849	(26,864 to 37,551)	2387.5	(2024.0 to 2826.6)
			2013	34,058	(28,417 to 40,052)	1185.7	(1011.5 to 1370.0)
	YLDs	Male	1990	82	(55 to 117)	10.9	(7.2 to 15.6)
			2013	231	(148 to 330)	11.5	(7.5 to 16.5)
		Female	1990	60	(38 to 86)	8.4	(5.4 to 12.2)
			2013	167	(107 to 242)	8.5	(5.5 to 12.2)
		Both	1990	143	(97 to 192)	9.7	(6.6 to 12.9)
			2013	399	(267 to 555)	10	(6.8 to 14.0)
Kazakhstan	Deaths	Male	1990	8,538	(7,210 to 9,893)	254.6	(216.4 to 293.0)
			2013	11,472	(9,488 to 13,816)	268.8	(225.8 to 317.8)
		Female	1990	14,183	(12,408 to 16,250)	211.2	(183.9 to 242.6)
			2013	15,612	(11,789 to 18,007)	216.1	(162.3 to 248.6)
		Both	1990	22,722	(20,364 to 25,226)	227.7	(204.2 to 252.5)

			2013	27,085	(22,740 to 30,505)	238.5	(199.0 to 266.3)
	DALYs	Male	1990	190,471	(159,236 to 224,952)	4429.6	(3724.1 to 5142.3)
			2013	261,742	(212,199 to 324,703)	4820.2	(3969.6 to 5844.1)
		Female	1990	225,828	(198,131 to 260,790)	3241.5	(2845.8 to 3751.6)
			2013	254,157	(191,015 to 299,243)	3259.1	(2448.1 to 3819.5)
		Both	1990	416,300	(376,414 to 464,537)	3711.4	(3347.8 to 4137.9)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
			2013	515,899	(439,453 to 590,439)	3920.2	(3332.9 to 4464.7)
	YLDs	Male	1990	891	(566 to 1,257)	17.4	(11.3 to 24.3)
			2013	1,053	(700 to 1,509)	17.3	(11.6 to 24.7)
		Female	1990	936	(621 to 1,329)	13.1	(8.6 to 18.6)
			2013	1,015	(675 to 1,404)	12.4	(8.2 to 17.2)
		Both	1990	1,827	(1,253 to 2,477)	15	(10.2 to 20.3)
			2013	2,069	(1,439 to 2,783)	14.4	(10.0 to 19.3)
Kenya	Deaths	Male	1990	2,895	(2,359 to 3,420)	96.3	(78.5 to 115.8)
			2013	5,427	(4,276 to 6,671)	95.8	(75.4 to 115.1)
		Female	1990	3,774	(2,807 to 4,585)	120.2	(88.8 to 148.9)
			2013	6,116	(4,293 to 7,913)	98.2	(69.3 to 127.1)
		Both	1990	6,669	(5,470 to 7,742)	109.1	(88.5 to 127.9)
			2013	11,544	(9,319 to 13,732)	97.5	(78.7 to 115.9)
	DALYs	Male	1990	59,969	(48,476 to 71,010)	1560.5	(1270.9 to 1843.3)
			2013	113,668	(88,542 to 144,285)	1522.9	(1194.2 to 1879.5)
		Female	1990	69,679	(52,561 to 83,220)	1803.6	(1352.9 to 2184.7)
			2013	106,376	(75,882 to 140,935)	1345.8	(960.1 to 1774.4)
		Both	1990	129,648	(106,806 to 149,325)	1688.8	(1390.3 to 1954.0)
			2013	220,045	(177,305 to 267,448)	1432.2	(1168.2 to 1708.5)
	YLDs	Male	1990	631	(399 to 897)	13	(8.2 to 18.7)
			2013	1,426	(866 to 2,109)	14	(8.3 to 20.5)

		Female	1990	454	(290 to 682)	9.1	(5.8 to 13.7)
			2013	1,001	(642 to 1,447)	9.2	(6.0 to 13.2)
		Both	1990	1,086	(712 to 1,510)	11	(7.2 to 15.7)
			2013	2,427	(1,580 to 3,335)	11.4	(7.5 to 15.9)
Kiribati	Deaths	Male	1990	37	(28 to 47)	266.6	(206.6 to 325.3)
			2013	76	(51 to 104)	308.5	(218.2 to 398.0)
		Female	1990	18	(13 to 27)	110.2	(80.5 to 168.9)
			2013	37	(22 to 55)	137.7	(82.3 to 200.2)
		Both	1990	55	(45 to 68)	178.2	(147.1 to 218.0)
			2013	114	(83 to 148)	212.2	(158.8 to 266.1)
	DALYs	Male	1990	1,241	(940 to 1,567)	6309	(4786.9 to 7949.4)
			2013	2,402	(1,615 to 3,364)	7187.7	(4905.6 to 9766.7)
		Female	1990	496	(361 to 764)	2374.9	(1719.5 to 3655.2)
			2013	914	(530 to 1,387)	2704	(1590.8 to 4023.4)
		Both	1990	1,737	(1,401 to 2,150)	4180.2	(3401.0 to 5152.6)
			2013	3,317	(2,359 to 4,439)	4787.5	(3492.4 to 6258.2)
	YLDs	Male	1990	7	(5 to 10)	56.5	(37.8 to 77.4)
			2013	14	(9 to 19)	58.1	(38.3 to 80.0)
		Female	1990	6	(4 to 9)	38.4	(25.2 to 53.2)
			2013	12	(8 to 17)	43.1	(29.1 to 59.3)
		Both	1990	14	(9 to 19)	46.1	(31.8 to 62.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	27	(18 to 36)	49.6	(34.3 to 65.5)
Kuwait	Deaths	Male	1990	173	(144 to 204)	81.1	(67.4 to 97.2)
			2013	343	(276 to 413)	65.1	(52.1 to 77.9)
		Female	1990	144	(118 to 177)	84.6	(69.9 to 105.4)
			2013	199	(149 to 224)	71.5	(52.7 to 82.0)
		Both	1990	317	(280 to 356)	82.8	(73.3 to 95.2)

DALYs Male 1990 4,843 (4,012 to 5,729) 1170.7 2013 7,136 (5,851 to 8,704) 865.5	(55.6 to 76.1) (981.2 to 1378.0) (702.6 to 1035.2) (999.5 to 1472.7) (654.2 to 980.6)
2013 7,136 (5,851 to 8,704) 865.5	(702.6 to 1035.2) (999.5 to 1472.7) (654.2 to 980.6)
	(999.5 to 1472.7) (654.2 to 980.6)
Female 1990 3,800 (3,039 to 4,573) 1208.7	(654.2 to 980.6)
	•
2013 4,340 (3,227 to 4,965) 872.2	(4040 4 ±= 4333 0)
Both 1990 8,643 (7,568 to 9,768) 1184.9	(1048.4 to 1323.9)
2013 11,476 (9,741 to 13,158) 861.6	(725.7 to 981.4)
YLDs Male 1990 61 (39 to 90) 10.7	(7.0 to 15.4)
2013 134 (88 to 191) 11.3	(7.5 to 15.6)
Female 1990 32 (20 to 46) 9	(5.7 to 12.9)
2013 61 (39 to 85) 8.9	(5.8 to 12.4)
Both 1990 94 (62 to 129) 10	(6.7 to 14.0)
2013 196 (131 to 265) 10.4	(7.1 to 13.9)
Kyrgyzstan Deaths Male 1990 2,244 (1,935 to 2,544) 263.3	(225.7 to 297.8)
2013 2,976 (2,451 to 3,597) 251.1	(207.6 to 301.0)
Female 1990 3,397 (2,818 to 3,729) 227.2	(188.5 to 247.4)
2013 3,076 (2,598 to 3,745) 193.7	(163.1 to 238.2)
Both 1990 5,641 (5,006 to 6,078) 241.9	(212.6 to 260.7)
2013 6,052 (5,362 to 6,941) 219	(195.1 to 252.9)
DALYs Male 1990 49,325 (42,539 to 56,617) 4671	(4019.6 to 5310.9)
2013 69,483 (57,160 to 84,374) 4768.1	(3942.4 to 5768.1)
Female 1990 56,450 (48,760 to 64,093) 3611.5	(3094.1 to 4077.5)
2013 56,398 (46,953 to 67,880) 3179.8	(2655.4 to 3837.4)
Both 1990 105,776 (96,248 to 115,432) 4048.3	(3656.7 to 4396.0)
2013 125,881 (110,275 to 143,367) 3888.6	(3417.0 to 4415.4)
YLDs Male 1990 203 (134 to 280) 16.4	(10.9 to 22.8)
2013 279 (184 to 386) 16.6	(11.3 to 22.9)
Female 1990 209 (132 to 292) 12.9	(8.1 to 18.2)

			2013	249	(162 to 343)	12.7	(8.3 to 17.6)
		Both	1990	413	(269 to 551)	14.4	(9.6 to 19.4)
			2013	528	(352 to 718)	14.5	(9.8 to 19.6)
Laos	Deaths	Male	1990	1,332	(824 to 2,055)	179.2	(116.3 to 271.0)
			2013	2,284	(1,511 to 3,461)	181.4	(127.7 to 261.0)
		Female	1990	1,480	(972 to 2,275)	173.2	(118.1 to 259.9)
			2013	2,733	(1,684 to 3,983)	187.7	(122.5 to 266.3)
		Both	1990	2,812	(1,845 to 4,324)	176.7	(121.8 to 261.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	5,018	(3,329 to 7,363)	185.7	(130.5 to 263.1)
	DALYs	Male	1990	39,785	(24,665 to 60,899)	3656.7	(2262.1 to 5693.8)
			2013	59,030	(37,289 to 92,624)	3433.4	(2238.9 to 5263.9)
		Female	1990	37,391	(22,937 to 57,238)	3247.2	(2066.5 to 5061.1)
			2013	59,002	(34,046 to 90,833)	3211.1	(1930.4 to 4804.3)
		Both	1990	77,177	(49,226 to 117,129)	3449	(2252.1 to 5314.9)
			2013	118,033	(73,542 to 179,193)	3326.3	(2159.5 to 4968.9)
	YLDs	Male	1990	479	(320 to 651)	58.6	(39.1 to 80.0)
			2013	832	(550 to 1,157)	55.8	(36.1 to 78.2)
		Female	1990	415	(277 to 559)	42.3	(28.7 to 56.5)
			2013	786	(515 to 1,076)	46.5	(30.8 to 64.9)
		Both	1990	894	(614 to 1,201)	49.4	(33.9 to 65.2)
			2013	1,618	(1,107 to 2,142)	50.8	(34.9 to 68.1)
Latvia	Deaths	Male	1990	2,342	(2,035 to 2,710)	232.6	(201.7 to 268.5)
			2013	1,906	(1,610 to 2,187)	159.1	(134.2 to 182.9)
		Female	1990	4,493	(3,878 to 5,129)	198.2	(171.0 to 225.9)
			2013	3,168	(2,802 to 3,931)	108.2	(96.0 to 134.2)
		Both	1990	6,835	(6,166 to 7,509)	211.4	(190.5 to 232.4)
			2013	5,074	(4,607 to 5,869)	127	(115.3 to 146.4)

	DALYs	Male	1990	41,111	(35,530 to 48,193)	3697.5	(3216.0 to 4301.5)
			2013	30,684	(25,897 to 35,175)	2500.5	(2103.6 to 2875.1)
		Female	1990	58,019	(50,128 to 65,451)	2716.2	(2338.0 to 3061.2)
			2013	33,710	(30,016 to 41,396)	1352.7	(1194.0 to 1671.1)
		Both	1990	99,131	(90,203 to 108,850)	3092.3	(2818.6 to 3402.3)
			2013	64,394	(58,216 to 73,002)	1798.4	(1618.1 to 2053.2)
	YLDs	Male	1990	465	(294 to 662)	37.8	(24.3 to 53.9)
			2013	554	(349 to 805)	46.5	(29.6 to 66.9)
		Female	1990	509	(329 to 705)	26.6	(17.2 to 36.9)
			2013	490	(312 to 708)	25.8	(16.5 to 37.2)
		Both	1990	975	(646 to 1,317)	31.3	(20.8 to 42.2)
			2013	1,045	(684 to 1,439)	34.3	(22.4 to 47.1)
Lebanon	Deaths	Male	1990	450	(352 to 588)	66.4	(52.0 to 86.2)
			2013	359	(246 to 527)	20.5	(14.0 to 30.1)
		Female	1990	1,091	(765 to 1,398)	136.3	(96.3 to 174.3)
			2013	1,711	(678 to 2,457)	85.6	(33.9 to 123.0)
		Both	1990	1,542	(1,222 to 1,857)	103.3	(81.6 to 124.0)
			2013	2,071	(1,019 to 2,829)	54.8	(26.9 to 74.6)
	DALYs	Male	1990	9,320	(7,204 to 12,502)	1034.9	(806.3 to 1371.8)
			2013	5,606	(3,868 to 8,192)	270.7	(187.6 to 391.6)
		Female	1990	21,079	(14,081 to 27,368)	2201.4	(1503.6 to 2837.5)
			2013	23,682	(9,102 to 34,677)	1131.6	(435.4 to 1654.3)
		Both	1990	30,399	(23,435 to 37,193)	1634.7	(1278.6 to 1981.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	29,289	(14,415 to 41,125)	704.1	(348.6 to 982.9)
	YLDs	Male	1990	94	(61 to 131)	10	(6.6 to 13.8)
			2013	245	(159 to 351)	10.6	(6.9 to 15.0)
		Female	1990	78	(52 to 110)	8	(5.4 to 11.1)

			2013	186	(120 to 269)	8.7	(5.6 to 12.7)
		Both	1990	173	(118 to 231)	9	(6.2 to 11.9)
			2013	432	(294 to 591)	9.7	(6.6 to 13.2)
Lesotho	Deaths	Male	1990	391	(296 to 485)	138.9	(104.9 to 172.0)
			2013	689	(540 to 860)	185.5	(145.9 to 230.0)
		Female	1990	408	(293 to 509)	112.9	(82.9 to 140.7)
			2013	651	(465 to 968)	117.8	(84.8 to 174.5)
		Both	1990	799	(640 to 943)	124.2	(98.7 to 146.4)
			2013	1,340	(1,089 to 1,699)	144.5	(117.3 to 183.9)
	DALYs	Male	1990	8,873	(6,745 to 11,097)	2554.8	(1931.6 to 3168.4)
			2013	17,058	(13,314 to 21,195)	3740.2	(2919.0 to 4688.5)
		Female	1990	8,044	(5,623 to 10,187)	1833	(1311.8 to 2299.4)
			2013	13,743	(9,534 to 21,638)	2088.2	(1470.9 to 3191.1)
		Both	1990	16,917	(13,519 to 20,085)	2146.1	(1718.0 to 2543.6)
			2013	30,801	(24,836 to 39,437)	2750.3	(2225.2 to 3505.4)
	YLDs	Male	1990	90	(61 to 125)	24.7	(17.0 to 34.6)
			2013	111	(73 to 153)	24.4	(16.0 to 33.5)
		Female	1990	112	(72 to 160)	23.8	(15.5 to 33.8)
			2013	143	(93 to 203)	23.1	(15.0 to 33.0)
		Both	1990	203	(138 to 276)	24.2	(16.5 to 33.0)
			2013	254	(170 to 343)	23.7	(15.8 to 32.3)
Liberia	Deaths	Male	1990	422	(326 to 515)	127.3	(100.8 to 152.9)
			2013	1,149	(909 to 1,419)	185.7	(148.0 to 222.9)
		Female	1990	496	(397 to 678)	148.4	(120.9 to 199.0)
			2013	1,272	(910 to 1,880)	188.4	(135.7 to 276.1)
		Both	1990	919	(789 to 1,089)	138.5	(120.0 to 162.9)
			2013	2,422	(2,001 to 3,040)	187.4	(155.5 to 237.3)
	DALYs	Male	1990	13,672	(10,200 to 17,422)	2376.9	(1855.0 to 2897.0)

			2013	28,507	(21,943 to 36,434)	3231.7	(2549.4 to 3974.8)
		Female	1990	13,928	(10,442 to 20,195)	2601.1	(2101.4 to 3521.7)
			2013	28,069	(20,005 to 42,862)	3115.3	(2231.1 to 4608.0)
		Both	1990	27,600	(22,241 to 33,944)	2497.1	(2151.2 to 2943.8)
			2013	56,576	(46,321 to 71,348)	3176.1	(2628.0 to 3999.6)
	YLDs	Male	1990	95	(62 to 138)	20	(13.3 to 28.6)
			2013	189	(126 to 263)	19.1	(12.7 to 26.7)
		Female	1990	74	(49 to 106)	14.4	(9.6 to 20.7)
			2013	156	(103 to 217)	14.5	(9.6 to 20.1)
		Both	1990	169	(115 to 240)	17.1	(11.6 to 24.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	346	(235 to 468)	16.7	(11.3 to 22.7)
Libya	Deaths	Male	1990	826	(636 to 1,076)	116.4	(88.7 to 147.7)
			2013	1,567	(1,186 to 2,014)	116	(88.1 to 147.4)
		Female	1990	1,026	(789 to 1,293)	145.9	(114.3 to 180.9)
			2013	2,110	(1,522 to 2,756)	133.2	(96.1 to 173.9)
		Both	1990	1,853	(1,555 to 2,203)	131.5	(111.1 to 156.3)
			2013	3,677	(2,959 to 4,477)	125.7	(101.0 to 152.0)
	DALYs	Male	1990	28,452	(21,228 to 38,466)	2018.5	(1554.2 to 2632.4)
			2013	32,217	(24,458 to 42,254)	1761.3	(1331.0 to 2285.7)
		Female	1990	31,712	(22,765 to 41,376)	2507.8	(1917.5 to 3174.7)
			2013	37,765	(26,925 to 49,589)	1936.5	(1401.3 to 2552.3)
		Both	1990	60,165	(48,457 to 73,395)	2255.5	(1888.4 to 2681.5)
			2013	69,983	(56,354 to 85,590)	1855.4	(1502.6 to 2262.9)
	YLDs	Male	1990	30	(18 to 47)	2.8	(1.7 to 4.3)
			2013	49	(31 to 72)	2.4	(1.6 to 3.6)
		Female	1990	22	(13 to 34)	2.4	(1.5 to 3.6)
			2013	45	(28 to 68)	2.2	(1.3 to 3.3)

		Both	1990	52	(33 to 77)	2.6	(1.7 to 3.8)
			2013	94	(61 to 136)	2.3	(1.5 to 3.3)
Lithuania	Deaths	Male	1990	1,814	(1,532 to 2,237)	122.2	(103.1 to 150.6)
			2013	1,778	(1,433 to 2,113)	96.9	(78.3 to 115.2)
		Female	1990	2,976	(2,674 to 3,735)	110	(99.4 to 137.9)
			2013	3,416	(2,590 to 3,881)	82.6	(63.5 to 93.2)
		Both	1990	4,791	(4,366 to 5,659)	115.3	(105.1 to 136.0)
			2013	5,194	(4,258 to 5,780)	89.6	(73.9 to 99.4)
	DALYs	Male	1990	33,059	(28,114 to 40,329)	2094	(1777.6 to 2551.3)
			2013	29,777	(24,251 to 35,287)	1642.7	(1335.8 to 1950.7)
		Female	1990	41,536	(37,125 to 50,712)	1623.9	(1445.1 to 1973.8)
			2013	37,332	(29,465 to 41,763)	1082.3	(869.3 to 1224.9)
		Both	1990	74,596	(68,206 to 86,334)	1823	(1665.9 to 2113.6)
			2013	67,110	(57,035 to 74,450)	1323.4	(1137.8 to 1475.5)
	YLDs	Male	1990	538	(347 to 772)	32	(20.8 to 45.9)
			2013	636	(402 to 911)	36.3	(23.2 to 51.5)
		Female	1990	521	(342 to 734)	21.6	(14.2 to 30.5)
			2013	595	(394 to 873)	21.8	(14.2 to 32.0)
		Both	1990	1,060	(717 to 1,461)	26	(17.7 to 35.8)
			2013	1,231	(854 to 1,695)	27.9	(19.3 to 38.2)
Luxembourg	Deaths	Male	1990	219	(182 to 262)	120.2	(99.1 to 143.5)
			2013	187	(147 to 229)	54.1	(42.5 to 66.2)
		Female	1990	367	(266 to 412)	110.4	(80.2 to 123.6)
			2013	260	(195 to 405)	46.9	(35.2 to 73.6)
		Both	1990	587	(483 to 649)	114.9	(94.7 to 127.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	447	(369 to 597)	50.4	(41.9 to 67.1)
	DALYs	Male	1990	3,519	(2,970 to 4,178)	1795	(1514.6 to 2128.5)

			2013	2,694	(2,175 to 3,241)	793.2	(640.9 to 953.5)
		Female	1990	4,455	(3,333 to 4,953)	1465.2	(1086.0 to 1632.9)
			2013	2,670	(2,061 to 4,038)	583.3	(451.1 to 886.3)
		Both	1990	7,975	(6,708 to 8,772)	1604.6	(1353.7 to 1763.5)
			2013	5,364	(4,543 to 6,789)	681.6	(576.5 to 853.7)
	YLDs	Male	1990	208	(141 to 290)	103.2	(70.1 to 143.6)
			2013	366	(248 to 513)	108.6	(73.8 to 152.5)
		Female	1990	208	(135 to 297)	72.3	(47.7 to 102.1)
			2013	302	(204 to 411)	74.1	(50.0 to 100.2)
		Both	1990	417	(286 to 570)	84.9	(58.6 to 115.7)
			2013	669	(461 to 891)	89.9	(62.1 to 119.8)
Macedonia	Deaths	Male	1990	1,828	(1,644 to 2,020)	288.9	(260.0 to 317.6)
			2013	2,500	(2,148 to 2,839)	240.8	(207.7 to 273.0)
		Female	1990	1,759	(1,604 to 1,987)	246	(224.9 to 277.7)
			2013	2,840	(2,340 to 3,218)	206.7	(170.0 to 233.8)
		Both	1990	3,587	(3,358 to 3,855)	266.1	(248.9 to 285.9)
			2013	5,341	(4,740 to 5,797)	221.8	(196.6 to 240.3)
	DALYs	Male	1990	31,104	(27,796 to 34,776)	4158	(3717.2 to 4613.7)
			2013	37,679	(32,174 to 42,664)	3280.5	(2813.9 to 3726.8)
		Female	1990	28,002	(25,396 to 31,971)	3412.8	(3111.6 to 3883.1)
			2013	36,264	(30,148 to 41,105)	2592	(2149.7 to 2938.1)
		Both	1990	59,106	(55,233 to 64,076)	3765.2	(3526.7 to 4048.9)
			2013	73,944	(65,784 to 80,699)	2903.1	(2586.7 to 3165.1)
	YLDs	Male	1990	245	(158 to 346)	28	(18.1 to 39.5)
			2013	392	(254 to 565)	31.4	(20.5 to 44.9)
		Female	1990	172	(111 to 242)	18.9	(12.4 to 26.5)
			2013	246	(157 to 358)	18.1	(11.6 to 26.0)
		Both	1990	418	(279 to 566)	23.3	(15.6 to 31.5)

			2013	638	(427 to 869)	24.4	(16.4 to 33.2)
Madagascar	Deaths	Male	1990	7,694	(6,587 to 8,871)	374	(323.7 to 426.8)
			2013	13,350	(10,486 to 18,134)	358.7	(296.9 to 473.4)
		Female	1990	6,496	(5,329 to 7,525)	356.7	(296.1 to 408.9)
			2013	13,318	(9,543 to 17,231)	355.9	(263.2 to 448.9)
		Both	1990	14,190	(12,522 to 15,802)	365.8	(323.5 to 404.4)
			2013	26,668	(21,649 to 32,954)	357.9	(297.8 to 432.0)
	DALYs	Male	1990	203,274	(170,864 to 237,195)	7554.4	(6444.9 to 8766.3)
			2013	353,194	(256,349 to 486,633)	7007.9	(5442.1 to 9609.5)
		Female	1990	151,915	(121,808 to 178,295)	6385.8	(5196.6 to 7410.7)
			2013	293,260	(202,255 to 395,435)	6113.5	(4329.8 to 7991.4)
		Both	1990	355,189	(308,000 to 398,119)	6969.7	(6132.7 to 7786.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	646,454	(498,966 to 819,891)	6559.5	(5243.0 to 8169.3)
	YLDs	Male	1990	384	(244 to 557)	13.6	(8.7 to 19.6)
			2013	788	(505 to 1,113)	14.6	(9.5 to 20.9)
		Female	1990	242	(149 to 359)	9	(5.6 to 13.2)
			2013	552	(359 to 801)	9.5	(6.1 to 13.7)
		Both	1990	627	(411 to 875)	11.3	(7.5 to 15.8)
			2013	1,340	(900 to 1,835)	11.9	(8.0 to 16.4)
Malawi	Deaths	Male	1990	913	(712 to 1,116)	75.8	(57.7 to 93.6)
			2013	2,407	(1,843 to 3,108)	107.5	(83.1 to 136.6)
		Female	1990	1,671	(1,241 to 2,028)	123.2	(89.7 to 151.2)
			2013	3,093	(2,303 to 3,939)	122.2	(91.0 to 155.5)
		Both	1990	2,584	(2,040 to 2,988)	102	(78.5 to 119.6)
			2013	5,500	(4,442 to 6,623)	115.9	(93.3 to 138.9)
	DALYs	Male	1990	25,443	(19,704 to 31,631)	1330.7	(1036.8 to 1619.5)
			2013	52,856	(39,902 to 68,728)	1781.4	(1359.5 to 2316.7)

		Female	1990	38,869	(29,452 to 47,832)	2042.8	(1527.8 to 2465.1)
			2013	58,474	(43,528 to 75,785)	1819.7	(1348.9 to 2343.1)
		Both	1990	64,312	(52,264 to 74,653)	1721.8	(1366.9 to 1992.2)
			2013	111,331	(90,476 to 134,501)	1807.8	(1459.3 to 2182.1)
	YLDs	Male	1990	259	(165 to 383)	12.8	(8.1 to 18.7)
			2013	468	(299 to 673)	12.8	(7.9 to 18.7)
		Female	1990	194	(123 to 275)	8.7	(5.5 to 12.6)
			2013	357	(224 to 519)	8.9	(5.6 to 13.1)
		Both	1990	454	(304 to 634)	10.6	(7.1 to 14.7)
			2013	825	(555 to 1,143)	10.7	(7.2 to 14.9)
Malaysia	Deaths	Male	1990	4,751	(3,965 to 5,587)	138	(112.2 to 164.9)
			2013	8,550	(6,946 to 10,922)	102.8	(83.9 to 132.1)
		Female	1990	4,942	(3,739 to 5,538)	145.2	(108.6 to 165.1)
			2013	6,381	(5,357 to 8,301)	86.8	(72.2 to 114.2)
		Both	1990	9,693	(8,070 to 10,785)	142	(116.7 to 158.8)
			2013	14,932	(12,963 to 18,389)	94.4	(81.5 to 116.7)
	DALYs	Male	1990	114,278	(94,102 to 133,279)	2520.8	(2094.5 to 2952.6)
			2013	191,927	(155,584 to 243,052)	1882.9	(1533.4 to 2386.0)
		Female	1990	103,868	(80,937 to 114,958)	2394.9	(1841.8 to 2661.5)
			2013	127,449	(106,085 to 165,228)	1326.2	(1116.1 to 1708.8)
		Both	1990	218,146	(185,222 to 242,202)	2460.8	(2062.1 to 2736.1)
			2013	319,376	(277,623 to 390,623)	1597	(1394.9 to 1955.5)
	YLDs	Male	1990	2,352	(1,592 to 3,334)	57	(38.2 to 81.0)
			2013	6,366	(4,257 to 8,780)	67.1	(44.8 to 93.3)
		Female	1990	2,086	(1,420 to 2,853)	50.7	(34.2 to 71.0)
			2013	4,993	(3,289 to 6,785)	51.3	(34.4 to 69.4)
		Both	1990	4,438	(3,091 to 5,965)	53.8	(37.4 to 71.9)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval

			2013	11,359	(7,911 to 14,912)	59.2	(41.9 to 77.4)
Maldives	Deaths	Male	1990	97	(64 to 113)	222.7	(144.6 to 263.1)
			2013	74	(59 to 107)	80.8	(63.1 to 118.4)
		Female	1990	39	(22 to 70)	145.4	(94.9 to 233.8)
			2013	58	(35 to 71)	82.4	(50.5 to 100.4)
		Both	1990	136	(108 to 164)	192.7	(151.3 to 225.9)
			2013	133	(106 to 157)	80.7	(64.2 to 95.2)
	DALYs	Male	1990	3,078	(2,221 to 3,646)	4328.4	(2874.0 to 5060.4)
			2013	1,502	(1,227 to 1,967)	1324.9	(1076.9 to 1815.7)
		Female	1990	1,168	(582 to 2,456)	2628.6	(1556.8 to 4511.5)
			2013	1,015	(637 to 1,254)	1087.5	(679.5 to 1341.5)
		Both	1990	4,246	(3,408 to 5,392)	3613.5	(2878.2 to 4330.4)
			2013	2,517	(2,059 to 2,917)	1203.7	(978.1 to 1408.1)
	YLDs	Male	1990	28	(18 to 39)	55.6	(37.3 to 75.4)
			2013	61	(40 to 87)	59.9	(39.1 to 86.4)
		Female	1990	16	(11 to 21)	47.4	(32.4 to 63.7)
			2013	43	(29 to 59)	46.9	(32.0 to 64.4)
		Both	1990	44	(30 to 59)	52.1	(36.4 to 69.4)
			2013	104	(73 to 141)	53.6	(37.4 to 72.3)
Mali	Deaths	Male	1990	1,337	(1,068 to 1,607)	91.7	(72.1 to 110.9)
			2013	2,401	(1,841 to 3,078)	117.5	(90.9 to 150.6)
		Female	1990	2,714	(1,915 to 3,215)	163.9	(116.3 to 197.4)
			2013	4,617	(3,297 to 5,966)	191.6	(137.2 to 247.3)
		Both	1990	4,051	(3,179 to 4,639)	130.9	(101.6 to 151.5)
			2013	7,018	(5,550 to 8,548)	157.9	(125.9 to 191.8)
	DALYs	Male	1990	37,546	(29,988 to 45,243)	1738.4	(1390.4 to 2084.2)
			2013	66,336	(49,512 to 85,150)	2111.7	(1615.0 to 2726.8)
		Female	1990	70,232	(48,646 to 84,755)	3115.1	(2207.6 to 3710.8)

			2013	116,383	(80,681 to 149,732)	3516	(2463.4 to 4529.0)
		Both	1990	107,778	(83,817 to 124,815)	2478.4	(1931.5 to 2852.5)
			2013	182,720	(141,892 to 221,174)	2869.4	(2265.7 to 3509.3)
	YLDs	Male	1990	381	(259 to 530)	20.2	(13.5 to 28.0)
			2013	559	(369 to 802)	18.7	(12.3 to 26.5)
		Female	1990	319	(217 to 449)	14.8	(10.1 to 20.8)
			2013	506	(325 to 711)	15	(9.7 to 21.0)
		Both	1990	701	(491 to 935)	17.3	(12.1 to 22.9)
			2013	1,066	(733 to 1,452)	16.7	(11.3 to 22.8)
Malta	Deaths	Male	1990	115	(96 to 144)	86.5	(71.9 to 107.7)
			2013	129	(101 to 159)	51	(40.2 to 62.8)
		Female	1990	159	(125 to 184)	80.9	(63.9 to 94.2)
			2013	175	(133 to 213)	49.1	(37.4 to 59.8)
		Both	1990	275	(241 to 312)	83.6	(73.2 to 94.8)
							Uncertainty
Country							
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Country	measure	sex	year 2013	number 304	Uncertainty Interval (255 to 352)	Rate 50.5	(42.0 to 58.4)
Country	DALYs	sex Male			· · · · · · · · · · · · · · · · · · ·		
Country			2013	304	(255 to 352)	50.5	(42.0 to 58.4)
Country			2013 1990	304 1,945	(255 to 352) (1,647 to 2,367)	50.5 1318.8	(42.0 to 58.4) (1121.6 to 1605.5)
Country		Male	2013 1990 2013	304 1,945 2,113	(255 to 352) (1,647 to 2,367) (1,697 to 2,566)	50.5 1318.8 731.8	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4)
Country		Male	2013 1990 2013 1990	304 1,945 2,113 2,146	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443)	50.5 1318.8 731.8 1095.3	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3)
Country		Male Female	2013 1990 2013 1990 2013	304 1,945 2,113 2,146 1,982	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426)	50.5 1318.8 731.8 1095.3 557.3	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8)
Country		Male Female	2013 1990 2013 1990 2013 1990	304 1,945 2,113 2,146 1,982 4,091	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426) (3,641 to 4,594)	50.5 1318.8 731.8 1095.3 557.3 1194	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8) (1061.2 to 1341.4)
Country	DALYs	Male Female Both	2013 1990 2013 1990 2013 1990 2013	304 1,945 2,113 2,146 1,982 4,091 4,096	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426) (3,641 to 4,594) (3,493 to 4,727)	50.5 1318.8 731.8 1095.3 557.3 1194 641.1	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8) (1061.2 to 1341.4) (545.3 to 736.3)
Country	DALYs	Male Female Both	2013 1990 2013 1990 2013 1990 2013 1990	304 1,945 2,113 2,146 1,982 4,091 4,096 149	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426) (3,641 to 4,594) (3,493 to 4,727) (97 to 200)	50.5 1318.8 731.8 1095.3 557.3 1194 641.1 99.5	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8) (1061.2 to 1341.4) (545.3 to 736.3) (65.1 to 133.7)
Country	DALYs	Male Female Both Male	2013 1990 2013 1990 2013 1990 2013 1990 2013	304 1,945 2,113 2,146 1,982 4,091 4,096 149 324	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426) (3,641 to 4,594) (3,493 to 4,727) (97 to 200) (203 to 467)	50.5 1318.8 731.8 1095.3 557.3 1194 641.1 99.5 105.7	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8) (1061.2 to 1341.4) (545.3 to 736.3) (65.1 to 133.7) (66.2 to 151.8)
Country	DALYs	Male Female Both Male	2013 1990 2013 1990 2013 1990 2013 1990 2013	304 1,945 2,113 2,146 1,982 4,091 4,096 149 324 142	(255 to 352) (1,647 to 2,367) (1,697 to 2,566) (1,736 to 2,443) (1,616 to 2,426) (3,641 to 4,594) (3,493 to 4,727) (97 to 200) (203 to 467) (99 to 198)	50.5 1318.8 731.8 1095.3 557.3 1194 641.1 99.5 105.7 72.6	(42.0 to 58.4) (1121.6 to 1605.5) (588.5 to 883.4) (885.9 to 1246.3) (457.3 to 681.8) (1061.2 to 1341.4) (545.3 to 736.3) (65.1 to 133.7) (66.2 to 151.8) (50.4 to 101.1)

Marshall							
Islands	Deaths	Male	1990	15	(10 to 20)	219	(141.1 to 274.2)
			2013	28	(18 to 40)	209.5	(139.4 to 277.7)
		Female	1990	7	(5 to 10)	109.2	(82.7 to 149.5)
			2013	19	(10 to 27)	131.2	(77.2 to 185.5)
		Both	1990	23	(18 to 28)	159.7	(126.0 to 189.1)
			2013	48	(32 to 63)	166.8	(114.6 to 211.4)
	DALYs	Male	1990	472	(328 to 610)	4457.8	(2921.6 to 5690.5)
			2013	821	(530 to 1,200)	4279.1	(2813.0 to 5960.0)
		Female	1990	203	(151 to 266)	2005.7	(1496.3 to 2640.0)
			2013	459	(241 to 679)	2408.8	(1333.6 to 3449.3)
		Both	1990	676	(519 to 816)	3191	(2459.8 to 3846.1)
			2013	1,281	(849 to 1,744)	3302.1	(2228.9 to 4358.0)
	YLDs	Male	1990	4	(3 to 6)	61.4	(41.8 to 86.1)
			2013	8	(5 to 11)	53.8	(35.8 to 73.5)
		Female	1990	3	(2 to 4)	41.5	(27.9 to 56.3)
			2013	6	(4 to 9)	40.3	(27.0 to 57.3)
		Both	1990	8	(5 to 11)	50.6	(35.4 to 68.0)
			2013	15	(10 to 20)	46.5	(32.5 to 61.2)
Mauritania	Deaths	Male	1990	346	(268 to 421)	119.6	(92.9 to 145.5)
			2013	883	(679 to 1,113)	153.6	(122.1 to 187.3)
		Female	1990	515	(412 to 662)	151.4	(122.1 to 195.8)
			2013	1,157	(832 to 1,591)	170.3	(123.9 to 233.1)
		Both	1990	862	(742 to 1,026)	137.7	(118.1 to 164.8)
			2013	2,040	(1,664 to 2,534)	163.6	(134.7 to 201.7)
	DALYs	Male	1990	8,891	(6,872 to 10,893)	2149.1	(1663.1 to 2613.0)
			2013	21,695	(16,112 to 28,493)	2664.1	(2050.6 to 3352.0)
		Female	1990	11,622	(9,067 to 14,824)	2584	(2056.4 to 3315.5)
			2013	24,091	(17,041 to 34,095)	2762.6	(1989.9 to 3823.1)

		Both	1990	20,513	(17,460 to 24,165)	2395.3	(2059.2 to 2843.2)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
			2013	45,787	(36,173 to 57,207)	2731	(2217.7 to 3389.8)
	YLDs	Male	1990	85	(56 to 119)	19.6	(13.2 to 27.0)
			2013	191	(123 to 269)	20.5	(13.3 to 29.0)
		Female	1990	73	(47 to 104)	14.7	(9.5 to 20.9)
			2013	154	(101 to 217)	14.8	(9.8 to 20.7)
		Both	1990	159	(108 to 214)	16.9	(11.7 to 22.7)
			2013	345	(229 to 478)	17.4	(11.8 to 23.9)
Mauritius	Deaths	Male	1990	515	(448 to 605)	251.7	(218.1 to 293.1)
			2013	733	(579 to 844)	155.7	(124.3 to 178.8)
		Female	1990	439	(365 to 504)	158	(133.5 to 184.0)
			2013	553	(458 to 760)	89.6	(74.1 to 121.5)
		Both	1990	954	(866 to 1,072)	196.7	(178.8 to 221.5)
			2013	1,286	(1,137 to 1,502)	118.4	(104.7 to 138.5)
	DALYs	Male	1990	12,221	(10,613 to 14,475)	4558.1	(3981.1 to 5334.6)
			2013	15,370	(12,213 to 17,875)	2700.2	(2141.5 to 3122.0)
		Female	1990	8,235	(6,811 to 9,380)	2491.9	(2059.1 to 2834.2)
			2013	8,463	(7,004 to 11,929)	1273.8	(1057.8 to 1792.8)
		Both	1990	20,457	(18,545 to 22,956)	3387.6	(3084.4 to 3790.6)
			2013	23,833	(20,808 to 27,739)	1921.1	(1686.7 to 2236.1)
	YLDs	Male	1990	148	(98 to 207)	57.4	(37.6 to 79.6)
			2013	340	(235 to 473)	61.6	(42.0 to 85.7)
		Female	1990	151	(102 to 209)	46	(31.4 to 63.6)
			2013	333	(227 to 452)	49.4	(33.6 to 67.2)
		Both	1990	300	(209 to 406)	50.8	(35.7 to 68.3)
			2013	673	(473 to 890)	54.8	(38.7 to 72.1)
Mexico	Deaths	Male	1990	10,399	(8,779 to 12,005)	60.5	(50.7 to 70.0)

			2013	19,386	(16,443 to 23,716)	54.7	(46.3 to 67.5)
		Female	1990	12,210	(9,763 to 14,472)	57.3	(45.6 to 67.9)
			2013	21,332	(17,312 to 26,536)	45.7	(37.0 to 56.9)
		Both	1990	22,609	(19,911 to 25,260)	58.7	(51.5 to 66.0)
			2013	40,719	(36,791 to 47,691)	49.8	(44.9 to 58.3)
	DALYs	Male	1990	231,431	(198,098 to 267,910)	1052.9	(896.4 to 1219.8)
			2013	374,777	(319,833 to 455,876)	914.5	(783.1 to 1118.1)
		Female	1990	240,799	(194,061 to 285,285)	949.9	(764.6 to 1122.1)
			2013	349,050	(289,812 to 448,323)	695.6	(576.3 to 888.0)
		Both	1990	472,231	(419,185 to 524,405)	995.7	(886.6 to 1107.7)
			2013	723,827	(655,880 to 848,261)	794.8	(721.5 to 931.4)
	YLDs	Male	1990	5,584	(3,940 to 7,314)	26.3	(18.6 to 34.3)
			2013	11,738	(8,246 to 15,343)	28	(19.9 to 36.5)
		Female	1990	4,399	(3,082 to 5,708)	18.1	(12.7 to 23.4)
			2013	9,322	(6,497 to 12,174)	18.2	(12.7 to 23.7)
		Both	1990	9,983	(7,001 to 13,009)	21.9	(15.4 to 28.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	21,060	(14,827 to 27,480)	22.6	(15.9 to 29.4)
Moldova	Deaths	Male	1990	2,544	(2,186 to 2,959)	189.7	(162.6 to 220.9)
			2013	2,837	(2,441 to 3,294)	184.3	(157.5 to 213.7)
		Female	1990	3,649	(3,011 to 4,095)	170.4	(140.4 to 190.2)
			2013	3,217	(2,690 to 3,719)	124.8	(104.4 to 144.1)
		Both	1990	6,194	(5,476 to 6,739)	178.5	(156.0 to 194.1)
			2013	6,054	(5,466 to 6,666)	148.6	(133.8 to 164.1)
	DALYs	Male	1990	53,190	(45,763 to 61,300)	3375.4	(2920.3 to 3902.1)
			2013	57,094	(49,189 to 66,292)	3251.8	(2807.0 to 3761.5)
		Female	1990	64,736	(54,382 to 73,623)	2813.5	(2363.3 to 3213.7)
			2013	48,639	(40,983 to 57,550)	1910.1	(1609.0 to 2267.7)

			2013	752	(622 to 886)	231.7	(191.8 to 272.0)
Montenegro	Deaths	Male	1990	529	(424 to 649)	247.8	(199.1 to 301.6)
			2013	260	(176 to 352)	13.9	(9.4 to 19.0)
		Both	1990	174	(119 to 231)	14.8	(10.2 to 19.7)
			2013	119	(77 to 170)	12.3	(7.9 to 17.9)
		Female	1990	75	(49 to 106)	12.4	(8.2 to 17.4)
			2013	141	(87 to 199)	15.8	(9.8 to 22.6)
	YLDs	Male	1990	98	(66 to 135)	17.5	(12.0 to 24.0)
			2013	92,583	(79,737 to 107,258)	5228.8	(4548.3 to 5942.8)
		Both	1990	53,351	(47,562 to 60,078)	4556.9	(4063.6 to 5155.9)
			2013	42,925	(33,264 to 53,021)	4522	(3517.5 to 5464.3)
		Female	1990	26,426	(22,750 to 31,783)	4271.1	(3671.1 to 5203.5)
			2013	49,658	(39,931 to 61,993)	6071	(5045.0 to 7365.0)
	DALYs	Male	1990	26,924	(22,228 to 31,321)	4877.1	(4013.6 to 5639.3)
			2013	3,612	(3,144 to 4,112)	272	(233.3 to 306.3)
		Both	1990	2,168	(1,936 to 2,458)	235.3	(210.2 to 267.2)
			2013	1,792	(1,397 to 2,158)	245.7	(191.4 to 290.6)
		Female	1990	1,117	(962 to 1,354)	224.9	(192.9 to 274.5)
			2013	1,819	(1,506 to 2,221)	303	(256.3 to 360.3)
Mongolia	Deaths	Male	1990	1,051	(870 to 1,218)	247	(206.8 to 286.1)
			2013	1,439	(955 to 1,975)	33.7	(22.2 to 46.1)
		Both	1990	1,316	(878 to 1,832)	32.2	(21.4 to 44.9)
			2013	636	(405 to 904)	26.2	(16.7 to 36.9)
		Female	1990	594	(358 to 894)	25.4	(15.4 to 38.1)
			2013	802	(515 to 1,146)	43.4	(28.0 to 61.3)
	YLDs	Male	1990	722	(473 to 1,023)	41.2	(27.1 to 58.4)
			2013	105,733	(96,171 to 117,354)	2476.5	(2252.7 to 2745.9)
		Both	1990	117,926	(105,278 to 129,260)	3047.1	(2714.7 to 3335.4)

		Female	1990	767	(620 to 941)	229.3	(184.6 to 282.1)
			2013	1,028	(853 to 1,176)	227.7	(188.5 to 261.2)
		Both	1990	1,297	(1,114 to 1,504)	238.5	(205.4 to 276.7)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	1,780	(1,563 to 1,984)	230.9	(201.9 to 257.0)
	DALYs	Male	1990	10,404	(8,248 to 12,967)	4266.2	(3403.8 to 5270.0)
			2013	12,323	(10,065 to 14,605)	3517.6	(2890.7 to 4174.1)
		Female	1990	10,834	(8,274 to 13,476)	3423.5	(2614.5 to 4266.3)
			2013	13,432	(11,184 to 15,563)	3007.3	(2522.5 to 3493.3)
		Both	1990	21,238	(17,899 to 24,531)	3812.8	(3217.1 to 4415.9)
			2013	25,756	(22,431 to 29,159)	3253.1	(2844.6 to 3671.9)
	YLDs	Male	1990	75	(51 to 106)	29.2	(19.6 to 41.2)
			2013	113	(72 to 160)	30.9	(19.8 to 43.5)
		Female	1990	60	(39 to 83)	19	(12.5 to 26.4)
			2013	80	(51 to 113)	18.8	(12.1 to 26.3)
		Both	1990	135	(93 to 180)	23.5	(16.3 to 31.3)
			2013	193	(131 to 266)	24.4	(16.7 to 33.3)
Morocco	Deaths	Male	1990	4,195	(3,418 to 5,130)	106.4	(86.5 to 129.6)
			2013	5,574	(4,359 to 7,429)	71.5	(55.8 to 92.9)
		Female	1990	5,310	(4,270 to 6,170)	112.5	(88.1 to 131.5)
			2013	7,988	(6,126 to 10,658)	79.8	(60.6 to 106.6)
		Both	1990	9,506	(8,228 to 10,725)	110.2	(92.8 to 124.2)
			2013	13,563	(11,340 to 16,747)	76.4	(63.9 to 94.0)
	DALYs	Male	1990	123,550	(98,622 to 154,666)	1905.7	(1556.9 to 2339.9)
			2013	126,103	(97,267 to 170,035)	1204.7	(939.6 to 1612.0)
		Female	1990	131,741	(105,954 to 162,773)	1882.5	(1527.7 to 2188.7)
			2013	151,748	(115,384 to 203,480)	1271.7	(968.0 to 1710.9)
		Both	1990	255,291	(220,434 to 298,168)	1897.4	(1642.8 to 2141.2)

			2013	277,851	(228,842 to 343,127)	1244	(1039.6 to 1534.7)
	YLDs	Male	1990	591	(373 to 837)	10.3	(6.4 to 14.7)
			2013	1,222	(783 to 1,744)	10.8	(7.0 to 15.4)
		Female	1990	597	(387 to 840)	9	(5.9 to 12.5)
			2013	1,194	(747 to 1,667)	9.4	(5.9 to 13.0)
		Both	1990	1,189	(791 to 1,621)	9.6	(6.4 to 13.2)
			2013	2,417	(1,621 to 3,259)	10	(6.7 to 13.4)
Mozambique	Deaths	Male	1990	1,930	(1,424 to 2,403)	90.8	(66.7 to 113.3)
			2013	5,112	(3,883 to 6,614)	130.6	(100.7 to 168.6)
		Female	1990	2,328	(1,679 to 2,959)	92.6	(65.3 to 118.8)
			2013	4,899	(3,573 to 6,275)	105.4	(76.0 to 134.9)
		Both	1990	4,258	(3,407 to 5,066)	92.6	(72.8 to 111.7)
			2013	10,011	(8,100 to 12,099)	117	(93.7 to 142.1)
	DALYs	Male	1990	50,061	(36,848 to 62,652)	1803.4	(1333.8 to 2249.3)
			2013	120,353	(91,639 to 156,383)	2495.8	(1907.3 to 3226.1)
		Female	1990	50,271	(36,043 to 63,993)	1554.1	(1125.9 to 1973.3)
			2013	91,954	(68,019 to 117,463)	1634.8	(1199.3 to 2096.2)
		Both	1990	100,332	(79,478 to 119,164)	1670.8	(1331.3 to 1990.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	212,308	(174,432 to 257,217)	2023.2	(1648.1 to 2448.1)
	YLDs	Male	1990	388	(259 to 544)	12.9	(8.5 to 17.8)
			2013	785	(488 to 1,164)	13.7	(8.6 to 20.2)
		Female	1990	337	(213 to 486)	9.1	(5.7 to 13.1)
			2013	661	(407 to 955)	9.6	(6.0 to 13.9)
		Both	1990	725	(489 to 992)	10.7	(7.2 to 14.8)
			2013	1,446	(951 to 2,042)	11.4	(7.6 to 15.9)
Myanmar	Deaths	Male	1990	19,818	(12,959 to 28,520)	248.9	(172.4 to 342.4)
			2013	27,881	(17,852 to 39,638)	213.5	(146.5 to 285.9)

		Female	1990	21,659	(13,316 to 33,203)	232	(150.6 to 342.6)
			2013	28,585	(17,956 to 41,501)	182.3	(120.3 to 255.5)
		Both	1990	41,478	(27,235 to 59,616)	241	(166.5 to 334.8)
			2013	56,467	(36,157 to 78,944)	196.8	(134.8 to 264.1)
	DALYs	Male	1990	491,968	(314,423 to 720,548)	4698.9	(3045.2 to 6779.1)
			2013	632,649	(375,906 to 962,469)	3756.5	(2375.5 to 5368.9)
		Female	1990	425,349	(245,042 to 697,593)	3799.4	(2283.3 to 5967.7)
			2013	512,382	(306,053 to 790,897)	2734.2	(1701.5 to 4058.0)
		Both	1990	917,318	(581,331 to 1,368,588)	4237.3	(2759.7 to 6205.8)
			2013	1,145,032	(691,108 to 1,699,167)	3207.3	(2015.9 to 4572.0)
	YLDs	Male	1990	5,223	(3,368 to 7,213)	54.4	(35.0 to 75.8)
			2013	9,667	(6,346 to 13,524)	60.4	(40.7 to 85.0)
		Female	1990	4,924	(3,251 to 6,903)	42.9	(28.9 to 60.7)
			2013	8,866	(6,111 to 12,188)	45.4	(31.0 to 62.1)
		Both	1990	10,147	(7,079 to 13,522)	48	(33.7 to 64.3)
			2013	18,534	(13,073 to 24,639)	52	(36.9 to 68.6)
Namibia	Deaths	Male	1990	321	(256 to 394)	154.8	(122.9 to 190.5)
			2013	597	(478 to 743)	161.6	(130.2 to 196.5)
		Female	1990	436	(346 to 517)	166.6	(133.2 to 197.9)
			2013	734	(590 to 954)	154	(125.1 to 199.0)
		Both	1990	758	(635 to 865)	162.5	(136.4 to 185.1)
			2013	1,332	(1,127 to 1,588)	158.1	(134.1 to 187.8)
	DALYs	Male	1990	7,532	(5,945 to 9,293)	2786.3	(2218.6 to 3425.3)
			2013	14,527	(11,156 to 18,355)	2995.5	(2379.6 to 3741.6)
		Female	1990	8,750	(6,735 to 10,538)	2757.2	(2167.0 to 3287.1)
			2013	14,373	(11,174 to 19,222)	2455.3	(1947.2 to 3217.3)
		Both	1990	16,283	(13,473 to 18,942)	2781.7	(2327.8 to 3189.0)
			2013	28,900	(24,010 to 34,880)	2700	(2274.9 to 3236.3)

	YLDs	Male	1990	74	(48 to 103)	24.5	(15.9 to 34.5)
			2013	130	(84 to 183)	24.1	(15.8 to 34.0)
		Female	1990	79	(51 to 110)	22.2	(14.4 to 31.4)
			2013	146	(97 to 212)	21.4	(14.0 to 31.2)
		Both	1990	153	(105 to 207)	23.3	(16.0 to 31.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	276	(189 to 378)	22.5	(15.2 to 30.9)
Nepal	Deaths	Male	1990	3,163	(2,295 to 4,126)	104.5	(76.2 to 134.8)
			2013	7,774	(5,745 to 10,025)	127.2	(93.6 to 162.5)
		Female	1990	2,938	(2,006 to 4,713)	89.7	(60.1 to 140.4)
			2013	8,322	(5,494 to 11,025)	125	(83.0 to 163.0)
		Both	1990	6,101	(4,900 to 7,826)	96.6	(77.2 to 121.6)
			2013	16,097	(12,249 to 19,583)	126	(96.8 to 152.5)
	DALYs	Male	1990	79,305	(58,283 to 104,099)	1857.4	(1356.4 to 2420.1)
			2013	158,217	(116,624 to 205,185)	2030.4	(1507.0 to 2600.6)
		Female	1990	66,029	(45,670 to 107,800)	1513.7	(1052.6 to 2425.8)
			2013	152,903	(98,498 to 206,526)	1890.6	(1252.2 to 2524.5)
		Both	1990	145,335	(117,284 to 188,869)	1678.4	(1355.9 to 2155.7)
			2013	311,121	(237,131 to 383,394)	1956	(1489.3 to 2387.3)
	YLDs	Male	1990	1,911	(1,259 to 2,733)	41.6	(26.7 to 59.9)
			2013	3,841	(2,549 to 5,376)	43.2	(28.7 to 60.1)
		Female	1990	1,742	(1,116 to 2,416)	35.3	(22.5 to 48.8)
			2013	3,272	(2,164 to 4,555)	34.5	(22.8 to 47.9)
		Both	1990	3,653	(2,441 to 5,037)	38.3	(25.6 to 52.6)
			2013	7,113	(4,993 to 9,526)	38.7	(26.8 to 51.4)
Netherlands	Deaths	Male	1990	5,167	(4,286 to 6,269)	74.1	(61.1 to 90.1)
			2013	4,546	(3,522 to 5,582)	38.4	(29.7 to 47.1)
		Female	1990	7,542	(5,888 to 8,571)	61.2	(47.9 to 69.7)

			2013	7,337	(5,898 to 10,206)	40.1	(32.5 to 55.2)
		Both	1990	12,709	(10,984 to 14,357)	66.7	(57.7 to 75.4)
			2013	11,884	(9,822 to 14,487)	39.8	(33.0 to 48.2)
	DALYs	Male	1990	83,402	(70,753 to 100,459)	1120.2	(949.1 to 1342.9)
			2013	66,800	(53,565 to 79,748)	557.3	(447.2 to 666.8)
		Female	1990	90,201	(72,528 to 103,044)	824.9	(668.6 to 958.3)
			2013	79,081	(65,660 to 104,815)	514.3	(423.6 to 680.9)
		Both	1990	173,604	(154,230 to 196,034)	949.1	(846.3 to 1073.4)
			2013	145,882	(124,352 to 171,203)	535	(455.3 to 623.7)
	YLDs	Male	1990	6,004	(4,068 to 8,272)	79	(53.3 to 108.5)
			2013	8,790	(5,631 to 12,463)	72.7	(46.9 to 102.4)
		Female	1990	6,387	(4,295 to 9,103)	61.1	(41.2 to 86.2)
			2013	8,105	(5,474 to 11,035)	56.3	(38.1 to 76.2)
		Both	1990	12,391	(8,393 to 16,869)	68.6	(46.5 to 93.2)
			2013	16,895	(11,273 to 22,788)	63.9	(42.8 to 86.2)
New Zealand	Deaths	Male	1990	1,067	(894 to 1,306)	76.4	(63.6 to 93.3)
			2013	1,131	(868 to 1,415)	38.5	(29.5 to 48.0)
		Female	1990	1,609	(1,320 to 1,972)	74.1	(61.0 to 90.8)
			2013	1,768	(1,306 to 2,133)	44.5	(33.2 to 53.2)
		Both	1990	2,677	(2,395 to 3,104)	75.7	(67.4 to 87.9)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	2,900	(2,336 to 3,349)	42.1	(34.2 to 48.5)
	DALYs	Male	1990	17,892	(15,229 to 21,603)	1170.7	(995.9 to 1414.2)
			2013	15,050	(11,930 to 18,303)	533.4	(422.7 to 647.6)
		Female	1990	20,636	(17,414 to 24,817)	1016.5	(860.4 to 1220.9)
			2013	17,682	(13,629 to 20,807)	513.5	(396.4 to 604.4)
		Both	1990	38,528	(34,944 to 44,194)	1085.7	(983.7 to 1244.6)
			2013	32,733	(27,590 to 37,014)	524.7	(442.3 to 593.3)

			2013	2,584	(1,216 to 3,742)	111	(55.1 to 159.2)
		Female	1990	1,319	(973 to 1,743)	114	(84.6 to 148.4)
			2013	1,845	(1,282 to 2,530)	79.8	(56.2 to 109.2)
Niger	Deaths	Male	1990	1,059	(827 to 1,326)	98.9	(75.3 to 122.9)
			2013	782	(550 to 1,054)	21.3	(15.0 to 28.7)
		Both	1990	314	(219 to 413)	17.3	(12.3 to 22.8)
			2013	353	(239 to 498)	18.3	(12.1 to 26.0)
		Female	1990	153	(101 to 211)	16.3	(10.8 to 22.4)
			2013	429	(292 to 595)	24.6	(16.7 to 34.3)
	YLDs	Male	1990	161	(112 to 219)	18.4	(13.0 to 25.2)
			2013	33,260	(29,398 to 38,682)	964.5	(852.9 to 1124.5)
		Both	1990	27,228	(22,617 to 30,055)	1563.3	(1306.5 to 1723.4)
			2013	15,500	(13,037 to 19,894)	841.1	(709.7 to 1082.0)
		Female	1990	14,452	(10,493 to 16,216)	1558.5	(1137.0 to 1741.4)
			2013	17,760	(14,530 to 21,164)	1103.9	(908.7 to 1317.1)
	DALYs	Male	1990	12,775	(10,733 to 15,090)	1571.2	(1302.8 to 1839.7)
			2013	2,049	(1,811 to 2,402)	66.5	(58.9 to 78.2)
		Both	1990	1,230	(1,024 to 1,354)	98.1	(80.7 to 108.3)
			2013	1,038	(880 to 1,328)	61	(51.7 to 78.0)
		Female	1990	687	(504 to 772)	98.7	(72.5 to 111.1)
			2013	1,011	(822 to 1,233)	72.8	(59.2 to 89.1)
Nicaragua	Deaths	Male	1990	542	(448 to 634)	96.9	(78.9 to 113.6)
			2013	2,901	(1,981 to 3,919)	49.1	(33.5 to 66.1)
		Both	1990	2,340	(1,653 to 3,052)	66.1	(46.8 to 86.2)
			2013	1,255	(838 to 1,730)	40.1	(27.0 to 55.2)
		Female	1990	1,087	(756 to 1,433)	54.6	(38.1 to 71.9)
			2013	1,645	(1,103 to 2,298)	59.3	(39.9 to 82.6)
	YLDs	Male	1990	1,252	(872 to 1,670)	80.3	(56.2 to 106.8)

		Both	1990	2,379	(1,926 to 2,814)	107.5	(87.3 to 127.9)
			2013	4,430	(2,874 to 5,792)	96.4	(62.8 to 125.2)
	DALYs	Male	1990	37,070	(28,619 to 48,990)	1915.2	(1488.8 to 2371.8)
			2013	48,064	(33,702 to 68,246)	1369.8	(958.8 to 1877.3)
		Female	1990	41,712	(29,756 to 56,625)	2076.3	(1537.8 to 2704.3)
			2013	60,364	(27,619 to 88,865)	1815.5	(851.9 to 2614.4)
		Both	1990	78,783	(62,757 to 97,472)	2005.9	(1627.6 to 2387.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	108,429	(68,680 to 143,681)	1599.5	(1042.8 to 2093.2)
	YLDs	Male	1990	308	(200 to 437)	19.6	(12.9 to 28.1)
			2013	740	(469 to 1,039)	19.1	(12.3 to 26.8)
		Female	1990	251	(164 to 355)	14.4	(9.5 to 20.0)
			2013	547	(359 to 754)	14.5	(9.7 to 20.3)
		Both	1990	560	(377 to 766)	16.8	(11.4 to 23.0)
			2013	1,287	(854 to 1,727)	16.8	(11.1 to 22.4)
Nigeria	Deaths	Male	1990	13,319	(10,594 to 16,401)	105.8	(83.8 to 126.8)
			2013	25,266	(19,537 to 32,558)	119.8	(95.1 to 148.5)
		Female	1990	18,758	(15,417 to 23,048)	134	(107.9 to 162.8)
			2013	30,149	(23,757 to 39,336)	139.4	(109.9 to 176.4)
		Both	1990	32,077	(27,593 to 36,926)	121.5	(103.2 to 140.6)
			2013	55,415	(46,776 to 66,946)	130.3	(111.1 to 153.8)
	DALYs	Male	1990	365,687	(287,127 to 450,793)	1753.8	(1387.4 to 2137.6)
			2013	629,801	(469,305 to 823,916)	1892.9	(1480.3 to 2427.4)
		Female	1990	455,988	(368,854 to 603,526)	2152.1	(1772.4 to 2647.9)
			2013	686,540	(532,025 to 946,477)	2090.7	(1653.3 to 2713.6)
		Both	1990	821,675	(706,415 to 973,879)	1971.3	(1681.4 to 2271.6)
					(1,086,684 to		
			2013	1,316,341	1,637,082)	1998.2	(1691.3 to 2402.7)

	YLDs	Male	1990	4,258	(2,790 to 6,128)	19.5	(12.6 to 27.6)
			2013	7,429	(4,572 to 10,870)	18.5	(11.3 to 27.8)
		Female	1990	3,339	(2,192 to 4,569)	14.3	(9.5 to 19.8)
			2013	5,890	(3,879 to 8,463)	14.6	(9.6 to 21.1)
		Both	1990	7,598	(5,269 to 10,153)	16.8	(11.8 to 22.2)
			2013	13,319	(8,634 to 18,750)	16.5	(10.8 to 23.5)
North Korea	Deaths	Male	1990	7,969	(5,320 to 11,355)	241.2	(175.6 to 320.0)
			2013	19,242	(14,582 to 23,756)	258.5	(202.5 to 313.3)
		Female	1990	12,935	(9,104 to 18,023)	210.7	(152.3 to 283.8)
			2013	25,833	(19,976 to 31,867)	196.1	(152.7 to 241.9)
		Both	1990	20,904	(15,997 to 26,795)	221.3	(173.0 to 275.9)
			2013	45,075	(38,317 to 53,206)	219.9	(187.6 to 259.0)
	DALYs	Male	1990	227,930	(150,317 to 336,162)	4413.6	(3045.5 to 6135.4)
			2013	466,318	(349,147 to 591,456)	4676.5	(3609.8 to 5772.2)
		Female	1990	271,032	(178,684 to 388,092)	3591.6	(2471.7 to 5047.9)
			2013	446,528	(340,151 to 560,238)	3154.2	(2405.8 to 3952.2)
		Both	1990	498,963	(375,186 to 656,316)	3916.7	(3010.1 to 5026.5)
			2013	912,846	(762,536 to 1,088,037)	3797.8	(3211.0 to 4510.3)
	YLDs	Male	1990	3,020	(1,963 to 4,282)	73.3	(47.5 to 106.4)
			2013	6,760	(4,532 to 9,411)	76.7	(51.0 to 106.7)
		Female	1990	3,305	(2,243 to 4,529)	45.3	(31.0 to 61.3)
			2013	6,430	(4,307 to 8,750)	45.6	(30.8 to 61.7)
		Both	1990	6,325	(4,322 to 8,656)	53.9	(36.9 to 73.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	13,190	(9,052 to 17,785)	56.2	(38.8 to 75.7)
Norway	Deaths	Male	1990	2,225	(1,838 to 2,783)	85.6	(70.7 to 106.7)
			2013	1,733	(1,282 to 2,097)	49.5	(36.7 to 59.6)
		Female	1990	3,227	(2,515 to 3,671)	75.1	(59.2 to 85.6)

			2013	2,286	(1,871 to 3,190)	40.7	(33.8 to 57.4)
		Both	1990	5,452	(4,768 to 6,112)	80.1	(70.1 to 89.4)
			2013	4,019	(3,449 to 4,993)	44.8	(38.5 to 55.2)
	DALYs	Male	1990	33,432	(28,117 to 42,034)	1283.5	(1080.3 to 1615.5)
			2013	23,430	(17,539 to 27,732)	687.3	(514.2 to 813.5)
		Female	1990	36,568	(29,240 to 41,466)	977.7	(781.9 to 1126.1)
			2013	21,540	(18,027 to 29,672)	470.3	(395.2 to 646.6)
		Both	1990	70,000	(62,562 to 78,721)	1114.8	(999.7 to 1261.1)
			2013	44,971	(38,690 to 53,506)	571.1	(491.7 to 678.2)
	YLDs	Male	1990	2,521	(1,679 to 3,478)	97.2	(65.3 to 134.0)
			2013	3,168	(2,066 to 4,434)	93.8	(61.0 to 131.4)
		Female	1990	2,600	(1,717 to 3,698)	74.1	(49.2 to 104.7)
			2013	2,698	(1,771 to 3,805)	68.1	(45.4 to 94.7)
		Both	1990	5,121	(3,557 to 6,871)	84.2	(58.4 to 112.1)
			2013	5,866	(3,957 to 8,021)	80.2	(53.9 to 109.3)
Oman	Deaths	Male	1990	221	(151 to 308)	103	(73.9 to 138.8)
			2013	529	(385 to 668)	96.2	(71.8 to 118.7)
		Female	1990	283	(202 to 411)	104.9	(75.8 to 149.6)
			2013	414	(288 to 528)	86	(60.4 to 108.5)
		Both	1990	504	(392 to 666)	103.2	(81.5 to 132.0)
			2013	943	(756 to 1,130)	91.4	(74.0 to 108.8)
	DALYs	Male	1990	6,569	(4,435 to 9,662)	1684.6	(1157.3 to 2336.8)
			2013	10,709	(7,558 to 14,238)	1438.3	(1037.4 to 1840.4)
		Female	1990	8,767	(5,699 to 13,224)	1916.7	(1347.6 to 2805.6)
			2013	7,499	(5,149 to 9,829)	1244.9	(858.9 to 1603.7)
		Both	1990	15,337	(11,428 to 21,185)	1779.1	(1378.0 to 2347.8)
			2013	18,209	(14,135 to 22,416)	1346.7	(1070.1 to 1628.2)
	YLDs	Male	1990	43	(28 to 62)	10.6	(6.9 to 15.0)

			2013	113	(74 to 161)	10.5	(6.9 to 15.1)
		Female	1990	29	(19 to 41)	8.6	(5.6 to 11.9)
			2013	59	(40 to 81)	8.7	(6.0 to 11.9)
		Both	1990	73	(49 to 99)	9.6	(6.5 to 12.8)
			2013	172	(118 to 238)	9.7	(6.8 to 13.3)
Pakistan	Deaths	Male	1990	22,608	(16,836 to 29,202)	100.3	(74.4 to 128.7)
			2013	57,255	(41,474 to 74,014)	141.9	(105.3 to 180.1)
		Female	1990	28,693	(20,081 to 35,313)	154.4	(108.4 to 190.5)
			2013	61,288	(39,423 to 86,657)	164.3	(107.7 to 227.4)
		Both	1990	51,302	(38,975 to 61,313)	124.7	(94.6 to 148.9)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	118,544	(89,238 to 147,413)	152.7	(115.2 to 187.5)
	DALYs	Male	1990	491,740	(369,609 to 639,686)	1697.3	(1265.9 to 2190.9)
			2013	1,203,732	(829,626 to 1,619,971)	2389.7	(1690.3 to 3129.6)
		Female	1990	597,171	(423,690 to 742,821)	2546.9	(1781.0 to 3146.9)
			2013	1,196,097	(768,952 to 1,727,168)	2592.8	(1666.2 to 3693.6)
		Both	1990	1,088,911	(842,868 to 1,303,849)	2082.3	(1594.2 to 2491.0)
			2013	2,399,830	(1,819,573 to 3,051,319)	2488.1	(1878.7 to 3128.0)
	YLDs	Male	1990	13,104	(8,506 to 17,936)	42.4	(28.0 to 57.6)
			2013	29,125	(19,381 to 39,989)	54.9	(36.8 to 75.6)
		Female	1990	8,559	(5,556 to 11,845)	32.2	(20.6 to 44.6)
			2013	16,127	(10,371 to 22,809)	29.9	(19.1 to 42.8)
		Both	1990	21,663	(14,749 to 28,869)	37.7	(25.4 to 50.0)
			2013	45,253	(31,560 to 61,131)	42.7	(30.0 to 57.6)
Palestine	Deaths	Male	1990	350	(250 to 468)	145.5	(105.5 to 193.1)
			2013	802	(569 to 1,089)	135.3	(98.1 to 179.7)
		Female	1990	468	(344 to 639)	174	(126.5 to 230.8)

			2013	769	(595 to 1,038)	126.1	(97.9 to 166.4)
		Doth			· · · · · ·		
		Both	1990	819	(661 to 1,017)	161.3	(129.5 to 196.3)
			2013	1,571	(1,281 to 1,923)	131.5	(108.2 to 159.0)
	DALYs	Male	1990	10,975	(7,911 to 14,820)	2482.5	(1738.0 to 3372.8)
			2013	19,370	(13,674 to 26,151)	2193.7	(1553.1 to 2990.7)
		Female	1990	13,485	(9,461 to 18,662)	2838.7	(2027.9 to 3925.2)
			2013	15,339	(11,565 to 20,574)	1729	(1311.7 to 2356.8)
		Both	1990	24,461	(19,238 to 31,234)	2672.1	(2128.7 to 3330.2)
			2013	34,710	(27,644 to 42,826)	1959.8	(1580.2 to 2414.2)
	YLDs	Male	1990	51	(34 to 71)	13.4	(9.0 to 18.6)
			2013	133	(89 to 189)	13.7	(9.1 to 19.5)
		Female	1990	42	(29 to 60)	10.6	(7.2 to 14.9)
			2013	110	(72 to 154)	11.1	(7.5 to 15.5)
		Both	1990	94	(65 to 128)	11.9	(8.3 to 16.1)
			2013	244	(167 to 331)	12.3	(8.4 to 16.8)
Panama	Deaths	Male	1990	561	(477 to 676)	103.4	(87.2 to 124.8)
			2013	1,126	(903 to 1,367)	82.4	(66.4 to 99.9)
		Female	1990	563	(471 to 669)	92.7	(77.6 to 110.9)
			2013	1,040	(829 to 1,259)	66.6	(53.3 to 80.6)
		Both	1990	1,124	(1,007 to 1,274)	97.9	(87.7 to 111.0)
			2013	2,166	(1,876 to 2,487)	74.2	(64.3 to 85.2)
	DALYs	Male	1990	10,032	(8,605 to 11,968)	1539.5	(1320.6 to 1848.4)
			2013	17,823	(14,107 to 22,127)	1205.5	(955.2 to 1486.3)
		Female	1990	8,873	(7,334 to 10,193)	1283.1	(1068.6 to 1487.0)
			2013	13,028	(10,552 to 16,283)	815.9	(661.2 to 1018.9)
		Both	1990	18,906	(16,941 to 21,050)	1407.6	(1264.2 to 1576.4)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	30,852	(26,632 to 35,930)	1004.5	(869.4 to 1167.5)

	YLDs	Male	1990	179	(120 to 247)	25.3	(16.8 to 35.1)
			2013	420	(278 to 586)	27.5	(18.2 to 38.8)
		Female	1990	132	(89 to 181)	18.3	(12.4 to 25.6)
			2013	294	(197 to 406)	18.1	(12.2 to 25.0)
		Both	1990	311	(213 to 417)	21.8	(15.1 to 29.1)
			2013	714	(490 to 966)	22.7	(15.4 to 30.6)
Papua New							
Guinea	Deaths	Male	1990	522	(317 to 767)	82	(51.6 to 113.8)
			2013	1,096	(669 to 1,627)	86	(55.4 to 119.2)
		Female	1990	273	(170 to 442)	40.3	(26.0 to 64.5)
			2013	576	(367 to 900)	40.6	(26.7 to 62.6)
		Both	1990	796	(514 to 1,158)	58.2	(40.0 to 79.9)
			2013	1,672	(1,067 to 2,408)	60	(40.5 to 81.6)
	DALYs	Male	1990	17,945	(11,268 to 26,613)	1817.5	(1120.6 to 2611.2)
			2013	34,819	(21,458 to 52,977)	1844.8	(1160.6 to 2681.9)
		Female	1990	8,121	(5,193 to 13,217)	857.1	(545.0 to 1362.5)
			2013	15,496	(9,685 to 24,249)	823.3	(532.8 to 1269.6)
		Both	1990	26,067	(17,145 to 38,223)	1308.1	(872.6 to 1863.2)
			2013	50,316	(32,658 to 75,055)	1295.3	(848.2 to 1836.3)
	YLDs	Male	1990	396	(260 to 552)	57.8	(38.2 to 80.1)
			2013	827	(554 to 1,127)	58.1	(39.5 to 80.6)
		Female	1990	311	(207 to 427)	41.3	(27.6 to 57.3)
			2013	619	(421 to 847)	37.8	(25.5 to 50.9)
		Both	1990	707	(481 to 957)	48.5	(33.4 to 65.4)
			2013	1,446	(997 to 1,926)	46.7	(32.2 to 62.0)
Paraguay	Deaths	Male	1990	927	(796 to 1,104)	123.9	(105.5 to 147.3)
			2013	2,091	(1,667 to 2,578)	115.9	(92.8 to 142.5)
		Female	1990	1,088	(880 to 1,227)	121.7	(97.9 to 137.3)
			2013	2,197	(1,673 to 2,807)	109.1	(82.8 to 138.6)

Both 1990 2,016 (1,784 to 2,224) 122.8 (108.0 to 135.6)								
DALYS Male 1990 19,514 (16,852 to 22,925) 2010 (1737.2 to 2379.6)			Both	1990	2,016	(1,784 to 2,224)	122.8	(108.0 to 135.6)
2013 44,482 (32,899 to 51,677) 1962.2 (1568.9 to 2432.3)				2013	4,289	(3,608 to 5,119)	112.7	(94.7 to 134.5)
Female 1990 20,423 (16,587 to 23,205) 1925.8 (1562.4 to 2177.7)		DALYs	Male	1990	19,514	(16,852 to 22,925)	2010	(1737.2 to 2379.6)
Both 1990 39,937 (35,529 to 48,736) 1703.1 (1282.8 to 2213.8)				2013	41,482	(32,899 to 51,677)	1962.2	(1568.9 to 2432.3)
Both 1990 39,937 (35,529 to 43,965) 1964.3 (1739.7 to 2160.6)			Female	1990	20,423	(16,587 to 23,205)	1925.8	(1562.4 to 2177.7)
YLDS				2013	37,443	(28,272 to 48,736)	1703.1	(1282.8 to 2213.8)
YLDs Male 1990 439 (293 to 608) 43.4 (29.3 to 60.5) 2013 926 (615 to 1,288) 42.4 (28.1 to 58.2) Female 1990 307 (209 to 442) 28 (18.8 to 41.0) Both 1990 747 (515 to 1,028) 35.3 (24.5 to 48.5) Uncertainty Measure sex year number Uncertainty Interval Rate Interval Uncertainty Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,951 (2,228 to 3,381) 63.4 (46.3 to 71.3) Female 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) DALYS Male 1990			Both	1990	39,937	(35,529 to 43,965)	1964.3	(1739.7 to 2160.6)
Country Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4)				2013	78,926	(66,081 to 94,133)	1832.7	(1536.1 to 2188.4)
Female 1990 307 (209 to 442) 28 (18.8 to 41.0)		YLDs	Male	1990	439	(293 to 608)	43.4	(29.3 to 60.5)
Both 1990 747 (515 to 1,028) 35.3 (24.5 to 48.5) Uncertainty				2013	926	(615 to 1,288)	42.4	(28.1 to 58.2)
Country measure sex year number Uncertainty Interval Rate Interval Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Pemale 1990 2,551 (2,2131 to 2,981) 62.4 (50.7 to 73.4) Peru Pemale 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Pemale 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Pemale 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Pemale 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Pemale 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) Pemale 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) Pemale 1990 60,318 <td< td=""><td></td><td></td><td>Female</td><td>1990</td><td>307</td><td>(209 to 442)</td><td>28</td><td>(18.8 to 41.0)</td></td<>			Female	1990	307	(209 to 442)	28	(18.8 to 41.0)
Country measure sex year number Uncertainty Interval Rate Interval Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Pemale 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Pemale 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Pemale 1990 5,466 (4,596 to 6,007) 42.7 (33.6 to 61.7) Pemale 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Pemale 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) Pemale 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) Pemale 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) Pemale 1990 60,318 (49,195 to 74,9				2013	616	(415 to 854)	26.9	(17.8 to 37.3)
Country measure sex year number Uncertainty Interval Rate Interval Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Female 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Peru Both 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Peru Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Peru Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Peru Both 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) Peru Pemale 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) Peru Pemale 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.			Both	1990	747	(515 to 1,028)	35.3	(24.5 to 48.5)
Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) Peru Deaths 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Peru Both 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) Peru Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Peru Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) Peru Both 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) Peru 2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Peru 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) Peru								Uncertainty
Peru Deaths Male 1990 2,551 (2,131 to 2,981) 62.4 (50.7 to 73.4) 2013 4,368 (3,391 to 5,702) 48.8 (38.1 to 63.3) Female 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) 2013 4,615 (3,602 to 6,607) 42.7 (33.6 to 61.7) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) 2013 8,983 (7,502 to 11,373) 45.5 (38.1 to 58.5) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) 2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5	Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
2013 4,368 (3,391 to 5,702) 48.8 (38.1 to 63.3) Female 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) 2013 4,615 (3,602 to 6,607) 42.7 (33.6 to 61.7) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) 2013 8,983 (7,502 to 11,373) 45.5 (38.1 to 58.5) DALYs Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) 2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	1,543	(1,047 to 2,101)	34.5	(23.5 to 46.5)
Female 1990 2,915 (2,228 to 3,381) 63.4 (46.3 to 71.3) 2013 4,615 (3,602 to 6,607) 42.7 (33.6 to 61.7) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 60,318 (49,195 to 74,939) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)	Peru	Deaths	Male	1990	2,551	(2,131 to 2,981)	62.4	(50.7 to 73.4)
2013 4,615 (3,602 to 6,607) 42.7 (33.6 to 61.7) Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 60,318 (49,195 to 74,939) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	4,368	(3,391 to 5,702)	48.8	(38.1 to 63.3)
Both 1990 5,466 (4,596 to 6,056) 63.2 (51.5 to 70.1) 2013 8,983 (7,502 to 11,373) 45.5 (38.1 to 58.5) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) 2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)			Female	1990	2,915	(2,228 to 3,381)	63.4	(46.3 to 71.3)
DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) DALYS Male 1990 64,067 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) DALYS Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) DALYS Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) DALYS 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	4,615	(3,602 to 6,607)	42.7	(33.6 to 61.7)
DALYs Male 1990 64,067 (53,947 to 74,715) 1088.2 (913.9 to 1268.4) 2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)			Both	1990	5,466	(4,596 to 6,056)	63.2	(51.5 to 70.1)
2013 89,074 (66,887 to 117,581) 815.9 (622.0 to 1068.8) Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	8,983	(7,502 to 11,373)	45.5	(38.1 to 58.5)
Female 1990 60,318 (49,195 to 74,939) 994.3 (779.0 to 1181.6) 2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)		DALYs	Male	1990	64,067	(53,947 to 74,715)	1088.2	(913.9 to 1268.4)
2013 78,419 (58,779 to 108,392) 659.9 (504.7 to 924.3) Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	89,074	(66,887 to 117,581)	815.9	(622.0 to 1068.8)
Both 1990 124,386 (109,200 to 142,152) 1041.9 (891.1 to 1159.4) 2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)			Female	1990	60,318	(49,195 to 74,939)	994.3	(779.0 to 1181.6)
2013 167,494 (137,567 to 209,811) 734.5 (606.7 to 920.8)				2013	78,419	(58,779 to 108,392)	659.9	(504.7 to 924.3)
			Both	1990	124,386	(109,200 to 142,152)	1041.9	(891.1 to 1159.4)
YLDs Male 1990 1,080 (735 to 1,497) 18.7 (12.7 to 25.6)				2013	167,494	(137,567 to 209,811)	734.5	(606.7 to 920.8)
		YLDs	Male	1990	1,080	(735 to 1,497)	18.7	(12.7 to 25.6)

			2013	2,393	(1,573 to 3,279)	21.7	(14.3 to 29.6)
		Female	1990	706	(473 to 977)	11.4	(7.7 to 15.8)
			2013	1,361	(924 to 1,872)	11.2	(7.6 to 15.2)
		Both	1990	1,786	(1,259 to 2,382)	14.9	(10.6 to 20.0)
			2013	3,754	(2,608 to 4,985)	16.1	(11.2 to 21.5)
Philippines	Deaths	Male	1990	13,968	(12,057 to 16,246)	140	(120.5 to 163.8)
			2013	34,642	(26,741 to 43,959)	179.7	(142.4 to 224.6)
		Female	1990	10,740	(9,003 to 12,823)	97.4	(79.5 to 116.2)
			2013	27,407	(20,348 to 35,174)	122.5	(91.0 to 154.7)
		Both	1990	24,709	(22,490 to 27,611)	116.6	(105.3 to 130.5)
			2013	62,049	(51,832 to 74,052)	147.7	(125.0 to 173.0)
	DALYs	Male	1990	392,931	(340,870 to 460,986)	2777.9	(2404.3 to 3217.7)
			2013	913,931	(688,129 to 1,186,876)	3445.8	(2678.3 to 4359.0)
		Female	1990	243,820	(208,748 to 296,321)	1671.3	(1429.4 to 2014.8)
			2013	587,157	(437,286 to 772,439)	2045.7	(1523.9 to 2645.5)
		Both	1990	636,751	(578,081 to 725,732)	2189.4	(1997.8 to 2449.7)
					(1,234,567 to		
			2013	1,501,088	1,833,385)	2687	(2252.4 to 3215.2)
	YLDs	Male	1990	6,455	(4,323 to 8,845)	57.3	(39.1 to 79.0)
			2013	14,216	(9,698 to 19,291)	60.7	(42.1 to 82.3)
		Female	1990	5,805	(4,025 to 7,875)	43.7	(29.9 to 59.8)
			2013	12,156	(8,043 to 17,335)	44.2	(29.5 to 63.5)
		Both	1990	12,261	(8,419 to 16,148)	49.7	(34.8 to 65.7)
			2013	26,372	(18,249 to 35,492)	51.6	(36.0 to 69.4)
Poland	Deaths	Male	1990	31,416	(27,583 to 37,196)	230.1	(202.9 to 271.3)
			2013	30,556	(25,671 to 34,885)	138.7	(116.4 to 158.1)
		Female	1990	44,992	(36,651 to 48,247)	187.3	(152.8 to 200.8)
			2013	38,181	(33,045 to 48,685)	91.9	(80.3 to 116.6)
		Both	1990	76,409	(67,846 to 82,316)	205.3	(182.2 to 221.0)

							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	68,737	(62,031 to 79,522)	110.8	(100.5 to 127.7)
	DALYs	Male	1990	556,830	(487,133 to 677,579)	3595.1	(3147.3 to 4310.2)
			2013	474,039	(389,880 to 544,621)	2063.1	(1702.8 to 2360.9)
		Female	1990	582,467	(475,596 to 627,383)	2492.2	(2031.2 to 2692.4)
			2013	395,773	(349,967 to 506,360)	1097.6	(972.0 to 1413.3)
		Both	1990	1,139,297	(1,031,590 to 1,249,948)	2968.3	(2683.9 to 3251.2)
			2013	869,813	(786,957 to 991,229)	1510.2	(1364.9 to 1712.7)
	YLDs	Male	1990	3,924	(2,540 to 5,624)	22.6	(14.7 to 32.1)
			2013	6,437	(4,245 to 9,381)	27.2	(17.9 to 39.8)
		Female	1990	3,403	(2,209 to 4,869)	15.3	(10.0 to 21.8)
			2013	4,949	(3,060 to 7,154)	16.2	(10.1 to 23.1)
		Both	1990	7,327	(4,963 to 9,971)	18.6	(12.6 to 25.2)
			2013	11,386	(7,796 to 15,646)	21.1	(14.4 to 28.9)
Portugal	Deaths	Male	1990	10,421	(9,402 to 11,634)	206.9	(186.1 to 230.9)
			2013	8,857	(7,580 to 9,984)	104.9	(90.0 to 118.5)
		Female	1990	14,066	(11,483 to 15,155)	174	(142.6 to 187.8)
			2013	10,263	(8,396 to 14,588)	73.7	(60.7 to 104.2)
		Both	1990	24,487	(21,690 to 26,101)	188.8	(167.1 to 201.3)
			2013	19,121	(16,769 to 23,472)	87.1	(76.9 to 106.2)
	DALYs	Male	1990	165,948	(148,857 to 186,623)	3126.8	(2804.6 to 3521.6)
			2013	121,636	(105,003 to 136,333)	1513.6	(1299.7 to 1695.8)
		Female	1990	172,262	(142,273 to 184,235)	2269.4	(1864.3 to 2425.0)
			2013	106,157	(88,663 to 147,012)	896.3	(749.4 to 1251.3)
		Both	1990	338,211	(305,766 to 360,103)	2649.7	(2395.7 to 2824.4)
			2013	227,793	(203,520 to 269,945)	1170.8	(1046.2 to 1377.5)

	YLDs	Male	1990	6,195	(4,160 to 8,681)	114.6	(76.5 to 159.6)
			2013	8,289	(5,525 to 11,302)	105.4	(70.1 to 143.3)
		Female	1990	6,443	(4,311 to 9,134)	89.1	(59.7 to 125.4)
			2013	10,529	(6,965 to 14,220)	99.4	(65.6 to 132.5)
		Both	1990	12,639	(8,655 to 17,070)	100	(68.8 to 135.1)
			2013	18,819	(13,079 to 24,711)	102.2	(70.8 to 134.8)
Qatar	Deaths	Male	1990	44	(35 to 56)	64.1	(50.9 to 82.5)
			2013	82	(62 to 105)	34.6	(26.8 to 44.8)
		Female	1990	34	(23 to 43)	98.6	(66.5 to 123.0)
			2013	39	(29 to 54)	44.5	(33.8 to 60.2)
		Both	1990	79	(67 to 92)	77.7	(65.3 to 90.5)
			2013	122	(100 to 147)	39.6	(32.7 to 47.8)
	DALYs	Male	1990	1,523	(1,227 to 1,934)	983.3	(800.3 to 1245.3)
			2013	2,866	(2,176 to 3,701)	461.1	(362.3 to 578.1)
		Female	1990	1,054	(664 to 1,349)	1530.6	(989.1 to 1921.6)
			2013	977	(714 to 1,375)	519.7	(393.0 to 705.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	2,578	(2,161 to 3,040)	1173	(984.5 to 1363.1)
			2013	3,844	(3,115 to 4,707)	491.1	(410.0 to 582.0)
	YLDs	Male	1990	18	(11 to 27)	10.7	(6.9 to 15.7)
			2013	100	(64 to 147)	10.9	(7.2 to 15.7)
		Female	1990	5	(3 to 7)	8.3	(5.4 to 11.9)
			2013	21	(14 to 30)	8.7	(5.8 to 12.2)
		Both	1990	23	(15 to 33)	9.9	(6.6 to 14.1)
			2013	122	(80 to 173)	10.2	(7.0 to 14.3)
Romania	Deaths	Male	1990	25,677	(22,836 to 29,208)	262.8	(233.8 to 297.8)
			2013	25,036	(21,514 to 28,604)	190.4	(163.4 to 217.6)
		Female	1990	32,045	(28,326 to 36,004)	238.2	(210.1 to 269.1)

			2013	32,492	(27,781 to 36,541)	155.6	(133.3 to 174.7)
		Both	1990	57,723	(53,429 to 62,582)	249.4	(230.2 to 269.8)
			2013	57,528	(51,255 to 62,771)	170.6	(152.4 to 186.1)
	DALYs	Male	1990	445,959	(394,639 to 509,419)	4105.6	(3636.2 to 4675.6)
			2013	365,217	(312,418 to 414,788)	2728.8	(2338.6 to 3090.6)
		Female	1990	470,433	(415,531 to 529,713)	3379.9	(2987.0 to 3812.2)
			2013	368,938	(320,501 to 415,319)	1897.9	(1662.5 to 2159.0)
		Both	1990	916,392	(846,509 to 994,333)	3713.1	(3430.5 to 4026.0)
			2013	734,156	(662,476 to 801,789)	2266.3	(2050.4 to 2477.9)
	YLDs	Male	1990	3,287	(2,139 to 4,613)	28.2	(18.4 to 39.4)
			2013	4,164	(2,734 to 5,915)	31.3	(20.6 to 44.5)
		Female	1990	2,688	(1,754 to 3,737)	19.5	(12.7 to 27.0)
			2013	2,973	(1,913 to 4,332)	17.9	(11.7 to 26.0)
		Both	1990	5,975	(4,042 to 8,066)	23.5	(15.9 to 31.6)
			2013	7,137	(4,803 to 9,819)	24	(16.1 to 32.9)
Russia	Deaths	Male	1990	107,127	(91,270 to 121,713)	262.1	(225.6 to 296.0)
			2013	144,866	(125,706 to 168,925)	222.4	(193.4 to 257.6)
		Female	1990	239,766	(199,967 to 265,033)	222.6	(185.3 to 246.1)
			2013	215,421	(191,329 to 285,011)	149.2	(133.3 to 197.5)
		Both	1990	346,894	(300,196 to 372,526)	235.3	(203.6 to 253.1)
			2013	360,288	(327,838 to 438,804)	175.7	(160.1 to 213.0)
					(1,716,101 to		
	DALYs	Male	1990	2,046,862	2,353,786)	4026.1	(3412.5 to 4578.4)
					(2,291,804 to		/
			2013	2,649,544	3,095,248)	3689.6	(3206.1 to 4279.9)
		Female	1990	3,321,579	(2,756,493 to 3,654,762)	3123.1	(2588.8 to 3446.4)
		i ciliale	1330	3,341,373	(2,396,671 to	3123.1	(2300.0 to 3440.4)
			2013	2,680,162		2040.6	(1827.7 to 2722.2)
		-1		, , ,	. , ,	L	· , , , , , , , , , , , , , , , , , , ,

	_			(4,644,873 to		
	Both	1990	5,368,442	· · · · · · · · · · · · · · · · · · ·	3428.8	(2967.7 to 3695.1)
		2012	5 000 TOS	• • •	2675.4	(2425 6 + 2222 5)
		2013	5,329,706	6,415,056)	26/5.4	(2425.6 to 3202.5)
measure	sex	vear	number	Uncertainty Interval	Rate	Uncertainty Interval
YLDs	Male	1990	17,035	(10,601 to 25,494)	27	(17.1 to 39.5)
		2013	21,751	(13,250 to 33,393)	28.3	(17.4 to 43.2)
	Female	1990	20,848	(13,351 to 29,977)	20.8	(13.6 to 29.8)
		2013	21,780	(12,854 to 32,674)	18.8	(11.4 to 28.0)
	Both	1990	37,883	(25,376 to 51,570)	23.4	(15.7 to 32.0)
		2013	43,532	(28,729 to 62,132)	22.6	(15.1 to 32.0)
Deaths	Male	1990	2,492	(1,899 to 3,108)	295	(227.9 to 361.2)
		2013	2,072	(1,571 to 2,655)	147.4	(112.7 to 186.7)
	Female	1990	2,935	(2,000 to 3,942)	292.7	(201.3 to 385.7)
		2013	2,121	(1,428 to 3,130)	132.3	(89.1 to 190.3)
	Both	1990	5,428	(4,361 to 6,643)	294.8	(235.4 to 358.6)
		2013	4,193	(3,326 to 5,195)	139.2	(109.9 to 172.4)
DALYs	Male	1990	66,763	(49,555 to 85,219)	5672.4	(4320.8 to 7049.8)
		2013	45,062	(33,780 to 59,300)	2424.6	(1843.8 to 3117.6)
	Female	1990	69,783	(45,111 to 95,301)	5389.7	(3633.3 to 7255.2)
		2013	39,763	(26,905 to 60,409)	1981.9	(1347.6 to 2949.9)
	Both	1990	136,547	(107,638 to 170,144)	5537.5	(4434.0 to 6783.4)
		2013	84,825	(67,025 to 106,867)	2185.8	(1736.0 to 2703.7)
YLDs	Male	1990	181	(115 to 266)	12.6	(7.8 to 18.7)
		2013	337	(218 to 485)	13.5	(8.9 to 19.4)
	Female	1990	144	(93 to 203)	9.7	(6.4 to 13.9)
		2013	267	(163 to 389)	9.4	(5.5 to 13.9)
	Both	1990	326	(223 to 457)	11.1	(7.6 to 15.5)
	Deaths	YLDs Male Female Both Deaths Male Female Both DALYS Male Female Both YLDs Male Female Female	measure sex year YLDs Male 1990 2013 Female 1990 2013 Both 1990 2013 Both 1990 2013 Female 1990 2013 Female 1990 2013 Both 1990 2013 DALYs Male 1990 2013 Female 1990 2013 Both 1990 2013 Both 1990 2013 Pemale 1990 2013 Female 1990 2013 Female 1990 2013 Female 1990 2013 Pemale 1990 2013 2013 YLDs Male 1990 2013 2013	measure sex year number YLDs Male 1990 17,035 2013 21,751 2013 21,751 Female 1990 20,848 2013 21,780 2013 21,780 Both 1990 37,883 2013 43,532 Deaths Male 1990 2,492 2013 2,072 2013 2,072 Female 1990 2,935 2013 2,121 80th 1990 5,428 2013 4,193 45,062 2013 45,062 Female 1990 66,763 2013 39,763 Both 1990 69,783 2013 39,763 Both 1990 136,547 2013 84,825 YLDs Male 1990 181 2013 337 Female 1990 144 2013 2013 2013 2013	Both 1990 5,368,442 5,800,348) (4,841,366 to 6,415,056) 2013 5,329,706 6,415,056) measure sex year number Uncertainty Interval YLDS Male 1990 17,035 (10,601 to 25,494) 2013 21,751 (13,250 to 33,393) (12,854 to 32,674) Both 1990 20,848 (13,351 to 29,977) Both 1990 37,883 (25,376 to 51,570) Both 1990 37,883 (25,376 to 51,570) Deaths Male 1990 2,492 (1,899 to 3,108) Deaths Male 1990 2,492 (1,899 to 3,108) Female 1990 2,935 (2,000 to 3,942) Deaths Female 1990 2,935 (2,000 to 3,942) Both 1990 5,428 (4,361 to 6,643) Both 1990 5,428 (4,361 to 6,643) Both 1990 66,763 (49,555 to 85,219) Both 1990 69,7	Both 1990 5,368,442 5,800,348) 3428.8 measure sex year number Uncertainty Interval (10,601 to 25,494) Rate YLDs Male 1990 17,035 (10,601 to 25,494) 27 Female 1990 20,848 (13,351 to 29,977) 20.8 Female 1990 20,848 (13,351 to 29,977) 20.8 Both 1990 37,883 (25,376 to 51,570) 23.4 Both 1990 37,883 (25,376 to 51,570) 23.4 Deaths Male 1990 2,492 (1,899 to 3,108) 295 Deaths Male 1990 2,935 (2,000 to 3,942) 292.7 Female 1990 2,935 (2,000 to 3,942) 292.7 DALYS Male 1990 5,428 (4,361 to 6,643) 294.8 DALYS Male 1990 66,763 (49,555 to 85,219) 5672.4 Female 1990 69,783 (45,111 to 95,301) 5389.7

			2013	605	(396 to 837)	11.3	(7.6 to 15.5)
Saint Lucia	Deaths	Male	1990	65	(55 to 78)	155.4	(133.0 to 186.2)
			2013	81	(61 to 105)	108.5	(81.4 to 139.7)
		Female	1990	87	(72 to 100)	170.3	(142.2 to 196.9)
			2013	109	(81 to 138)	111.8	(82.8 to 141.6)
		Both	1990	153	(137 to 170)	164.7	(146.8 to 183.9)
			2013	191	(152 to 228)	111.2	(88.8 to 132.7)
	DALYs	Male	1990	1,306	(1,131 to 1,545)	2793.3	(2407.7 to 3310.1)
			2013	1,429	(1,071 to 1,877)	1819.9	(1366.7 to 2373.8)
		Female	1990	1,483	(1,230 to 1,708)	2795.6	(2325.1 to 3210.8)
			2013	1,489	(1,093 to 1,931)	1616.7	(1184.5 to 2103.8)
		Both	1990	2,790	(2,526 to 3,103)	2805.4	(2527.3 to 3112.6)
			2013	2,918	(2,359 to 3,485)	1720.2	(1387.3 to 2055.8)
	YLDs	Male	1990	27	(18 to 37)	61.8	(42.1 to 85.5)
			2013	47	(32 to 64)	60.9	(41.6 to 83.4)
		Female	1990	23	(14 to 32)	44.9	(29.2 to 62.5)
			2013	42	(28 to 57)	46.4	(30.4 to 62.7)
		Both	1990	50	(34 to 67)	52.7	(36.2 to 70.6)
			2013	90	(62 to 120)	53.1	(36.8 to 70.7)
							Uncertainty
Country Saint Vincent and the	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Grenadines	Deaths	Male	1990	39	(32 to 46)	143.8	(119.9 to 169.9)
			2013	40	(32 to 51)	110.8	(87.5 to 139.2)
		Female	1990	61	(49 to 70)	167.4	(134.2 to 190.6)
			2013	53	(40 to 64)	119.3	(91.6 to 144.5)
		Both	1990	101	(88 to 112)	158.2	(137.0 to 176.4)
			2013	93	(77 to 110)	116.5	(95.9 to 136.8)
					•		

930) 2549.1 1,028) 1903.7 1,178) 2773.5 1,022) 1806.3	(2149.6 to 2987.7) (1509.9 to 2400.5) (2267.6 to 3147.4)
1,178) 2773.5	
	(2267.6 to 3147.4)
1,022) 1806.3	
	(1377.9 to 2216.2)
to 2,010) 2680.7	(2330.9 to 2955.6)
to 1,932) 1864.5	(1557.3 to 2202.9)
23) 57.8	(37.5 to 82.6)
32) 55.5	(37.6 to 77.3)
20) 40.9	(28.1 to 56.7)
28) 44.1	(28.5 to 62.1)
48.4	(33.0 to 65.4)
58) 49.5	(33.6 to 66.6)
33) 210.6	(141.3 to 269.0)
37) 151.7	(108.8 to 191.5)
55) 102.5	(67.3 to 174.6)
36) 102.6	(72.4 to 145.9)
.28) 152	(116.2 to 202.5)
.62) 123.8	(95.9 to 156.5)
to 2,250) 4153	(2786.0 to 5478.8)
to 1,999) 2699.7	(1959.8 to 3481.0)
1,192) 1764.1	(1102.8 to 2978.5)
1,528) 1737.3	(1202.6 to 2469.7)
to 3,193) 2904.7	(2163.7 to 3887.9)
to 3,353) 2190.3	(1678.3 to 2812.6)
26) 54.9	(35.5 to 77.4)
10) 55.4	(37.0 to 76.9)
22) 43.2	(29.1 to 58.7)
34) 41.8	(26.9 to 57.9)
	to 2,010) 2680.7 to 1,932) 1864.5 23) 57.8 32) 55.5 20) 40.9 28) 44.1 42) 48.4 58) 49.5 33) 210.6 37) 151.7 55) 102.5 36) 102.6 128) 152 162) 123.8 to 2,250) 4153 to 1,999) 2699.7 1,192) 1764.1 1,528) 1737.3 to 3,193) 2904.7 to 3,353) 2190.3 26) 54.9 40) 55.4

		Both	1990	35	(23 to 47)	48.6	(33.1 to 64.8)
			2013	54	(37 to 74)	48	(32.8 to 65.5)
Sao Tome and							
Principe	Deaths	Male	1990	37	(30 to 44)	142.7	(114.9 to 172.6)
			2013	54	(37 to 71)	154.6	(111.7 to 200.3)
		Female	1990	48	(38 to 59)	161.9	(130.4 to 203.0)
			2013	65	(44 to 88)	168.1	(116.5 to 224.5)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	85	(72 to 100)	153.6	(132.0 to 183.2)
			2013	120	(89 to 150)	162.1	(126.0 to 200.5)
	DALYs	Male	1990	1,151	(916 to 1,421)	2943.3	(2385.2 to 3574.9)
			2013	1,502	(932 to 2,058)	2991.9	(1977.5 to 4070.4)
		Female	1990	1,222	(865 to 1,542)	3096.2	(2388.5 to 3868.8)
			2013	1,525	(964 to 2,074)	2960.7	(1948.0 to 4072.0)
		Both	1990	2,374	(1,902 to 2,805)	3027	(2569.4 to 3550.9)
			2013	3,027	(2,031 to 3,902)	2984.4	(2135.5 to 3821.9)
	YLDs	Male	1990	5	(3 to 7)	19.8	(13.5 to 27.6)
			2013	8	(5 to 11)	19.6	(12.8 to 27.9)
		Female	1990	4	(2 to 6)	14.3	(9.5 to 20.4)
			2013	7	(4 to 10)	14.7	(9.3 to 21.3)
		Both	1990	10	(7 to 13)	16.9	(11.8 to 22.6)
			2013	15	(10 to 21)	16.9	(11.5 to 23.0)
Saudi Arabia	Deaths	Male	1990	3,118	(2,295 to 4,178)	125.7	(94.4 to 165.6)
			2013	5,105	(3,963 to 6,736)	103.4	(81.2 to 134.8)
		Female	1990	2,729	(2,056 to 3,465)	119.8	(90.6 to 152.9)
			2013	4,392	(3,487 to 5,206)	86.8	(69.5 to 102.4)
		Both	1990	5,847	(4,846 to 7,038)	122.9	(102.9 to 147.4)
			2013	9,497	(8,118 to 11,188)	95.8	(82.5 to 112.5)

	DALYs	Male	1990	60,509	(43,104 to 84,824)	1837.7	(1317.8 to 2535.2)
			2013	102,564	(77,004 to 136,322)	1461	(1121.9 to 1950.6)
		Female	1990	48,305	(34,633 to 63,825)	1731.4	(1234.7 to 2250.4)
			2013	62,762	(49,376 to 76,388)	1093.2	(870.5 to 1318.7)
		Both	1990	108,814	(87,513 to 136,070)	1786.3	(1431.7 to 2216.9)
			2013	165,326	(136,894 to 202,063)	1292.1	(1080.7 to 1559.9)
	YLDs	Male	1990	222	(142 to 324)	5.6	(3.5 to 8.3)
			2013	673	(421 to 991)	6.7	(4.4 to 9.6)
		Female	1990	74	(47 to 103)	2.7	(1.7 to 3.8)
			2013	237	(150 to 341)	3.3	(2.1 to 4.9)
		Both	1990	296	(199 to 415)	4.3	(2.9 to 6.2)
			2013	910	(596 to 1,294)	5.2	(3.5 to 7.4)
Senegal	Deaths	Male	1990	1,311	(1,048 to 1,585)	123	(98.1 to 148.6)
			2013	2,606	(2,074 to 3,289)	142.3	(114.9 to 175.7)
		Female	1990	1,726	(1,257 to 2,150)	148.8	(111.1 to 188.7)
			2013	4,024	(2,760 to 5,363)	178.1	(125.4 to 237.0)
		Both	1990	3,037	(2,553 to 3,520)	137.9	(114.4 to 162.1)
			2013	6,631	(5,281 to 8,131)	163	(129.8 to 200.1)
	DALYs	Male	1990	34,198	(27,665 to 41,520)	2198.5	(1764.4 to 2644.3)
			2013	59,319	(45,825 to 75,786)	2424.6	(1919.0 to 3077.3)
		Female	1990	40,079	(29,979 to 50,097)	2456.1	(1791.8 to 3042.1)
			2013	80,409	(53,209 to 109,032)	2821.7	(1911.5 to 3767.4)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	74,278	(62,743 to 86,146)	2345.4	(1965.8 to 2709.5)
			2013	139,729	(108,244 to 171,029)	2654	(2096.4 to 3246.0)
	YLDs	Male	1990	321	(212 to 448)	19.4	(12.7 to 27.6)
			2013	548	(362 to 792)	19	(12.4 to 27.8)
		Female	1990	252	(167 to 360)	14.1	(9.2 to 20.3)

	DALYs	Male	1990	889	(753 to 1,051)	4063.9	(3435.9 to 4769.5)
			2013	92	(79 to 108)	117.4	(101.2 to 136.8)
		Both	1990	78	(69 to 93)	159.4	(141.3 to 190.5)
			2013	49	(38 to 61)	101	(77.4 to 125.6)
		Female	1990	37	(30 to 49)	123.6	(103.7 to 163.9)
			2013	43	(35 to 52)	136.3	(112.6 to 165.6)
Seychelles	Deaths	Male	1990	41	(34 to 48)	207.2	(174.7 to 244.1)
			2013	3,090	(2,127 to 4,161)	24	(16.4 to 32.3)
		Both	1990	2,304	(1,557 to 3,120)	22.6	(15.2 to 30.4)
			2013	1,243	(814 to 1,758)	17.8	(11.7 to 25.1)
		Female	1990	1,000	(649 to 1,394)	18.1	(11.9 to 25.5)
			2013	1,846	(1,224 to 2,557)	31.3	(20.8 to 43.4)
	YLDs	Male	1990	1,303	(876 to 1,804)	27.8	(18.8 to 38.3)
			2013	218,494	(196,209 to 239,416)	1610.7	(1446.5 to 1763.6)
		Both	1990	228,481	(186,736 to 268,472)	2393.8	(1964.9 to 2798.8)
			2013	110,505	(97,851 to 128,631)	1405.3	(1253.0 to 1640.0)
		Female	1990	115,115	(85,631 to 140,754)	2146.6	(1609.0 to 2617.8)
			2013	107,989	(90,185 to 123,809)	1859.4	(1549.5 to 2131.5)
	DALYs	Male	1990	113,366	(87,760 to 143,459)	2711.2	(2117.0 to 3417.0)
			2013	16,353	(14,569 to 17,960)	118	(105.0 to 129.7)
		Both	1990	14,019	(11,628 to 16,305)	159.6	(132.3 to 185.3)
			2013	9,295	(8,047 to 10,635)	109.7	(95.1 to 125.8)
		Female	1990	7,634	(5,812 to 9,312)	148.4	(113.7 to 181.4)
	2 3 3 4 1 1 2	1110110	2013	7,058	(5,864 to 8,172)	128	(106.0 to 149.2)
Serbia	Deaths	Male	1990	6,385	(5,025 to 7,934)	175	(137.9 to 216.5)
		200	2013	1,056	(711 to 1,458)	16.5	(11.1 to 22.7)
		Both	1990	573	(387 to 779)	16.6	(11.1 to 22.6)
			2013	508	(328 to 721)	14.5	(9.3 to 20.4)

			2013	910	(740 to 1,103)	2397.7	(1975.3 to 2878.6)
		Female	1990	590	(501 to 767)	2071.6	(1766.7 to 2690.7)
			2013	697	(536 to 870)	1487.8	(1144.3 to 1859.2)
		Both	1990	1,480	(1,315 to 1,731)	2975.3	(2645.0 to 3470.9)
			2013	1,608	(1,392 to 1,857)	1918.6	(1665.6 to 2222.5)
	YLDs	Male	1990	12	(8 to 17)	61.6	(41.4 to 83.6)
			2013	21	(14 to 30)	61.3	(42.0 to 84.2)
		Female	1990	12	(8 to 16)	43.7	(29.6 to 59.5)
			2013	20	(13 to 27)	45	(29.6 to 60.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	25	(17 to 33)	51.4	(35.4 to 69.6)
			2013	42	(29 to 55)	52.1	(36.6 to 69.0)
Sierra Leone	Deaths	Male	1990	1,058	(837 to 1,350)	151	(121.7 to 189.6)
			2013	1,590	(1,228 to 2,046)	187.5	(147.9 to 233.6)
		Female	1990	835	(638 to 1,056)	134.1	(104.4 to 172.3)
			2013	1,320	(965 to 1,790)	166	(124.2 to 220.9)
		Both	1990	1,893	(1,599 to 2,265)	142.4	(121.4 to 169.8)
			2013	2,910	(2,386 to 3,538)	176.2	(145.8 to 210.4)
	DALYs	Male	1990	32,561	(25,585 to 40,965)	2989.6	(2387.6 to 3801.9)
			2013	44,060	(33,332 to 57,307)	3473.1	(2693.4 to 4429.5)
		Female	1990	24,444	(17,441 to 31,183)	2374.7	(1840.4 to 3005.0)
			2013	34,078	(24,701 to 46,595)	2767.1	(2057.3 to 3732.3)
		Both	1990	57,006	(47,232 to 68,515)	2679.9	(2263.8 to 3199.8)
			2013	78,139	(62,886 to 95,432)	3107.2	(2554.2 to 3759.4)
	YLDs	Male	1990	191	(119 to 271)	19.6	(12.2 to 27.6)
			2013	265	(171 to 374)	19.5	(12.6 to 27.3)
		Female	1990	146	(97 to 212)	14.8	(9.7 to 21.3)
			2013	203	(129 to 291)	13.8	(8.7 to 19.9)

		Both	1990	338	(228 to 468)	17.2	(11.6 to 23.9)
			2013	469	(312 to 644)	16.6	(11.0 to 22.8)
Singapore	Deaths	Male	1990	725	(612 to 862)	98.5	(82.5 to 117.1)
			2013	1,044	(839 to 1,288)	52.9	(42.6 to 65.4)
		Female	1990	955	(736 to 1,073)	105.3	(80.5 to 118.6)
			2013	1,449	(1,040 to 2,021)	50.1	(36.2 to 70.0)
		Both	1990	1,680	(1,442 to 1,842)	103.3	(87.6 to 113.3)
			2013	2,493	(2,025 to 3,132)	51.9	(42.3 to 64.9)
	DALYs	Male	1990	15,742	(13,387 to 18,485)	1755.2	(1495.8 to 2061.6)
			2013	17,780	(14,771 to 21,385)	839.2	(697.1 to 1012.3)
		Female	1990	15,875	(12,432 to 17,528)	1610.1	(1262.3 to 1780.0)
			2013	16,132	(12,590 to 22,343)	618.2	(485.3 to 852.0)
		Both	1990	31,618	(27,549 to 34,676)	1687.7	(1463.2 to 1844.5)
			2013	33,913	(28,984 to 41,063)	724.9	(617.3 to 878.2)
	YLDs	Male	1990	1,130	(758 to 1,570)	130.1	(86.9 to 179.8)
			2013	2,603	(1,771 to 3,531)	122.7	(83.9 to 165.7)
		Female	1990	790	(501 to 1,136)	78.3	(49.8 to 113.2)
			2013	1,963	(1,253 to 2,805)	77.4	(49.1 to 110.7)
		Both	1990	1,921	(1,308 to 2,545)	101.7	(69.3 to 135.3)
			2013	4,566	(3,094 to 6,198)	98.2	(66.8 to 133.9)
Slovakia	Deaths	Male	1990	3,923	(3,383 to 4,553)	200.5	(172.1 to 232.7)
			2013	3,217	(2,718 to 3,881)	124.1	(104.5 to 149.6)
		Female	1990	4,528	(3,592 to 5,046)	146.5	(116.8 to 163.4)
			2013	3,939	(3,366 to 4,963)	82.2	(70.6 to 103.1)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
,		Both	1990	8,452	(7,399 to 9,274)	168.6	(147.5 to 185.3)
			2013	7,157	(6,368 to 8,344)	98.4	(87.6 to 114.6)
	DALYs	Male	1990	68,183	(58,510 to 80,176)	3155.9	(2715.4 to 3681.8)

		1				ı	1
			2013	52,554	(44,034 to 62,866)	1822	(1533.2 to 2183.4)
		Female	1990	62,038	(49,703 to 68,252)	2023.7	(1620.1 to 2228.1)
			2013	45,685	(39,523 to 55,939)	1024.7	(883.1 to 1250.7)
		Both	1990	130,222	(116,301 to 143,090)	2509	(2239.5 to 2749.2)
			2013	98,239	(87,289 to 112,156)	1357.6	(1204.8 to 1547.6)
	YLDs	Male	1990	670	(449 to 942)	28.8	(19.3 to 40.5)
			2013	961	(636 to 1,323)	30.4	(20.2 to 42.0)
		Female	1990	569	(381 to 803)	19.2	(12.9 to 26.7)
			2013	744	(479 to 1,053)	18.5	(12.1 to 26.2)
		Both	1990	1,240	(840 to 1,695)	23.4	(15.9 to 31.9)
			2013	1,705	(1,166 to 2,306)	23.8	(16.3 to 32.1)
Slovenia	Deaths	Male	1990	1,196	(1,017 to 1,393)	159.1	(135.3 to 186.4)
			2013	1,029	(811 to 1,233)	73.9	(58.2 to 88.7)
		Female	1990	1,678	(1,346 to 1,843)	119.5	(96.1 to 131.1)
			2013	1,512	(1,228 to 1,869)	57.1	(46.8 to 70.7)
		Both	1990	2,874	(2,580 to 3,141)	134.6	(120.5 to 146.7)
			2013	2,541	(2,164 to 2,906)	64.4	(54.7 to 73.4)
	DALYs	Male	1990	20,710	(17,841 to 24,114)	2450.7	(2106.6 to 2841.4)
			2013	14,191	(11,468 to 16,941)	985.7	(796.1 to 1178.9)
		Female	1990	21,686	(17,345 to 23,677)	1604.9	(1282.6 to 1753.8)
			2013	13,869	(11,465 to 17,018)	623.5	(516.1 to 765.5)
		Both	1990	42,396	(38,252 to 46,205)	1946.5	(1754.4 to 2119.5)
			2013	28,060	(24,048 to 31,900)	784.2	(669.9 to 891.4)
	YLDs	Male	1990	277	(177 to 385)	29.1	(18.7 to 41.3)
			2013	421	(265 to 628)	29.4	(18.6 to 43.6)
		Female	1990	257	(169 to 364)	20.1	(13.3 to 28.6)
			2013	316	(207 to 450)	18.4	(12.0 to 26.0)
1		Both	1990	534	(359 to 730)	24	(16.2 to 32.8)

			2013	738	(498 to 1,004)	23.6	(15.9 to 32.1)
Solomon							
Islands	Deaths	Male	1990	120	(76 to 169)	212	(138.1 to 283.8)
			2013	244	(163 to 354)	217.4	(153.6 to 298.5)
		Female	1990	66	(41 to 99)	142.1	(93.7 to 210.9)
			2013	175	(106 to 261)	171.4	(108.6 to 243.3)
		Both	1990	187	(125 to 260)	179.9	(127.1 to 239.2)
			2013	419	(275 to 606)	194.7	(136.4 to 265.7)
	DALYs	Male	1990	3,371	(2,072 to 4,958)	4584.8	(2869.2 to 6518.7)
			2013	6,702	(4,180 to 10,159)	4574.4	(3006.3 to 6695.3)
		Female	1990	1,798	(1,111 to 2,768)	2855.1	(1823.1 to 4291.6)
			2013	4,376	(2,520 to 6,913)	3282.3	(1987.1 to 4962.5)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	5,170	(3,352 to 7,580)	3766.1	(2527.7 to 5268.7)
			2013	11,079	(6,747 to 16,607)	3934.7	(2549.8 to 5721.2)
	YLDs	Male	1990	35	(23 to 49)	59.5	(39.1 to 82.7)
			2013	66	(45 to 89)	55.3	(37.6 to 74.6)
		Female	1990	22	(15 to 31)	44	(28.6 to 61.3)
			2013	50	(34 to 69)	43.6	(29.4 to 60.4)
		Both	1990	58	(40 to 78)	52.4	(35.8 to 71.0)
			2013	117	(81 to 152)	49.6	(34.6 to 65.2)
Somalia	Deaths	Male	1990	1,556	(956 to 2,513)	158	(101.3 to 247.8)
			2013	1,968	(1,242 to 2,999)	136.7	(90.3 to 199.5)
		Female	1990	2,213	(1,173 to 3,350)	200	(111.3 to 295.0)
			2013	2,461	(1,444 to 3,613)	152.8	(93.2 to 215.3)
		Both	1990	3,770	(2,260 to 5,669)	181.6	(114.2 to 265.2)
			2013	4,429	(2,713 to 6,569)	145.9	(94.0 to 208.3)
	DALYs	Male	1990	38,563	(23,404 to 62,194)	2874.3	(1754.7 to 4623.7)

			2013	46,076	(27,849 to 72,197)	2442.9	(1530.2 to 3747.2)
		Female	1990	48,700	(24,976 to 75,877)	3459.5	(1814.6 to 5286.6)
			2013	50,766	(28,454 to 76,263)	2548.2	(1475.5 to 3779.4)
		Both	1990	87,263	(50,461 to 132,314)	3197.7	(1901.8 to 4829.7)
			2013	96,843	(56,992 to 146,783)	2507.3	(1509.0 to 3766.9)
	YLDs	Male	1990	204	(129 to 291)	13.4	(8.6 to 18.8)
			2013	308	(196 to 452)	13.3	(8.6 to 19.6)
		Female	1990	141	(91 to 208)	8.6	(5.5 to 12.5)
			2013	242	(155 to 351)	9.8	(6.2 to 14.2)
		Both	1990	346	(235 to 472)	10.9	(7.3 to 14.9)
			2013	551	(366 to 785)	11.5	(7.6 to 16.0)
South Africa	Deaths	Male	1990	8,931	(7,177 to 11,099)	124.1	(100.5 to 154.4)
			2013	12,479	(9,909 to 15,246)	105.6	(83.6 to 128.5)
		Female	1990	12,428	(9,939 to 14,626)	128.5	(105.6 to 151.1)
			2013	22,316	(17,629 to 27,283)	108.5	(85.8 to 132.4)
		Both	1990	21,359	(18,395 to 24,277)	127.8	(112.0 to 145.3)
			2013	34,795	(29,650 to 40,164)	109.2	(93.1 to 125.7)
	DALYs	Male	1990	214,340	(170,665 to 266,137)	2288	(1840.6 to 2853.2)
			2013	249,691	(196,490 to 309,141)	1747.2	(1393.2 to 2130.9)
		Female	1990	237,531	(167,931 to 286,654)	2152	(1616.0 to 2553.2)
			2013	314,725	(249,185 to 397,534)	1525.8	(1204.6 to 1917.8)
		Both	1990	451,871	(367,441 to 525,154)	2220.4	(1886.2 to 2545.7)
			2013	564,416	(480,913 to 660,243)	1635.9	(1393.1 to 1898.9)
	YLDs	Male	1990	2,314	(1,530 to 3,205)	23.3	(15.2 to 32.2)
			2013	4,188	(2,763 to 5,872)	25.4	(16.8 to 35.7)
		Female	1990	2,609	(1,765 to 3,658)	22.6	(15.1 to 31.8)
			2013	4,848	(3,140 to 6,794)	21.9	(14.2 to 31.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval

		Female	2013 1990	3,347 1,318	(2,460 to 4,334) (814 to 1,983)	180.8 159.1	(136.0 to 230.5) (102.8 to 228.9)
South Suudh	Deatils	iviale			-		-
South Sudan	Deaths	Male	2013 1990	71,807 1,653	(50,035 to 97,245) (1,048 to 2,479)	116.2 194.5	(80.9 to 157.0) (128.3 to 282.3)
		Both	1990	25,256	(17,473 to 33,983)	96.7	(66.2 to 130.1)
		5.4	2013	32,474	(22,358 to 43,953)	93.1	(64.3 to 126.5)
		Female	1990	11,996	(8,115 to 16,206)	80	(53.5 to 108.5)
			2013	39,332	(26,924 to 53,591)	145.6	(99.2 to 195.5)
	YLDs	Male	1990	13,260	(8,983 to 18,221)	122.8	(81.1 to 169.5)
			2013	710,619	(634,794 to 832,550)	1166.5	(1040.9 to 1368.9)
		Both	1990	1,016,028	(904,856 to 1,089,070)	3874.9	(3448.4 to 4127.7)
			2013	328,467	(279,389 to 448,025)	919.1	(780.8 to 1257.5)
		Female	1990	474,863	(365,311 to 509,374)	3197.5	(2490.6 to 3425.7)
			2013	382,152	(334,242 to 430,533)	1469.4	(1295.2 to 1646.8)
	DALYs	Male	1990	541,164	(488,230 to 631,627)	4849.6	(4397.1 to 5616.6)
			2013	50,362	(44,666 to 60,650)	86.6	(76.8 to 104.6)
		Both	1990	49,045	(43,749 to 52,142)	234.5	(207.5 to 249.6)
			2013	28,810	(24,346 to 38,787)	77.6	(65.6 to 104.7)
		Female	1990	26,217	(20,484 to 28,110)	205.6	(161.5 to 221.2)
			2013	21,552	(18,827 to 24,329)	97.7	(85.2 to 111.2)
South Korea	Deaths	Male	1990	22,828	(20,721 to 26,464)	281.2	(254.5 to 326.9)
			2013	9,036	(6,138 to 12,115)	23.3	(15.8 to 31.4)
		Both	1990	4,924	(3,427 to 6,648)	23	(16.0 to 31.1)

		Female	1990	29,916	(17,513 to 46,082)	2553.1	(1576.2 to 3823.5)
			2013	52,271	(38,569 to 70,374)	2158.6	(1583.9 to 2879.8)
		Both	1990	73,688	(50,280 to 100,704)	3076	(2125.1 to 4156.8)
			2013	129,813	(103,363 to 159,077)	2695	(2162.0 to 3286.2)
	YLDs	Male	1990	171	(110 to 259)	12.2	(8.0 to 18.0)
			2013	366	(227 to 533)	12.9	(8.1 to 18.4)
		Female	1990	127	(80 to 184)	8.8	(5.5 to 12.7)
			2013	277	(182 to 396)	9.2	(6.1 to 13.1)
		Both	1990	298	(202 to 423)	10.4	(7.1 to 14.6)
			2013	643	(437 to 894)	11	(7.5 to 15.1)
Spain	Deaths	Male	1990	18,162	(15,516 to 21,244)	91.6	(78.1 to 107.3)
			2013	17,321	(13,802 to 20,699)	44.9	(36.0 to 53.4)
		Female	1990	27,640	(21,267 to 30,563)	84.4	(65.3 to 93.0)
			2013	25,537	(20,770 to 35,745)	40.2	(33.3 to 56.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	45,802	(39,238 to 50,406)	88.3	(75.7 to 97.2)
			2013	42,858	(36,881 to 53,318)	42.6	(36.9 to 52.2)
	DALYs	Male	1990	292,780	(253,246 to 338,878)	1419.4	(1226.5 to 1644.5)
			2013	225 672	/404045	670.4	(FC2 0 + 202 C)
			2013	235,673	(194,947 to 275,426)	678.4	(562.8 to 793.6)
		Female	1990	322,028	(194,947 to 275,426) (259,399 to 353,051)	1086.5	(877.4 to 1198.9)
		Female					
		Female Both	1990	322,028	(259,399 to 353,051)	1086.5	(877.4 to 1198.9)
			1990 2013	322,028 242,326	(259,399 to 353,051) (203,483 to 326,878)	1086.5 487.7	(877.4 to 1198.9) (414.2 to 655.8)
	YLDs		1990 2013 1990	322,028 242,326 614,809	(259,399 to 353,051) (203,483 to 326,878) (547,639 to 672,636)	1086.5 487.7 1241.3	(877.4 to 1198.9) (414.2 to 655.8) (1106.6 to 1356.8)
	YLDs	Both	1990 2013 1990 2013	322,028 242,326 614,809 478,000	(259,399 to 353,051) (203,483 to 326,878) (547,639 to 672,636) (421,838 to 564,425)	1086.5 487.7 1241.3 576.5	(877.4 to 1198.9) (414.2 to 655.8) (1106.6 to 1356.8) (510.2 to 674.3)
	YLDs	Both	1990 2013 1990 2013 1990	322,028 242,326 614,809 478,000 20,483	(259,399 to 353,051) (203,483 to 326,878) (547,639 to 672,636) (421,838 to 564,425) (14,096 to 27,678)	1086.5 487.7 1241.3 576.5 96.3	(877.4 to 1198.9) (414.2 to 655.8) (1106.6 to 1356.8) (510.2 to 674.3) (66.6 to 129.6)
	YLDs	Both Male	1990 2013 1990 2013 1990 2013	322,028 242,326 614,809 478,000 20,483 34,234	(259,399 to 353,051) (203,483 to 326,878) (547,639 to 672,636) (421,838 to 564,425) (14,096 to 27,678) (23,520 to 47,340)	1086.5 487.7 1241.3 576.5 96.3 102	(877.4 to 1198.9) (414.2 to 655.8) (1106.6 to 1356.8) (510.2 to 674.3) (66.6 to 129.6) (70.0 to 141.3)

		1	1			ı	
			2013	65,810	(46,510 to 88,269)	87.1	(60.9 to 117.6)
Sri Lanka	Deaths	Male	1990	3,873	(3,244 to 4,796)	78.1	(64.6 to 97.2)
			2013	6,251	(4,806 to 7,979)	79.7	(60.9 to 101.7)
		Female	1990	2,636	(2,157 to 3,312)	57.7	(46.5 to 73.0)
			2013	6,086	(2,986 to 8,082)	67.7	(32.0 to 90.4)
		Both	1990	6,510	(5,777 to 7,767)	68	(60.1 to 81.2)
			2013	12,338	(8,658 to 15,122)	73.4	(50.4 to 90.3)
	DALYs	Male	1990	97,673	(82,889 to 117,366)	1599.5	(1357.7 to 1930.1)
			2013	126,673	(97,633 to 161,250)	1393.8	(1084.2 to 1767.9)
		Female	1990	58,988	(48,784 to 71,736)	1001.5	(837.7 to 1235.7)
			2013	92,927	(53,402 to 121,384)	918.4	(514.3 to 1200.0)
		Both	1990	156,661	(140,791 to 183,293)	1301.9	(1168.5 to 1530.5)
			2013	219,600	(166,891 to 263,232)	1141.1	(851.8 to 1365.9)
	YLDs	Male	1990	3,236	(2,181 to 4,441)	60.8	(41.0 to 83.1)
			2013	5,437	(3,645 to 7,391)	60.9	(40.5 to 82.9)
		Female	1990	2,538	(1,720 to 3,525)	48.6	(32.6 to 67.8)
			2013	5,219	(3,434 to 7,104)	50.2	(32.8 to 68.5)
		Both	1990	5,774	(4,017 to 7,660)	54.8	(38.3 to 72.5)
			2013	10,657	(7,237 to 14,238)	55.1	(37.7 to 73.5)
Sudan	Deaths	Male	1990	5,511	(4,159 to 6,875)	176.3	(135.2 to 215.9)
			2013	9,270	(6,789 to 11,884)	154.1	(114.3 to 191.9)
		Female	1990	6,707	(4,718 to 8,834)	191.7	(142.8 to 250.4)
			2013	10,170	(7,155 to 13,767)	155	(110.1 to 205.8)
		Both	1990	12,218	(9,783 to 14,545)	184.6	(152.8 to 219.1)
			2013	19,440	(15,416 to 23,965)	154.7	(124.5 to 186.5)
	DALYs	Male	1990	166,318	(119,494 to 216,939)	3266.7	(2456.6 to 4111.6)
			2013	232,696	(166,829 to 308,945)	2613.2	(1907.0 to 3372.8)
		Female	1990	194,017	(107,678 to 261,198)	3567.8	(2477.5 to 4725.1)

			2013	229,045	(154,363 to 320,673)	2500.3	(1730.7 to 3440.0)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
		Both	1990	360,336	(260,778 to 440,493)	3422.3	(2745.8 to 4095.1)
			2013	461,741	(362,590 to 578,955)	2556.5	(2015.4 to 3173.0)
	YLDs	Male	1990	468	(313 to 671)	10.8	(7.0 to 15.5)
			2013	1,001	(639 to 1,399)	11.1	(7.0 to 15.6)
		Female	1990	410	(265 to 585)	9.1	(5.9 to 13.0)
			2013	808	(516 to 1,172)	8.5	(5.5 to 12.3)
		Both	1990	879	(591 to 1,204)	9.9	(6.7 to 13.8)
			2013	1,810	(1,187 to 2,489)	9.7	(6.4 to 13.3)
Suriname	Deaths	Male	1990	138	(118 to 167)	144.4	(123.9 to 175.4)
			2013	254	(184 to 330)	155.2	(114.2 to 200.4)
		Female	1990	151	(133 to 192)	138.1	(121.4 to 176.1)
			2013	302	(217 to 387)	145.3	(104.3 to 185.3)
		Both	1990	290	(262 to 346)	141.7	(128.0 to 169.3)
			2013	556	(443 to 674)	151.2	(121.6 to 181.6)
	DALYs	Male	1990	3,282	(2,822 to 3,916)	2762.1	(2375.4 to 3337.2)
			2013	5,576	(3,973 to 7,367)	2847.8	(2059.6 to 3713.6)
		Female	1990	3,069	(2,713 to 3,747)	2413.2	(2132.9 to 2981.7)
			2013	5,116	(3,693 to 6,717)	2289	(1648.2 to 2986.9)
		Both	1990	6,352	(5,771 to 7,373)	2584.9	(2344.8 to 3047.1)
			2013	10,693	(8,457 to 13,182)	2562	(2037.7 to 3143.5)
	YLDs	Male	1990	64	(43 to 87)	61.2	(40.8 to 83.9)
			2013	112	(76 to 156)	61.2	(41.4 to 86.1)
		Female	1990	50	(33 to 69)	42.7	(28.6 to 58.5)
			2013	97	(63 to 137)	43.6	(28.2 to 62.4)
		Both	1990	114	(79 to 153)	51.3	(35.4 to 69.0)
			2013	209	(144 to 287)	51.4	(35.3 to 70.6)

Swaziland	Deaths	Male	1990	137	(100 to 174)	131.3	(98.3 to 165.0)
			2013	317	(250 to 397)	159.4	(128.2 to 198.7)
		Female	1990	196	(145 to 238)	149.6	(112.1 to 179.5)
			2013	373	(276 to 507)	150.7	(112.3 to 201.9)
		Both	1990	333	(268 to 386)	143	(115.0 to 165.4)
			2013	691	(568 to 845)	155.5	(128.4 to 190.2)
	DALYs	Male	1990	3,401	(2,522 to 4,374)	2369.7	(1746.6 to 3003.6)
			2013	8,282	(6,410 to 10,455)	3131.1	(2454.2 to 3948.3)
		Female	1990	4,217	(3,066 to 5,158)	2461.4	(1819.1 to 2980.1)
			2013	8,307	(6,173 to 11,542)	2564.9	(1911.1 to 3510.7)
		Both	1990	7,619	(6,002 to 8,917)	2432.8	(1957.2 to 2817.8)
			2013	16,589	(13,608 to 20,582)	2819.6	(2304.2 to 3447.6)
	YLDs	Male	1990	38	(25 to 53)	24.3	(16.2 to 33.5)
			2013	61	(39 to 89)	22.4	(14.1 to 32.9)
		Female	1990	44	(28 to 61)	22.2	(14.8 to 30.7)
			2013	74	(48 to 103)	22.3	(14.4 to 31.3)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	83	(57 to 114)	23.1	(15.9 to 31.5)
			2013	135	(91 to 188)	22.4	(15.0 to 31.0)
Sweden	Deaths	Male	1990	4,635	(3,828 to 5,742)	76.8	(63.4 to 95.4)
			2013	3,996	(2,981 to 4,946)	48.9	(36.5 to 60.3)
		Female	1990	6,264	(5,094 to 7,520)	65.3	(53.1 to 77.8)
			2013	5,432	(4,224 to 6,999)	43.5	(34.2 to 56.1)
		Both	1990	10,899	(9,701 to 12,501)	70.6	(63.4 to 81.1)
			2013	9,429	(7,773 to 11,079)	46.3	(38.5 to 54.1)
	DALYs	Male	1990	69,923	(59,842 to 84,853)	1186.3	(1015.1 to 1440.0)
			2013	52,794	(40,984 to 62,880)	686.6	(531.8 to 817.9)
		Female	1990	70,701	(59,601 to 83,177)	865.8	(731.4 to 1014.5)

			2013	50,822	(41,103 to 64,087)	502.2	(408.3 to 627.5)
		Both	1990	140,624	(127,691 to 157,530)	1011	(917.5 to 1135.5)
			2013	103,616	(88,353 to 118,810)	589.7	(497.8 to 671.6)
	YLDs	Male	1990	6,948	(4,477 to 9,792)	117.9	(76.3 to 165.3)
			2013	7,942	(5,144 to 11,053)	107	(69.9 to 148.8)
		Female	1990	5,954	(3,905 to 8,378)	79.4	(53.0 to 109.5)
			2013	6,407	(4,335 to 8,692)	74.5	(51.1 to 100.9)
		Both	1990	12,903	(8,717 to 17,649)	96.6	(65.0 to 130.9)
			2013	14,350	(9,951 to 19,070)	90.1	(62.5 to 120.5)
Switzerland	Deaths	Male	1990	2,708	(2,245 to 3,290)	72.3	(59.8 to 87.7)
			2013	2,654	(1,949 to 3,239)	42.6	(31.3 to 51.9)
		Female	1990	4,269	(3,406 to 4,849)	62.2	(49.9 to 70.3)
			2013	4,076	(3,267 to 5,525)	38.9	(31.3 to 52.6)
		Both	1990	6,977	(6,072 to 7,817)	66.7	(58.3 to 74.8)
			2013	6,730	(5,639 to 8,163)	40.8	(34.3 to 49.2)
	DALYs	Male	1990	39,644	(33,588 to 48,124)	1040.3	(880.5 to 1260.3)
			2013	33,868	(25,786 to 40,115)	565.4	(431.4 to 670.3)
		Female	1990	45,057	(37,138 to 50,472)	760.4	(625.5 to 853.7)
			2013	36,132	(29,820 to 46,846)	420.1	(349.7 to 540.6)
		Both	1990	84,702	(75,755 to 94,383)	880.3	(786.2 to 981.5)
			2013	70,000	(60,098 to 81,753)	487	(416.8 to 562.9)
	YLDs	Male	1990	3,946	(2,609 to 5,529)	103	(68.0 to 144.4)
			2013	6,124	(4,188 to 8,549)	105.4	(72.1 to 146.5)
		Female	1990	4,040	(2,702 to 5,595)	76	(51.7 to 104.0)
			2013	5,484	(3,661 to 7,621)	75.8	(51.1 to 104.5)
		Both	1990	7,986	(5,423 to 10,587)	87.5	(59.7 to 115.3)
			2013	11,608	(8,060 to 15,487)	89.7	(62.8 to 118.9)
Syria	Deaths	Male	1990	5,314	(4,122 to 6,735)	229.4	(177.8 to 293.9)
34114	Deaths	IVIAIC	1330	3,311	(1,122 to 0,733)	223.1	(177.0 to 233.3)

			2013	7,345	(5,393 to 9,518)	159.7	(116.7 to 204.1)
		Female	1990	3,647	(2,689 to 5,502)	169.3	(123.8 to 258.8)
			2013	5,837	(4,238 to 7,518)	130.9	(95.7 to 167.6)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	8,962	(7,305 to 11,444)	199	(162.2 to 258.8)
			2013	13,182	(10,692 to 15,776)	145.2	(117.0 to 172.7)
	DALYs	Male	1990	186,675	(144,005 to 239,811)	5025.3	(3842.7 to 6431.8)
			2013	173,258	(128,176 to 231,326)	2795.2	(2053.3 to 3652.3)
		Female	1990	112,305	(81,382 to 161,738)	3256.1	(2364.6 to 4961.8)
			2013	112,903	(82,367 to 146,597)	1937.5	(1413.4 to 2525.3)
		Both	1990	298,980	(238,581 to 376,190)	4144.4	(3346.9 to 5362.7)
			2013	286,162	(231,833 to 348,119)	2364.6	(1899.7 to 2843.9)
	YLDs	Male	1990	276	(186 to 392)	10.4	(7.0 to 14.9)
			2013	695	(459 to 969)	10.9	(7.3 to 15.1)
		Female	1990	229	(155 to 329)	8.7	(5.9 to 12.6)
			2013	552	(359 to 780)	8.7	(5.7 to 12.1)
		Both	1990	505	(348 to 687)	9.6	(6.7 to 13.0)
			2013	1,248	(848 to 1,664)	9.8	(6.7 to 13.0)
Taiwan	Deaths	Male	1990	6,983	(5,811 to 8,097)	115.8	(96.8 to 134.1)
			2013	8,157	(6,949 to 9,764)	56.6	(48.4 to 67.5)
		Female	1990	5,035	(3,624 to 6,033)	94.8	(67.6 to 114.7)
			2013	6,043	(4,992 to 7,679)	38.9	(32.2 to 49.3)
		Both	1990	12,018	(10,284 to 13,542)	106.4	(89.7 to 120.8)
			2013	14,201	(12,674 to 16,471)	47.3	(42.3 to 54.8)
	DALYs	Male	1990	161,675	(134,538 to 187,188)	2142.9	(1800.5 to 2467.2)
			2013	151,424	(132,200 to 176,352)	1065.9	(932.5 to 1235.3)
		Female	1990	94,822	(69,419 to 112,816)	1483.1	(1085.1 to 1773.5)
			2013	85,127	(73,331 to 105,378)	564.4	(489.2 to 702.1)

		Both	1990	256,498	(221,380 to 288,860)	1834.9	(1582.4 to 2054.7)
			2013	236,552	(214,892 to 267,771)	806.1	(732.0 to 912.2)
	YLDs	Male	1990	7,672	(5,295 to 10,244)	107.9	(74.5 to 143.4)
			2013	13,608	(9,097 to 18,704)	97	(65.2 to 133.7)
		Female	1990	4,600	(3,218 to 6,293)	71.4	(49.8 to 97.4)
			2013	9,576	(6,652 to 12,963)	62.8	(43.5 to 84.9)
		Both	1990	12,273	(8,631 to 16,187)	90	(63.7 to 118.1)
			2013	23,185	(16,099 to 31,100)	79.1	(55.0 to 106.0)
Tajikistan	Deaths	Male	1990	1,900	(1,628 to 2,177)	210.3	(179.3 to 240.5)
			2013	2,413	(1,925 to 2,940)	179.5	(144.2 to 217.2)
		Female	1990	2,301	(2,060 to 2,685)	180.6	(162.0 to 211.1)
			2013	2,759	(2,144 to 3,357)	169.1	(131.0 to 204.9)
		Both	1990	4,201	(3,841 to 4,655)	192.4	(175.5 to 213.2)
			2013	5,173	(4,344 to 5,924)	174.2	(145.9 to 198.3)
	DALYs	Male	1990	37,202	(31,808 to 42,802)	3467.3	(2974.9 to 3983.6)
			2013	47,358	(36,758 to 58,576)	2842.9	(2227.1 to 3475.8)
		Female	1990	40,277	(35,917 to 47,395)	2912.8	(2615.1 to 3408.1)
			2013	48,850	(37,630 to 60,198)	2568.1	(1982.6 to 3172.0)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
		Both	1990	77,479	(70,684 to 86,311)	3136.9	(2860.3 to 3484.1)
			2013	96,208	(80,643 to 111,930)	2697.9	(2270.7 to 3118.8)
	YLDs	Male	1990	221	(140 to 315)	17	(11.0 to 24.2)
			2013	357	(233 to 502)	16.3	(10.6 to 23.0)
		Female	1990	186	(122 to 260)	12.6	(8.3 to 17.6)
			2013	288	(184 to 405)	12.8	(8.3 to 17.9)
		Both	1990	408	(271 to 558)	14.7	(9.9 to 19.9)
			2013	645	(452 to 872)	14.5	(10.2 to 19.4)
Tanzania	Deaths	Male	1990	2,434	(1,722 to 3,079)	78.5	(54.1 to 99.4)

			2013	5,481	(4,023 to 7,009)	83.2	(60.6 to 105.8)
		Female	1990	2,670	(1,855 to 3,351)	71.2	(48.1 to 90.6)
		Terriale	2013	5,684	(3,896 to 7,599)	69.5	(46.8 to 93.5)
		Both	1990	5,105	(4,003 to 6,073)	74.4	(55.8 to 90.1)
		Dotti	2013	11,165	(8,575 to 13,535)	75.5	(57.7 to 92.1)
	DALYs	Male	1990	58,491	(42,834 to 73,711)	1291.4	(916.6 to 1631.3)
	DALIS	iviale	2013	112,910	(82,682 to 144,610)	1273.5	(937.3 to 1626.0)
		Female	1990	58,301	(42,854 to 73,443)	1147.1	(809.3 to 1433.1)
		Telliale	2013	111,327	(78,291 to 147,209)	1074	(748.5 to 1442.5)
		Both	1990	111,327	(93,769 to 138,114)	1213.9	(954.5 to 1437.6)
		BULII	2013		, ,		, ,
	VIDa	Mala		224,237 929	(175,698 to 271,617)	1163.8	(899.6 to 1405.4)
	YLDs	Male	1990		(582 to 1,363)	18	(11.4 to 26.2)
		F	2013	1,758	(1,114 to 2,608)	15.5	(9.4 to 23.2)
		Female	1990	754	(467 to 1,127)	13.2	(8.1 to 19.1)
			2013	1,376	(868 to 2,016)	11.4	(7.2 to 16.9)
		Both	1990	1,683	(1,122 to 2,353)	15.4	(10.3 to 21.8)
			2013	3,135	(2,097 to 4,438)	13.3	(8.7 to 19.0)
Thailand	Deaths	Male	1990	15,215	(12,881 to 18,422)	125.9	(107.9 to 147.4)
			2013	28,581	(22,746 to 36,721)	97.4	(78.2 to 124.9)
		Female	1990	17,907	(13,498 to 19,780)	130.1	(98.7 to 143.7)
			2013	33,118	(25,308 to 48,781)	91.7	(69.8 to 135.3)
		Both	1990	33,123	(28,971 to 36,552)	130.1	(112.0 to 141.7)
			2013	61,699	(50,902 to 79,803)	95	(78.1 to 124.2)
	DALYs	Male	1990	383,905	(317,175 to 493,591)	2219.4	(1874.6 to 2736.7)
			2013	550,386	(430,064 to 698,724)	1648.8	(1296.4 to 2092.9)
		Female	1990	296,003	(210,531 to 330,177)	1744.3	(1292.6 to 1926.9)
			2013	442,947	(342,219 to 621,909)	1166.4	(900.9 to 1656.0)
		Both	1990	679,908	(598,114 to 782,042)	1985.6	(1758.7 to 2215.6)

			2013	993,334	(827,295 to 1,222,054)	1400.3	(1167.6 to 1740.5)
	YLDs	Male	1990	8,827	(6,110 to 11,953)	59.6	(41.5 to 79.9)
	1.255	TVIGIC	2013	20,354	(14,227 to 27,539)	62	(43.3 to 83.8)
		Female	1990	7,570	(5,203 to 10,365)	44.5	(30.5 to 61.2)
			2013	17,432	(11,648 to 24,368)	44.8	(29.9 to 62.5)
				, -	()		Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	16,398	(11,515 to 21,456)	51.4	(36.0 to 67.0)
			2013	37,786	(26,595 to 50,001)	52.6	(36.8 to 69.9)
The Bahamas	Deaths	Male	1990	75	(62 to 90)	152.2	(124.7 to 183.7)
			2013	141	(101 to 185)	116.2	(84.9 to 148.9)
		Female	1990	94	(78 to 108)	143.9	(119.5 to 165.8)
			2013	180	(127 to 233)	104.2	(73.9 to 134.5)
		Both	1990	169	(150 to 190)	148.4	(130.5 to 166.4)
			2013	322	(256 to 384)	110.2	(88.7 to 130.4)
	DALYs	Male	1990	1,804	(1,503 to 2,138)	2774.6	(2324.0 to 3309.6)
			2013	3,013	(2,102 to 4,041)	2019	(1439.0 to 2657.1)
		Female	1990	1,802	(1,506 to 2,067)	2337.1	(1964.1 to 2678.0)
			2013	2,930	(2,033 to 3,932)	1585.7	(1109.0 to 2103.1)
		Both	1990	3,607	(3,238 to 4,029)	2541.5	(2262.8 to 2844.4)
			2013	5,943	(4,699 to 7,229)	1787.2	(1427.8 to 2157.4)
	YLDs	Male	1990	36	(24 to 50)	61.7	(41.2 to 86.6)
			2013	89	(59 to 125)	63	(41.4 to 88.5)
		Female	1990	31	(21 to 42)	42.6	(28.9 to 58.9)
			2013	79	(52 to 117)	43.1	(28.4 to 63.5)
		Both	1990	67	(46 to 91)	50.8	(34.7 to 68.9)
			2013	169	(119 to 227)	51.6	(36.3 to 69.6)
The Gambia	Deaths	Male	1990	161	(105 to 257)	120.7	(81.9 to 185.6)
			2013	351	(227 to 535)	151.2	(105.2 to 216.7)

		Female	1990	168	(108 to 261)	145.4	(99.6 to 214.9)
			2013	358	(228 to 551)	164.4	(108.3 to 242.3)
		Both	1990	330	(231 to 504)	132.6	(96.0 to 195.9)
			2013	709	(464 to 1,048)	157.9	(111.1 to 222.9)
	DALYs	Male	1990	4,469	(2,910 to 6,905)	2179.9	(1422.2 to 3484.0)
			2013	8,310	(5,150 to 13,087)	2565.9	(1661.6 to 3930.5)
		Female	1990	4,278	(2,646 to 6,545)	2480.8	(1585.3 to 3876.2)
			2013	7,970	(4,737 to 12,570)	2636.9	(1650.7 to 4061.9)
		Both	1990	8,747	(5,996 to 13,015)	2322	(1598.0 to 3578.0)
			2013	16,281	(9,995 to 25,144)	2601.5	(1682.9 to 3863.1)
	YLDs	Male	1990	39	(26 to 54)	19.9	(13.2 to 27.7)
			2013	73	(48 to 100)	19.6	(13.0 to 27.2)
		Female	1990	25	(17 to 37)	14.8	(9.9 to 21.6)
			2013	57	(38 to 81)	15.1	(10.1 to 21.6)
		Both	1990	65	(44 to 89)	17.5	(11.8 to 24.3)
			2013	130	(90 to 176)	17.4	(12.1 to 23.4)
Timor-Leste	Deaths	Male	1990	140	(102 to 182)	135.3	(100.4 to 172.6)
			2013	235	(172 to 306)	143.3	(104.7 to 183.4)
		Female	1990	130	(92 to 185)	143.5	(102.3 to 200.9)
			2013	247	(167 to 355)	141.2	(95.5 to 198.9)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	270	(212 to 346)	139.2	(110.2 to 176.2)
			2013	483	(383 to 617)	142.7	(111.8 to 179.7)
	DALYs	Male	1990	4,890	(3,547 to 6,523)	2608.3	(1927.4 to 3367.6)
			2013	6,034	(4,458 to 7,889)	2473.9	(1813.3 to 3200.3)
		Female	1990	4,263	(2,641 to 6,152)	2639.8	(1913.9 to 3734.1)
					1	i e	
			2013	5,264	(3,608 to 7,766)	2209.7	(1530.9 to 3178.0)

		Female	1990	25	(19 to 34)	119.2	(91.6 to 155.5)
			2013	28	(20 to 36)	105.9	(77.6 to 135.3)
Tonga	Deaths	Male	1990	20	(14 to 26)	103.7	(77.0 to 132.4)
			2013	522	(357 to 715)	16.5	(11.2 to 22.4)
		Both	1990	286	(192 to 393)	16.6	(11.0 to 22.9)
			2013	238	(155 to 328)	14.2	(9.3 to 19.5)
		Female	1990	134	(87 to 191)	14.9	(9.4 to 21.6)
			2013	283	(186 to 410)	19.3	(12.7 to 27.5)
	YLDs	Male	1990	152	(94 to 218)	18.4	(11.6 to 26.8)
			2013	61,476	(50,900 to 75,424)	2271.1	(1924.0 to 2747.2)
		Both	1990	42,590	(36,081 to 49,870)	2482.1	(2116.3 to 2914.6)
			2013	27,807	(21,234 to 37,694)	1988.4	(1577.0 to 2611.8)
		Female	1990	22,213	(17,158 to 28,299)	2507.1	(1987.4 to 3228.1)
			2013	33,669	(25,304 to 43,381)	2602.7	(2020.4 to 3255.9)
	DALYs	Male	1990	20,376	(16,326 to 24,869)	2428.5	(1964.3 to 2964.1)
			2013	2,536	(2,146 to 3,078)	139.5	(118.9 to 167.3)
		Both	1990	1,626	(1,392 to 1,908)	141.2	(121.2 to 167.9)
			2013	1,223	(968 to 1,610)	128.8	(102.2 to 168.4)
		Female	1990	911	(724 to 1,172)	147.5	(117.6 to 193.3)
			2013	1,312	(1,017 to 1,650)	152.2	(121.3 to 187.2)
Togo	Deaths	Male	1990	715	(580 to 872)	131.5	(107.0 to 160.2)
			2013	238	(161 to 322)	54	(36.9 to 73.3)
		Both	1990	124	(84 to 164)	52.3	(36.6 to 69.6)
			2013	110	(73 to 153)	48.7	(32.7 to 67.6)
		Female	1990	52	(33 to 72)	45.1	(30.0 to 62.3)
			2013	127	(84 to 176)	59.9	(40.8 to 82.4)
	YLDs	Male	1990	72	(48 to 97)	59.1	(40.4 to 79.5)
			2013	11,299	(8,902 to 14,565)	2344.6	(1856.8 to 2989.4)

			2013	38	(28 to 49)	97.2	(71.4 to 125.7)
		Both	1990	46	(37 to 56)	112.8	(92.1 to 136.2)
			2013	66	(54 to 80)	102.5	(83.2 to 123.2)
	DALYs	Male	1990	465	(332 to 608)	1880.3	(1370.7 to 2458.3)
			2013	607	(437 to 805)	1945.3	(1398.8 to 2548.2)
		Female	1990	486	(352 to 666)	1896.4	(1407.3 to 2577.7)
			2013	574	(415 to 765)	1485.1	(1053.0 to 1971.0)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	952	(757 to 1,190)	1897.1	(1525.2 to 2367.1)
			2013	1,181	(958 to 1,447)	1709.3	(1383.0 to 2091.7)
	YLDs	Male	1990	12	(8 to 17)	54	(35.6 to 75.2)
			2013	16	(11 to 23)	58.3	(38.1 to 81.5)
		Female	1990	10	(6 to 14)	39.7	(25.1 to 57.9)
			2013	15	(10 to 21)	42.6	(27.3 to 59.1)
		Both	1990	22	(15 to 29)	46.3	(31.8 to 62.0)
			2013	32	(22 to 44)	49.4	(34.3 to 66.8)
Trinidad and							
Tobago	Deaths	Male	1990	478	(408 to 556)	159.6	(136.4 to 186.9)
			2013	715	(567 to 874)	140	(111.8 to 169.9)
		Female	1990	514	(424 to 596)	132.5	(109.5 to 153.5)
			2013	623	(499 to 806)	92.5	(74.2 to 119.6)
		Both	1990	992	(891 to 1,096)	144.5	(129.6 to 160.1)
			2013	1,338	(1,154 to 1,565)	113.2	(98.1 to 132.8)
	DALYs	Male	1990	9,851	(8,402 to 11,486)	2787.7	(2392.7 to 3230.7)
			2013	14,651	(11,622 to 18,035)	2423.3	(1949.6 to 2950.9)
		Female	1990	9,415	(7,772 to 10,871)	2226.9	(1839.1 to 2587.2)
			2013	10,178	(8,122 to 13,260)	1429.5	(1146.3 to 1852.9)
		Both	1990	19,267	(17,298 to 21,299)	2474.8	(2225.6 to 2741.5)
					-		

			2013	24,829	(21,221 to 29,124)	1879.8	(1615.8 to 2196.5)
	YLDs	Male	1990	222	(145 to 307)	65.8	(43.1 to 91.6)
			2013	368	(241 to 519)	62.3	(40.6 to 87.8)
		Female	1990	174	(118 to 239)	42.6	(29.1 to 58.7)
			2013	311	(211 to 424)	43.9	(29.9 to 59.7)
		Both	1990	397	(269 to 533)	52.9	(36.0 to 71.7)
			2013	680	(465 to 921)	52.3	(35.7 to 71.2)
Tunisia	Deaths	Male	1990	1,793	(1,417 to 2,267)	116.5	(91.9 to 149.1)
			2013	3,104	(2,182 to 4,252)	87.2	(61.8 to 118.6)
		Female	1990	2,147	(1,577 to 2,619)	140.3	(103.0 to 170.8)
			2013	4,467	(2,874 to 5,903)	104.4	(67.0 to 137.4)
		Both	1990	3,941	(3,259 to 4,647)	128.6	(105.7 to 152.1)
			2013	7,571	(5,774 to 9,396)	97	(73.7 to 119.9)
	DALYs	Male	1990	40,701	(31,515 to 52,541)	1812.7	(1430.8 to 2297.1)
			2013	51,519	(35,248 to 71,403)	1239.5	(858.4 to 1716.8)
		Female	1990	46,937	(33,420 to 58,464)	2106.3	(1546.4 to 2573.0)
			2013	62,753	(40,250 to 86,424)	1341.5	(863.0 to 1837.4)
		Both	1990	87,638	(71,395 to 104,556)	1960.3	(1616.1 to 2320.1)
			2013	114,272	(86,765 to 145,772)	1296.6	(993.7 to 1640.6)
	YLDs	Male	1990	304	(201 to 425)	13.4	(8.9 to 18.7)
			2013	521	(345 to 758)	11.1	(7.3 to 15.9)
		Female	1990	248	(160 to 348)	11	(7.1 to 15.3)
			2013	461	(300 to 655)	9	(5.9 to 12.8)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	552	(373 to 734)	12.2	(8.2 to 16.2)
			2013	983	(669 to 1,382)	10	(6.9 to 14.0)
Turkey	Deaths	Male	1990	17,268	(14,304 to 20,683)	162.5	(134.5 to 194.2)
			2013	18,368	(14,652 to 23,627)	85.3	(67.9 to 109.4)

		Female	1990	18,697	(15,552 to 24,936)	129.7	(107.2 to 172.0)
			2013	26,609	(20,234 to 30,981)	83.7	(63.9 to 97.9)
		Both	1990	35,966	(31,775 to 43,273)	143.3	(126.3 to 171.5)
			2013	44,978	(38,621 to 51,245)	84.7	(72.3 to 96.8)
	DALYs	Male	1990	422,955	(347,485 to 522,130)	2706.2	(2246.8 to 3276.4)
			2013	323,237	(256,139 to 415,571)	1225	(983.3 to 1582.2)
		Female	1990	368,039	(288,121 to 509,986)	2042.2	(1685.2 to 2806.6)
			2013	367,410	(274,618 to 437,797)	1094.9	(821.6 to 1297.7)
		Both	1990	790,995	(671,784 to 974,619)	2339.8	(2049.4 to 2853.7)
			2013	690,647	(578,004 to 801,453)	1153	(970.7 to 1332.3)
	YLDs	Male	1990	1,805	(1,187 to 2,552)	11.6	(7.7 to 16.3)
			2013	3,297	(2,109 to 4,546)	11.2	(7.2 to 15.4)
		Female	1990	1,578	(1,033 to 2,201)	9	(5.8 to 12.7)
			2013	3,045	(2,005 to 4,318)	8.6	(5.7 to 12.3)
		Both	1990	3,384	(2,286 to 4,697)	10.2	(7.0 to 14.1)
			2013	6,343	(4,311 to 8,590)	9.8	(6.6 to 13.2)
Turkmenistan	Deaths	Male	1990	1,286	(1,116 to 1,504)	204.2	(176.1 to 239.2)
			2013	2,352	(1,770 to 3,182)	206.5	(160.7 to 271.1)
		Female	1990	1,558	(1,389 to 1,850)	166.8	(150.0 to 198.7)
			2013	2,228	(1,756 to 3,021)	148.2	(118.4 to 200.1)
		Both	1990	2,844	(2,621 to 3,165)	181.9	(167.5 to 204.3)
			2013	4,581	(3,785 to 5,694)	173.7	(146.3 to 213.4)
	DALYs	Male	1990	33,144	(28,583 to 38,752)	4010.7	(3475.7 to 4677.4)
			2013	59,909	(43,362 to 82,712)	4016.2	(3008.7 to 5447.6)
		Female	1990	33,112	(28,827 to 38,683)	3071.3	(2699.2 to 3595.2)
			2013	45,112	(33,944 to 63,340)	2542.9	(1964.8 to 3505.1)
		Both	1990	66,256	(60,349 to 74,129)	3468	(3182.1 to 3860.0)
			2013	105,021	(83,904 to 133,838)	3202.4	(2605.0 to 4038.3)

	YLDs	Male	1990	146	(97 to 199)	16.5	(11.1 to 22.8)
			2013	264	(174 to 370)	16	(10.6 to 22.2)
		Female	1990	133	(86 to 187)	12.4	(8.0 to 17.5)
			2013	221	(139 to 318)	11.8	(7.5 to 17.3)
		Both	1990	279	(191 to 374)	14.2	(9.8 to 19.1)
			2013	486	(328 to 656)	13.7	(9.2 to 18.9)
Uganda	Deaths	Male	1990	2,949	(2,089 to 3,836)	126.6	(87.9 to 165.6)
			2013	4,362	(3,195 to 5,988)	99.5	(73.0 to 135.1)
		Female	1990	4,137	(2,233 to 5,233)	166	(90.4 to 211.0)
			2013	6,208	(3,848 to 8,126)	130.5	(82.7 to 170.5)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Both	1990	7,087	(4,856 to 8,707)	148.7	(100.8 to 183.4)
			2013	10,571	(8,004 to 13,062)	116.9	(88.0 to 145.4)
	DALYs	Male	1990	76,006	(53,939 to 100,513)	2328.1	(1654.8 to 3037.7)
			2013	97,952	(71,849 to 135,048)	1677.3	(1231.9 to 2322.4)
		Female	1990	90,329	(49,364 to 114,299)	2770.1	(1485.3 to 3508.0)
			2013	116,836	(71,736 to 153,148)	1975.1	(1220.1 to 2599.4)
		Both	1990	166,336	(116,276 to 206,085)	2576.1	(1766.8 to 3166.8)
			2013	214,788	(160,625 to 268,195)	1844.1	(1394.8 to 2286.2)
	YLDs	Male	1990	498	(321 to 708)	13.1	(8.4 to 18.5)
			2013	1,031	(675 to 1,502)	13.8	(9.0 to 20.2)
		Female	1990	364	(231 to 516)	9.2	(6.0 to 13.0)
			2013	717	(464 to 1,054)	9.1	(6.0 to 13.4)
		Both	1990	862	(585 to 1,185)	11	(7.5 to 15.2)
			2013	1,749	(1,169 to 2,444)	11.4	(7.6 to 16.1)
Ukraine	Deaths	Male	1990	39,012	(32,460 to 44,796)	205.6	(170.4 to 237.5)
			2013	44,497	(37,911 to 52,882)	176.7	(150.4 to 210.0)
		Female	1990	79,248	(61,733 to 85,400)	184.1	(143.1 to 198.4)

			2013	65,864	(57,949 to 85,188)	124.8	(110.2 to 160.8)
		Both	1990	118,260	(99,058 to 127,795)	193.1	(160.8 to 208.4)
			2013	110,361	(99,429 to 131,553)	144.7	(130.2 to 171.9)
	DALYs	Male	1990	743,611	(618,153 to 858,990)	3380.2	(2813.1 to 3900.6)
			2013	797,094	(680,359 to 944,166)	3035.9	(2602.4 to 3599.2)
		Female	1990	1,101,062	(860,742 to 1,181,623)	2633.1	(2048.9 to 2828.1)
			2013	836,252	(739,558 to 1,076,318)	1762.9	(1566.7 to 2253.3)
		Both	1990	1,844,674	(1,571,852 to 1,999,064)	2933.5	(2498.7 to 3178.0)
			2013	1,633,346	(1,467,323 to 1,911,644)	2272.7	(2038.5 to 2648.5)
	YLDs	Male	1990	11,912	(7,975 to 17,026)	49.5	(33.6 to 69.7)
			2013	14,572	(9,519 to 20,375)	55.5	(36.4 to 77.2)
		Female	1990	11,869	(7,848 to 16,480)	31.2	(20.7 to 42.5)
			2013	12,133	(8,153 to 17,104)	29.9	(19.8 to 41.4)
		Both	1990	23,782	(16,172 to 32,143)	38.6	(26.5 to 51.9)
			2013	26,706	(18,354 to 36,070)	40.3	(27.8 to 54.7)
United Arab Emirates	Deaths	Male	1990	274	(176 to 416)	141.2	(98.7 to 200.5)
			2013	585	(385 to 857)	108.8	(77.1 to 143.9)
		Female	1990	204	(100 to 306)	178.3	(96.0 to 259.4)
			2013	244	(127 to 381)	109.7	(62.8 to 157.1)
		Both	1990	479	(336 to 648)	159.4	(111.9 to 207.0)
			2013	829	(595 to 1,112)	110.2	(82.9 to 137.4)
	DALYs	Male	1990	9,003	(5,732 to 13,698)	2244.9	(1500.2 to 3335.9)
			2013	21,384	(13,699 to 32,715)	1576.1	(1087.5 to 2149.4)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	4,837	(2,081 to 7,574)	2765.6	(1386.4 to 4189.3)

			2013	7,185	(3,548 to 11,493)	1446.6	(801.7 to 2167.0)
		Both	1990	13,841	(9,529 to 19,270)	2471	(1723.2 to 3292.9)
			2013	28,569	(20,093 to 39,779)	1538.6	(1142.0 to 1967.0)
	YLDs	Male	1990	58	(37 to 84)	10.6	(6.9 to 14.5)
			2013	289	(178 to 419)	10.5	(6.9 to 14.6)
		Female	1990	18	(11 to 25)	8.2	(5.4 to 11.7)
			2013	90	(59 to 131)	9.2	(6.1 to 13.3)
		Both	1990	77	(50 to 106)	9.7	(6.6 to 13.0)
			2013	379	(247 to 535)	10.1	(6.8 to 13.8)
United							
Kingdom	Deaths	Male	1990	27,840	(23,714 to 33,478)	86.6	(73.5 to 104.2)
			2013	22,202	(17,010 to 26,516)	44.4	(34.0 to 52.9)
		Female	1990	48,096	(37,217 to 53,209)	82.3	(64.0 to 91.6)
			2013	34,944	(30,517 to 45,487)	46.4	(40.6 to 60.4)
		Both	1990	75,937	(65,252 to 83,106)	84.9	(73.1 to 93.3)
			2013	57,146	(50,461 to 67,587)	46	(40.9 to 54.0)
	DALYs	Male	1990	449,366	(386,885 to 537,577)	1351.4	(1165.9 to 1613.3)
			2013	311,623	(244,694 to 369,554)	669.7	(527.1 to 794.5)
		Female	1990	567,309	(455,041 to 631,985)	1128.8	(912.2 to 1268.7)
			2013	351,357	(310,354 to 446,756)	572.6	(500.7 to 723.6)
		Both	1990	1,016,675	(901,984 to 1,123,040)	1228.5	(1094.9 to 1360.1)
			2013	662,981	(589,970 to 752,428)	620.8	(550.4 to 702.2)
	YLDs	Male	1990	34,993	(24,685 to 46,264)	104	(73.4 to 137.3)
			2013	54,919	(38,580 to 71,446)	119.7	(83.9 to 155.7)
		Female	1990	36,261	(25,770 to 47,462)	76.7	(54.2 to 100.7)
			2013	46,560	(33,063 to 59,949)	83.9	(59.8 to 108.7)
		Both	1990	71,255	(50,547 to 93,020)	88.5	(62.7 to 115.6)
			2013	101,479	(72,315 to 130,842)	100.6	(71.6 to 129.9)
United States	Deaths	Male	1990	61,606	(50,656 to 78,465)	54.1	(44.3 to 69.2)

			2013	68,227	(51,583 to 85,185)	34.7	(26.3 to 43.2)
		Female	1990	88,130	(73,029 to 110,975)	46.1	(38.5 to 57.8)
			2013	96,005	(73,005 to 123,933)	33.1	(25.2 to 42.3)
		Both	1990	149,736	(133,496 to 176,913)	49.6	(44.3 to 58.6)
			2013	164,233	(133,123 to 194,315)	34	(27.9 to 40.0)
					(1,017,442 to		
	DALYs	Male	1990	1,199,220	1,464,489)	1002.3	(852.4 to 1220.5)
					(1,112,176 to		
			2013	1,358,955	1,617,059)	705.3	(576.4 to 838.6)
		Famala	1000	4 202 502	(1,120,609 to	775 7	(C72 4 + - OF 4 O)
		Female	1990	1,293,582	1,587,055) (1,213,431 to	775.7	(673.4 to 954.9)
					1,766,154)		
			2013	1,478,171	1,700,1547	620.7	(512.7 to 736.3)
				, ,			Uncertainty
							0110011011111
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
Country	measure	sex	year	number	Uncertainty Interval (2,244,755 to	Rate	•
Country	measure	sex Both	year 1990	number 2,492,803	(2,244,755 to 2,864,834)	Rate 873.2	•
Country	measure		1990	2,492,803	(2,244,755 to 2,864,834) (2,415,461 to	873.2	(787.7 to 1005.5)
Country		Both	1990 2013	2,492,803 2,837,127	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780)	873.2 659.7	(787.7 to 1005.5) (560.4 to 757.0)
Country	measure YLDs		1990 2013 1990	2,492,803 2,837,127 178,475	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611)	873.2 659.7 149.4	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2)
Country		Both Male	1990 2013 1990 2013	2,492,803 2,837,127 178,475 394,789	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525)	873.2 659.7 149.4 204.8	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1)
Country		Both	1990 2013 1990	2,492,803 2,837,127 178,475	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611)	873.2 659.7 149.4	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2)
Country		Both Male	1990 2013 1990 2013	2,492,803 2,837,127 178,475 394,789	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525)	873.2 659.7 149.4 204.8	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1)
Country		Both Male	1990 2013 1990 2013 1990	2,492,803 2,837,127 178,475 394,789 207,614	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525) (134,783 to 292,909)	873.2 659.7 149.4 204.8 123.1	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1) (80.2 to 175.5)
Country		Both Male Female	1990 2013 1990 2013 1990 2013	2,492,803 2,837,127 178,475 394,789 207,614 462,207	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525) (134,783 to 292,909) (330,876 to 611,053)	873.2 659.7 149.4 204.8 123.1 203	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1) (80.2 to 175.5) (145.6 to 268.6)
		Both Male Female	1990 2013 1990 2013 1990 2013 1990	2,492,803 2,837,127 178,475 394,789 207,614 462,207 386,089	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525) (134,783 to 292,909) (330,876 to 611,053) (262,598 to 521,404)	873.2 659.7 149.4 204.8 123.1 203 134	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1) (80.2 to 175.5) (145.6 to 268.6) (91.2 to 180.7)
	YLDs	Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013	2,492,803 2,837,127 178,475 394,789 207,614 462,207 386,089 856,996	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525) (134,783 to 292,909) (330,876 to 611,053) (262,598 to 521,404) (611,498 to 1,123,381)	873.2 659.7 149.4 204.8 123.1 203 134 203.2	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1) (80.2 to 175.5) (145.6 to 268.6) (91.2 to 180.7) (144.7 to 267.7)
Uruguay	YLDs	Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013 1990	2,492,803 2,837,127 178,475 394,789 207,614 462,207 386,089 856,996 1,710	(2,244,755 to 2,864,834) (2,415,461 to 3,245,780) (119,534 to 246,611) (279,181 to 511,525) (134,783 to 292,909) (330,876 to 611,053) (262,598 to 521,404) (611,498 to 1,123,381) (1,457 to 2,011)	873.2 659.7 149.4 204.8 123.1 203 134 203.2 124.7	(787.7 to 1005.5) (560.4 to 757.0) (100.4 to 205.2) (144.5 to 265.1) (80.2 to 175.5) (145.6 to 268.6) (91.2 to 180.7) (144.7 to 267.7) (105.9 to 146.7)

		Both	1990	3,919	(3,546 to 4,551)	115	(104.0 to 133.4)
			2013	4,113	(3,575 to 4,915)	78.8	(68.5 to 92.8)
	DALYs	Male	1990	30,090	(25,836 to 35,323)	2046.4	(1766.2 to 2398.4)
			2013	26,358	(21,041 to 31,962)	1433.5	(1142.0 to 1741.6)
		Female	1990	30,111	(26,863 to 37,159)	1544.9	(1375.6 to 1902.4)
			2013	25,415	(20,608 to 32,783)	915.8	(733.7 to 1179.5)
		Both	1990	60,201	(55,115 to 68,893)	1767.3	(1616.3 to 2018.2)
			2013	51,773	(44,898 to 60,015)	1137.6	(982.3 to 1316.3)
	YLDs	Male	1990	679	(456 to 945)	44.9	(30.1 to 62.5)
			2013	781	(520 to 1,104)	43	(28.7 to 60.8)
		Female	1990	485	(325 to 672)	25.5	(17.2 to 35.3)
			2013	587	(394 to 800)	24.5	(16.7 to 33.3)
		Both	1990	1,164	(798 to 1,554)	34	(23.4 to 45.3)
			2013	1,368	(945 to 1,816)	32.5	(22.5 to 43.2)
Uzbekistan	Deaths	Male	1990	6,855	(5,930 to 7,995)	189.2	(163.9 to 220.5)
			2013	12,887	(9,902 to 16,747)	203.7	(159.0 to 258.4)
		Female	1990	8,731	(7,844 to 10,297)	156.9	(141.1 to 185.2)
			2013	12,786	(9,866 to 16,046)	154.9	(119.9 to 192.4)
		Both	1990	15,587	(14,284 to 17,404)	169.9	(155.6 to 190.5)
			2013	25,673	(21,758 to 30,875)	177.2	(150.4 to 210.1)
	DALYs	Male	1990	144,118	(124,033 to 167,684)	3355.9	(2896.7 to 3886.0)
			2013	278,473	(207,371 to 373,536)	3564.3	(2726.2 to 4689.6)
		Female	1990	148,341	(129,572 to 170,862)	2549.6	(2245.4 to 2949.0)
			2013	211,172	(157,174 to 280,294)	2334.5	(1757.6 to 3032.1)
		Both	1990	292,460	(266,997 to 324,399)	2890.8	(2651.9 to 3220.4)
			2013	489,646	(403,778 to 603,288)	2903.5	(2438.1 to 3535.1)
	YLDs	Male	1990	849	(569 to 1,188)	16.8	(11.4 to 23.8)
			2013	1,535	(1,017 to 2,118)	16.7	(11.0 to 23.0)

		Female	1990	776	(519 to 1,077)	12.6	(8.3 to 17.6)
Country	measure	sex	year	number	Uncertainty Interval	Rate	Uncertainty Interval
			2013	1,253	(835 to 1,815)	12.3	(8.2 to 17.8)
		Both	1990	1,626	(1,094 to 2,245)	14.5	(9.8 to 19.9)
			2013	2,788	(1,913 to 3,798)	14.4	(9.9 to 19.7)
Vanuatu	Deaths	Male	1990	66	(43 to 92)	205.3	(136.8 to 276.6)
			2013	118	(76 to 166)	209.9	(140.7 to 279.8)
		Female	1990	33	(21 to 51)	131.6	(85.4 to 199.4)
			2013	83	(48 to 127)	158.6	(97.8 to 230.4)
		Both	1990	100	(68 to 139)	172.3	(122.4 to 233.1)
			2013	202	(130 to 287)	184.2	(125.3 to 249.0)
	DALYs	Male	1990	1,671	(1,045 to 2,435)	4311.8	(2748.8 to 6149.1)
			2013	3,159	(1,990 to 4,732)	4299.3	(2763.8 to 6129.1)
		Female	1990	794	(479 to 1,259)	2532.5	(1572.6 to 3943.5)
			2013	2,011	(1,081 to 3,211)	2923.8	(1654.5 to 4502.9)
		Both	1990	2,466	(1,579 to 3,641)	3493	(2302.1 to 5028.1)
			2013	5,170	(3,183 to 7,750)	3610.9	(2304.6 to 5212.7)
	YLDs	Male	1990	19	(13 to 27)	58.3	(39.0 to 79.7)
			2013	36	(23 to 50)	58.5	(39.0 to 80.3)
		Female	1990	11	(7 to 15)	40.6	(27.8 to 55.3)
			2013	26	(17 to 36)	42.7	(28.4 to 59.8)
		Both	1990	31	(21 to 42)	50.4	(34.1 to 67.9)
			2013	62	(42 to 84)	50.6	(34.4 to 67.8)
Venezuela	Deaths	Male	1990	3,229	(2,722 to 3,869)	95.4	(80.3 to 114.2)
			2013	6,458	(5,244 to 7,892)	73.9	(59.6 to 90.8)
		Female	1990	3,532	(3,008 to 4,271)	88.7	(75.7 to 107.3)
			2013	6,532	(5,059 to 7,839)	65.2	(50.0 to 78.1)
		Both	1990	6,762	(6,099 to 7,609)	92	(82.8 to 103.9)

			2013	12,991	(11,173 to 14,756)	69.5	(59.6 to 79.5)
	DALYs	Male	1990	71,687	(61,003 to 86,183)	1582.1	(1340.9 to 1894.1)
			2013	129,311	(103,919 to 157,907)	1210.2	(985.9 to 1465.7)
		Female	1990	70,405	(58,771 to 82,695)	1390.1	(1181.4 to 1660.1)
			2013	104,858	(84,417 to 125,570)	914.1	(734.2 to 1092.8)
		Both	1990	142,092	(127,114 to 160,566)	1480	(1328.9 to 1661.4)
			2013	234,169	(201,048 to 267,623)	1057.1	(913.9 to 1202.3)
	YLDs	Male	1990	1,319	(869 to 1,832)	27.6	(18.3 to 38.2)
			2013	3,091	(2,019 to 4,234)	27.9	(18.5 to 38.4)
		Female	1990	971	(655 to 1,383)	18.8	(12.5 to 27.2)
			2013	2,242	(1,479 to 3,136)	18.7	(12.5 to 26.0)
		Both	1990	2,290	(1,582 to 3,080)	22.9	(16.0 to 30.7)
			2013	5,333	(3,598 to 7,204)	23.1	(15.5 to 31.0)
Vietnam	Deaths	Male	1990	47,383	(39,465 to 55,004)	308.5	(259.5 to 353.8)
			2013	63,146	(48,875 to 81,044)	238.4	(186.2 to 304.5)
		Female	1990	57,247	(47,278 to 69,269)	242.2	(199.9 to 293.1)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
			2013	82,587	(66,342 to 97,666)	170.6	(135.5 to 205.2)
		Both	1990	104,631	(91,614 to 118,032)	272.7	(239.1 to 306.9)
			2013	145,734	(123,670 to 169,740)	198.7	(168.7 to 232.3)
	DALYs	Male	1990	1,058,216	(859,893 to 1,263,728)	5934.3	(4885.3 to 7011.3)
			2013	1,158,336	(848,611 to 1,549,263)	3799.4	(2821.9 to 5066.9)
		Female	1990	891,091	(718,272 to 1,103,721)	3803.7	(3068.5 to 4709.5)
			2013	949,834	(732,303 to 1,199,392)	2155	(1646.9 to 2768.8)
					(1,673,659 to		
		Both	1990	1,949,307	2,232,714)	4771.4	(4105.9 to 5434.5)
					(1,719,529 to		(
			2013	2,108,170	2,565,870)	2863.1	(2339.9 to 3487.0)

			2013	4,531	(3,528 to 5,686)	230.6	(181.0 to 286.7)
		Female	1990	2,000	(1,559 to 2,382)	178.1	(137.6 to 212.2)
			2013	2,215	(1,709 to 2,878)	130.4	(101.1 to 167.0)
Zambia	Deaths	Male	1990	530	(385 to 674)	53.6	(38.0 to 68.4)
			2013	1,066	(725 to 1,473)	9.7	(6.6 to 13.2)
		Both	1990	448	(304 to 613)	9.8	(6.7 to 13.4)
			2013	488	(314 to 695)	8.6	(5.7 to 12.3)
		Female	1990	217	(141 to 311)	8.6	(5.7 to 12.3)
			2013	578	(368 to 809)	10.8	(6.8 to 15.2)
	YLDs	Male	1990	231	(154 to 319)	11.3	(7.7 to 15.9)
			2013	397,121	(252,476 to 624,272)	3678.9	(2399.1 to 5690.0)
		Both	1990	251,176	(163,131 to 362,256)	4082.7	(2747.0 to 6154.3)
			2013	216,788	(131,158 to 355,866)	3950.2	(2449.2 to 6304.9)
		Female	1990	137,946	(78,516 to 209,276)	4191.4	(2634.1 to 6515.8)
			2013	180,332	(109,306 to 278,566)	3378.4	(2135.3 to 5100.9)
	DALYs	Male	1990	113,229	(73,231 to 166,545)	3941.5	(2611.3 to 5902.5)
			2013	15,307	(10,163 to 23,273)	207.2	(145.1 to 295.9)
		Both	1990	7,746	(5,305 to 11,450)	212.5	(150.5 to 309.4)
			2013	8,511	(5,366 to 13,360)	220.2	(146.1 to 329.9)
		Female	1990	4,473	(2,906 to 6,799)	217.1	(150.3 to 327.2)
			2013	6,796	(4,365 to 10,153)	192.9	(132.2 to 270.9)
Yemen	Deaths	Male	1990	3,272	(2,223 to 4,755)	206.3	(146.2 to 289.5)
			2013	36,009	(25,359 to 48,878)	49.3	(34.9 to 67.1)
		Both	1990	21,762	(15,505 to 28,245)	53.1	(37.8 to 69.3)
			2013	18,631	(12,542 to 25,997)	43.3	(29.6 to 60.7)
		Female	1990	10,902	(7,636 to 14,310)	45.9	(32.2 to 60.5)
			2013	17,378	(11,849 to 24,156)	57.6	(39.3 to 80.8)
	YLDs	Male	1990	10,859	(7,436 to 14,659)	63.4	(43.7 to 85.4)

		Both	1990	2,531	(2,037 to 2,924)	123.1	(97.8 to 144.7)
			2013	6,746	(5,598 to 8,096)	186.3	(155.5 to 221.4)
	DALYs	Male	1990	14,763	(10,960 to 19,047)	867.5	(633.8 to 1106.6)
			2013	47,073	(35,437 to 61,670)	2100.6	(1601.2 to 2733.2)
							Uncertainty
Country	measure	sex	year	number	Uncertainty Interval	Rate	Interval
		Female	1990	40,408	(31,684 to 48,441)	2754.9	(2142.6 to 3287.0)
			2013	84,031	(64,524 to 107,494)	3572.9	(2735.7 to 4531.2)
		Both	1990	55,172	(44,923 to 64,288)	1904.2	(1539.0 to 2204.7)
			2013	131,105	(106,690 to 157,261)	2902.4	(2388.8 to 3504.9)
	YLDs	Male	1990	222	(140 to 325)	13.1	(8.0 to 19.3)
			2013	394	(266 to 581)	12.8	(8.4 to 19.2)
		Female	1990	166	(107 to 243)	9.1	(5.9 to 13.3)
			2013	278	(174 to 421)	8.7	(5.4 to 13.8)
		Both	1990	389	(255 to 550)	11	(7.2 to 15.6)
			2013	673	(453 to 940)	10.7	(7.1 to 15.1)
Zimbabwe	Deaths	Male	1990	1,222	(974 to 1,611)	78.9	(62.4 to 106.0)
			2013	2,736	(1,798 to 3,745)	102.3	(65.0 to 140.6)
		Female	1990	2,400	(1,934 to 2,841)	143.9	(115.2 to 170.3)
			2013	4,280	(2,972 to 6,230)	128.4	(88.4 to 190.8)
		Both	1990	3,622	(3,150 to 4,200)	114.6	(99.5 to 132.8)
			2013	7,016	(5,361 to 8,999)	117.7	(88.8 to 152.9)
	DALYs	Male	1990	32,066	(25,579 to 42,249)	1464	(1172.8 to 1919.0)
			2013	74,853	(50,737 to 100,938)	1977.2	(1294.1 to 2723.6)
		Female	1990	52,145	(41,461 to 62,548)	2449	(1958.6 to 2891.2)
			2013	103,349	(70,757 to 143,609)	2257.6	(1550.1 to 3362.5)
		Both	1990	84,211	(72,757 to 99,460)	1992.4	(1726.9 to 2307.5)
			2013	178,202	(131,388 to 224,574)	2137.6	(1611.7 to 2764.0)
	YLDs	Male	1990	479	(320 to 652)	22.3	(14.5 to 30.7)

		2013	678	(452 to 945)	22.1	(14.4 to 31.0)
	Female	1990	506	(332 to 704)	21.7	(14.4 to 30.2)
		2013	809	(536 to 1,144)	22.3	(14.6 to 31.5)
	Both	1990	985	(674 to 1,324)	22	(15.2 to 30.0)
		2013	1,487	(1,012 to 2,020)	22.3	(15.0 to 30.3)

Web Appendix B. Country-specific age-standardized stroke burden per 100 000 in 2013 for males and females stratified by stroke type.

					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95%UI
Afghanistan	Deaths	Ischemic Stroke	Male	1990	1,014	(502 to 1,637)	112.1	(59.7 to 166.3)

			2013	3,545	(1,607 to 5,906)	116	(56.4 to 180.0)
		Female	1990	1,657	(878 to 2,810)	133	(77.6 to 194.8)
			2013	5,588	(2,818 to 9,586)	145.3	(78.2 to 220.0)
		Both	1990	2,672	(1,619 to 3,829)	123.8	(82.6 to 165.8)
			2013	9,133	(5,010 to 13,559)	132.2	(78.2 to 180.3)
	Hemorrhagic			-	,		,
Deaths	Stroke	Male	1990	1,343	(755 to 2,148)	105.3	(60.0 to 168.8)
			2013	4,716	(2,630 to 7,665)	115.1	(67.0 to 181.7)
		Female	1990	2,902	(1,284 to 5,445)	167	(84.4 to 296.0)
			2013	9,427	(4,480 to 18,278)	184.6	(97.0 to 344.6)
		Both	1990	4,245	(2,503 to 6,899)	139	(90.1 to 216.3)
			2013	14,144	(8,693 to 23,874)	152.1	(98.7 to 245.2)
DALYs	Ischemic Stroke	Male	1990	24,020	(11,076 to 42,231)	1807.4	(929.9 to 2808.7)
					(33,404 to		,
			2013	77,306	138,329)	1833.5	(841.7 to 3013.9)
		Female	1990	40,181	(19,212 to 76,378)	2288.3	(1253.0 to 3822.5)
					(56,702 to		
			2013	122,347	243,732)	2429.1	(1218.3 to 4213.1)
					(35,584 to		
		Both	1990	64,201	101,799)	2068.5	(1291.0 to 2889.0)
					(102,000 to		
			2013	199,654	320,874)	2149.3	(1186.6 to 3174.9)
	Hemorrhagic		4000	40.006	(22.205 (6.672)	2420	(4247.61.2465.7)
	Stroke	Male	1990	40,996	(22,305 to 66,672)	2138	(1217.6 to 3465.7)
			2013	132,739	(72,666 to 217,256)	2319.3	(1298.4 to 3760.8)
			2013	132,/39	(38,808 to	2319.3	(1230.4 (0 3/00.8)
		Female	1990	92,134	173,694)	3814.5	(1719.0 to 7117.5)
		· ciliaic	1330	32,131	(123,167 to	301.13	(=, 13.0 to , 11)
			2013	280,419	552,508)	4192.5	(1981.0 to 8144.9)

						(75,853 to		
			Both	1990	133,131	217,798)	3022.5	(1807.6 to 4835.6)
						(244,179 to		(22.2.2.)
				2013	413,158	700,287)	3282.7	(2017.3 to 5528.6)
	YLDs	Ischemic Stroke	Male	1990	180	(115 to 260)	9.2	(5.9 to 13.4)
				2013	564	(368 to 815)	9.6	(6.4 to 13.5)
			Female	1990	153	(95 to 223)	7.3	(4.7 to 10.5)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	468	(302 to 666)	7.6	(5.0 to 10.7)
			Both	1990	334	(215 to 477)	8.2	(5.4 to 11.4)
				2013	1,032	(695 to 1,431)	8.5	(5.9 to 11.8)
		Hemorrhagic						
		Stroke	Male	1990	31	(16 to 52)	1.2	(0.7 to 2.0)
				2013	87	(51 to 137)	1.3	(0.8 to 2.0)
			Female	1990	34	(20 to 52)	1.4	(0.8 to 2.0)
				2013	85	(46 to 132)	1.2	(0.7 to 1.9)
			Both	1990	66	(41 to 95)	1.3	(0.8 to 1.8)
			Both	2013	172	(107 to 253)	1.2	(0.8 to 1.8)
Albania	Deaths	Ischemic Stroke	Male	1990	303	(219 to 364)	43.4	(31.2 to 52.7)
				2013	663	(420 to 1,033)	45.2	(28.4 to 71.2)
			Female	1990	320	(238 to 387)	36.2	(26.6 to 43.6)
				2013	585	(329 to 832)	32.4	(18.4 to 45.9)
			Both	1990	624	(513 to 714)	39.4	(32.1 to 45.2)
				2013	1,248	(836 to 1,678)	38.3	(25.5 to 51.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,118	(989 to 1,280)	150.8	(133.5 to 173.4)
				2013	2,549	(1,958 to 3,250)	171	(132.1 to 214.0)
			Female	1990	1,290	(1,170 to 1,520)	142.5	(129.0 to 168.7)

				2013	2,471	(1,985 to 3,105)	137	(110.0 to 171.7)
			Both	1990	2,408	(2,237 to 2,700)	146.7	(135.2 to 165.0)
				2013	5,021	(4,226 to 5,963)	153.3	(129.9 to 181.2)
	DALYs	Ischemic Stroke	Male	1990	5,240	(3,855 to 6,369)	595.9	(433.8 to 717.1)
				2013	10,014	(6,510 to 15,222)	621.5	(406.6 to 945.5)
			Female	1990	4,685	(3,752 to 5,832)	467.9	(368.8 to 572.0)
				2013	7,131	(4,199 to 10,134)	393.4	(232.3 to 558.8)
			Both	1990	9,925	(8,342 to 11,512)	526.9	(437.2 to 607.4)
				2013	17,145	(11,763 to 23,096)	500.8	(345.3 to 671.5)
		Hemorrhagic						
		Stroke	Male	1990	21,568	(18,775 to 24,643)	2247.2	(1981.3 to 2562.7)
				2013	41,137	(30,705 to 53,818)	2533.1	(1917.9 to 3283.7)
			Female	1990	20,475	(18,337 to 24,109)	1977	(1777.6 to 2301.5)
				2013	31,400	(24,545 to 41,096)	1745.8	(1364.9 to 2279.5)
			Both	1990	42,044	(38,743 to 47,003)	2108.5	(1945.1 to 2344.9)
				2013	72,538	(59,562 to 87,117)	2124.4	(1749.7 to 2550.7)
	YLDs	Ischemic Stroke	Male	1990	238	(150 to 341)	22	(14.1 to 32.0)
				2013	424	(279 to 619)	24.7	(16.3 to 35.7)
			Female	1990	158	(102 to 226)	14.6	(9.3 to 20.9)
							14.3	
				2012	250	(165 to 385)		(0.2+24.4)
				2013	259	at a Allega and	Data	(9.2 to 21.1)
6		Charles Torre				nts, All ages		100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	396	(270 to 549)	18.2	(12.5 to 25.2)
		Homourbee's		2013	683	(454 to 957)	19.3	(12.9 to 27.0)
		Hemorrhagic Stroke	Male	1990	67	(39 to 106)	5.4	(3.1 to 8.5)
		JUOKC	IVIUIC	1000	0,	(33 to 100)	J.7	(3.1 to 0.5)

				2013	102	(62 to 154)	6	(3.7 to 9.2)
			Female	1990	49	(30 to 72)	4.1	(2.5 to 6.0)
				2013	71	(45 to 105)	4	(2.6 to 5.9)
			Both	1990	117	(76 to 166)	4.7	(3.1 to 6.7)
			Both	2013	174	(113 to 250)	5	(3.3 to 7.1)
Algeria	Deaths	Ischemic Stroke	Male	1990	2,894	(2,256 to 3,702)	89.8	(70.7 to 113.3)
				2013	5,553	(4,394 to 6,796)	69.1	(55.0 to 84.7)
			Female	1990	3,614	(2,981 to 4,598)	88.1	(72.2 to 110.3)
				2013	6,692	(5,295 to 8,455)	74.1	(58.0 to 92.8)
			Both	1990	6,508	(5,554 to 7,837)	88.9	(76.3 to 105.3)
				2013	12,246	(10,329 to 14,229)	71.9	(60.9 to 83.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,787	(2,110 to 3,523)	60.4	(43.7 to 78.3)
				2013	4,598	(3,541 to 5,803)	46.5	(35.3 to 59.5)
			Female	1990	3,654	(2,874 to 4,640)	66.3	(51.3 to 84.1)
				2013	5,601	(4,488 to 7,075)	52	(40.9 to 67.3)
			Both	1990	6,441	(5,335 to 7,648)	63.5	(51.9 to 75.5)
				2013	10,200	(8,713 to 12,092)	49.3	(42.0 to 58.8)
	DALYs	Ischemic Stroke	Male	1990	49,172	(37,261 to 65,122)	1197.6	(929.4 to 1542.4)
						(65,413 to		
				2013	85,358	105,207)	863.8	(677.0 to 1059.8)
			Female	1990	55,823	(45,248 to 72,912)	1131.1	(928.7 to 1439.8)
						(73,485 to		
				2013	93,596	118,906)	888.1	(702.7 to 1118.5)
						(88,708 to		
			Both	1990	104,996	131,084)	1161.4	(986.5 to 1409.6)
				2012	470.0	(149,241 to	07	(700 0 t 1000 t)
		11		2013	178,955	210,402)	877.7	(739.2 to 1023.1)
		Hemorrhagic	Mala	1000	02.471	(70,903 to	1220.0	(02F F to 1FFF 2)
		Stroke	Male	1990	92,471	118,982)	1220.9	(925.5 to 1555.3)

						(86,919 to		
				2013	112,617	142,811)	842.6	(653.8 to 1055.8)
						(87,426 to		
			Female	1990	115,984	157,031)	1402	(1087.2 to 1779.8)
						(106,682 to		(
				2013	132,233	167,875)	957.4	(771.5 to 1222.5)
						(169,268 to		
						251,244)		
			Both	1990	208,456		1313.3	(1082.1 to 1568.7)
			Dotti	1330	200,430		1313.3	(1002.1 to 1500.7)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				·		(207,936 to		
				2013	244,851	291,375)	900.5	(770.2 to 1064.9)
	YLDs	Ischemic Stroke	Male	1990	524	(343 to 745)	9.3	(6.1 to 13.2)
				2013	1,371	(856 to 1,939)	10.3	(6.6 to 14.6)
			Female	1990	485	(323 to 695)	7.9	(5.2 to 11.3)
				2013	1,022	(656 to 1,446)	7.6	(5.0 to 10.7)
			Both	1990	1,010	(686 to 1,390)	8.6	(5.8 to 11.7)
				2013	2,393	(1,580 to 3,262)	8.9	(5.9 to 12.3)
		Hemorrhagic						
		Stroke	Male	1990	89	(53 to 139)	1.3	(0.8 to 2.1)
				2013	178	(103 to 284)	1.2	(0.7 to 2.0)
			Female	1990	95	(55 to 147)	1.4	(0.8 to 2.1)
				2013	183	(109 to 279)	1.3	(0.8 to 1.9)
			Both	1990	184	(116 to 264)	1.3	(0.8 to 1.9)
			Both	2013	361	(228 to 528)	1.2	(0.8 to 1.8)
Andorra	Deaths	Ischemic Stroke	Male	1990	12	(8 to 17)	47.7	(33.6 to 67.0)
				2013	18	(13 to 26)	29.3	(20.4 to 41.4)

_								
			Female	1990	21	(13 to 29)	41.8	(26.6 to 57.9)
				2013	29	(17 to 44)	25.7	(15.3 to 38.9)
			Both	1990	33	(25 to 43)	44.5	(33.9 to 56.8)
				2013	48	(35 to 65)	27.5	(20.3 to 37.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	6	(3 to 10)	25.6	(14.6 to 40.1)
				2013	11	(6 to 16)	17.9	(10.7 to 26.3)
			Female	1990	10	(4 to 15)	21	(10.4 to 32.5)
				2013	15	(7 to 24)	14.5	(7.4 to 23.5)
			Both	1990	16	(9 to 23)	23.1	(13.7 to 32.4)
				2013	26	(17 to 37)	16.1	(10.4 to 22.3)
	DALYs	Ischemic Stroke	Male	1990	166	(119 to 227)	615.5	(442.1 to 842.3)
				2013	225	(165 to 301)	365.4	(269.0 to 487.8)
			Female	1990	194	(130 to 262)	415.2	(279.7 to 566.5)
				2013	232	(151 to 338)	241.9	(159.0 to 351.2)
			Both	1990	361	(284 to 457)	499.2	(391.1 to 634.5)
				2013	458	(357 to 603)	297.2	(231.9 to 389.8)
		Hemorrhagic						
		Stroke	Male	1990	116	(71 to 177)	423.6	(259.6 to 641.6)
				2013	152	(91 to 225)	262.9	(159.8 to 391.8)
			Female	1990	111	(57 to 167)	275.7	(143.5 to 408.5)
				2013	133	(74 to 208)	164.4	(93.5 to 248.3)
								(221.8 to 462.5)
			D. H	4000	220	(4.45) . 200)	242.2	
			Both	1990	228	(145 to 306)	342.2	
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

YLDS Ischemic Stroke Male 1990 25 (16 to 35) 91.7 (60.6 to 125.6)									
2013 49 (32 to 69) 85.2 (55.8 to 118.2)					2013	286	(194 to 380)	210.3	(146.2 to 278.6)
Female 1990 26		YLDs	Ischemic Stroke	Male	1990	25	(16 to 35)	91.7	(60.6 to 125.6)
Both 1990 51 (34 to 70) 76 (51.1 to 103.4)					2013	49	(32 to 69)	85.2	(55.8 to 118.2)
Both 1990 51 (34 to 70) 76 (51.1 to 103.4) 2013 94 (63 to 127) 71.2 (47.8 to 95.8)				Female	1990	26	(16 to 37)	64.7	(41.2 to 94.1)
Hemorrhagic Stroke Male 1990 5 (2 to 9) 19.1 (9.7 to 32.8)					2013	44	(28 to 61)	59.4	(38.4 to 82.1)
Hemorrhagic Stroke Male 1990 5 (2 to 9) 19.1 (9.7 to 32.8)				Both	1990	51	(34 to 70)	76	(51.1 to 103.4)
Stroke Male 1990 5 (2 to 9) 19.1 (9.7 to 32.8) 2013 10 (5 to 18) 17.7 (9.6 to 31.5) 4 2013 10 (5 to 18) 17.7 (9.6 to 31.5) 5 2013 9 (4 to 16) 13.5 (7.2 to 23.2) 6 Both 1990 9 (5 to 14) 15.2 (9.3 to 22.6) 8 Both 2013 19 (11 to 29) 15.4 (9.5 to 22.9) Angola Deaths Ischemic Stroke Male 1990 1,296 (756 to 1,904) 109.6 (65.5 to 159.5) 8 2013 2,352 (1,552 to 3,563) 106.5 (69.8 to 154.4) 9 1,431 (854 to 2,144) 109.3 (67.2 to 156.5) 1 2013 2,851 (1,925 to 4,327) 112.8 (79.1 to 167.9) 1 Both 1990 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) 1 2013 5,204 (3,77		_			2013	94	(63 to 127)	71.2	(47.8 to 95.8)
2013 10			Hemorrhagic						
Female 1990 4			Stroke	Male	1990	5	(2 to 9)	19.1	(9.7 to 32.8)
Both 1990 9 (5 to 14) 15.2 (9.3 to 22.6)					2013	10	(5 to 18)	17.7	(9.6 to 31.5)
Both 1990 9 (5 to 14) 15.2 (9.3 to 22.6)				Female	1990	4	(2 to 7)	12.5	(7.0 to 20.1)
Angola Deaths Ischemic Stroke Male 1990 1,296 (756 to 1,904) 109.6 (65.5 to 159.5) Angola Deaths Ischemic Stroke Male 1990 1,296 (756 to 1,904) 109.6 (65.5 to 159.5) 2013 2,352 (1,552 to 3,563) 106.5 (69.8 to 154.4) 109.3 (67.2 to 156.5) (67.2 to 156.5) 2013 2,851 (1,925 to 4,327) 112.8 (79.1 to 167.9) 109.0 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) 109.0 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) 109.0 2,727 (1,722 to 3,943) 110.6 (82.3 to 159.3) 109.0 1,792 (1,080 to 2,578) 10.6 (82.3 to 159.3) 109.0 1,792 (1,080 to 2,578) 10.5 (59.3 to 151.4) 109.0 2,083 (1,294 to 4,429) 91.1 (58.5 to 136.3) 109.0 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 109.0					2013	9	(4 to 16)	13.5	(7.2 to 23.2)
Angola Deaths Ischemic Stroke Male 1990 1,296 (756 to 1,904) 109.6 (65.5 to 159.5) 2013 2,352 (1,552 to 3,563) 106.5 (69.8 to 154.4) 1 Female 1990 1,431 (854 to 2,144) 109.3 (67.2 to 156.5) 2013 2,851 (1,925 to 4,327) 112.8 (79.1 to 167.9) Both 1990 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) Deaths Stroke Male 1990 1,792 (1,080 to 2,578) 10.5 (59.3 to 151.4) Deaths Stroke Male 1990 2,083 (1,274 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,				Both	1990	9	(5 to 14)	15.2	(9.3 to 22.6)
2013 2,352 (1,552 to 3,563) 106.5 (69.8 to 154.4)				Both	2013	19	(11 to 29)	15.4	(9.5 to 22.9)
Female 1990 1,431 (854 to 2,144) 109.3 (67.2 to 156.5) 2013 2,851 (1,925 to 4,327) 112.8 (79.1 to 167.9) Both 1990 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) 2013 5,204 (3,778 to 7,794) 110.6 (82.3 to 159.3) Hemorrhagic Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4) 2013 2,885 (1,974 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYS Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)	Angola	Deaths	Ischemic Stroke	Male	1990	1,296	(756 to 1,904)	109.6	(65.5 to 159.5)
2013 2,851 (1,925 to 4,327) 112.8 (79.1 to 167.9)					2013	2,352	(1,552 to 3,563)	106.5	(69.8 to 154.4)
Both 1990 2,727 (1,722 to 3,943) 110.2 (72.4 to 155.1) Deaths Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4) Deaths Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4) Deaths Stroke Male 1990 2,885 (1,974 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) Deaths Both 1990 3,875 (2,466 to 5,588) 98 (61.0 to 154.4) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)				Female	1990	1,431	(854 to 2,144)	109.3	(67.2 to 156.5)
Deaths Deaths Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4)					2013	2,851	(1,925 to 4,327)	112.8	(79.1 to 167.9)
Deaths Hemorrhagic Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4) 2013 2,885 (1,974 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)				Both	1990	2,727	(1,722 to 3,943)	110.2	(72.4 to 155.1)
Deaths Stroke Male 1990 1,792 (1,080 to 2,578) 105 (59.3 to 151.4) 2013 2,885 (1,974 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)					2013	5,204	(3,778 to 7,794)	110.6	(82.3 to 159.3)
2013 2,885 (1,974 to 4,429) 91.1 (58.5 to 136.3) Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYS Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)			Hemorrhagic						
Female 1990 2,083 (1,282 to 3,179) 115.1 (71.7 to 173.9) 2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4) Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)		Deaths	Stroke	Male	1990	1,792	(1,080 to 2,578)	105	(59.3 to 151.4)
2013 3,329 (2,095 to 5,268) 98 (61.0 to 154.4)					2013	2,885	(1,974 to 4,429)	91.1	(58.5 to 136.3)
Both 1990 3,875 (2,466 to 5,588) 111.4 (70.9 to 158.6) 2013 6,214 (4,254 to 9,353) 95.3 (63.7 to 142.1) DALYs Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)				Female	1990	2,083	(1,282 to 3,179)	115.1	(71.7 to 173.9)
DALYS Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)					2013	3,329	(2,095 to 5,268)	98	(61.0 to 154.4)
DALYS Ischemic Stroke Male 1990 31,048 (17,701 to 47,645) 1842.1 (1062.9 to 2697.6)				Both	1990	3,875	(2,466 to 5,588)	111.4	(70.9 to 158.6)
					2013	6,214	(4,254 to 9,353)	95.3	(63.7 to 142.1)
2013 50,669 (32,996 to 78,962) 1645.8 (1080.7 to 2498.2)		DALYs	Ischemic Stroke	Male	1990	31,048	(17,701 to 47,645)	1842.1	(1062.9 to 2697.6)
					2013	50,669	(32,996 to 78,962)	1645.8	(1080.7 to 2498.2)

			Female	1990	28,504	(15,743 to 46,315)	1593.9	(952.2 to 2384.1)
				2013	51,370	(33,383 to 82,521)	1559.1	(1044.9 to 2381.9)
			Both	1990	59,552	(36,912 to 86,597)	1716.6	(1086.9 to 2471.6)
						(71,527 to		
				2013	102,039	155,613)	1606	(1162.6 to 2417.7)
		Hemorrhagic						
		Stroke	Male	1990	59,406	(36,375 to 84,682)	2440.5	(1441.2 to 3521.6)
						(63,322 to		
				2013	90,628	136,738)	1968.2	(1336.3 to 3046.6)
			Female	1990	60,851	(35,988 to 92,656)	2389.9	(1457.6 to 3653.9)
						(57,339 to		
						146,727)		
				2012	01 211		1020.1	/420F 0 to 2002 0\
				2013	91,211		1928.1	(1205.0 to 3082.8)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Country	measure	Stroke Type		·		(78,496 to		
Country	measure	Stroke Type	sex Both	year 1990	Mean 120,258	(78,496 to 170,460)	Mean 2425.4	95% UI (1525.0 to 3506.0)
Country	measure	Stroke Type		1990	120,258	(78,496 to 170,460) (126,699 to	2425.4	(1525.0 to 3506.0)
Country			Both	1990 2013	120,258 181,840	(78,496 to 170,460) (126,699 to 272,666)	2425.4 1954.2	(1525.0 to 3506.0) (1329.8 to 2973.6)
Country	measure YLDs	Stroke Type Ischemic Stroke		1990 2013 1990	120,258 181,840 278	(78,496 to 170,460) (126,699 to 272,666) (176 to 412)	2425.4 1954.2 14.1	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6)
Country			Both Male	1990 2013 1990 2013	120,258 181,840 278 568	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812)	2425.4 1954.2 14.1 14.2	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1)
Country			Both	1990 2013 1990 2013 1990	120,258 181,840 278 568 239	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337)	2425.4 1954.2 14.1 14.2 10.6	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9)
Country			Both Male	1990 2013 1990 2013	120,258 181,840 278 568	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812)	2425.4 1954.2 14.1 14.2	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1)
Country			Both Male	1990 2013 1990 2013 1990	120,258 181,840 278 568 239	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337)	2425.4 1954.2 14.1 14.2 10.6	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9)
Country			Both Male Female	1990 2013 1990 2013 1990 2013	120,258 181,840 278 568 239 533	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337) (351 to 765)	2425.4 1954.2 14.1 14.2 10.6 11.9	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9) (7.7 to 17.1)
Country			Both Male Female	1990 2013 1990 2013 1990 2013 1990	120,258 181,840 278 568 239 533 517	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337) (351 to 765) (341 to 725)	2425.4 1954.2 14.1 14.2 10.6 11.9 12.1	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9) (7.7 to 17.1) (8.0 to 16.8)
Country		Ischemic Stroke	Both Male Female	1990 2013 1990 2013 1990 2013 1990	120,258 181,840 278 568 239 533 517	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337) (351 to 765) (341 to 725)	2425.4 1954.2 14.1 14.2 10.6 11.9 12.1	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9) (7.7 to 17.1) (8.0 to 16.8)
Country		Ischemic Stroke Hemorrhagic	Both Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013	120,258 181,840 278 568 239 533 517 1,102	(78,496 to 170,460) (126,699 to 272,666) (176 to 412) (367 to 812) (152 to 337) (351 to 765) (341 to 725) (746 to 1,526)	2425.4 1954.2 14.1 14.2 10.6 11.9 12.1 13	(1525.0 to 3506.0) (1329.8 to 2973.6) (9.0 to 20.6) (9.3 to 20.1) (6.8 to 14.9) (7.7 to 17.1) (8.0 to 16.8) (8.8 to 17.8)

			Female	1990	69	(42 to 102)	2.6	(1.6 to 3.7)
				2013	142	(89 to 207)	2.7	(1.7 to 3.9)
			Both	1990	155	(103 to 220)	3	(2.0 to 4.3)
			Both	2013	329	(212 to 461)	3.1	(2.0 to 4.4)
Antigua and								
Barbuda	Deaths	Ischemic Stroke	Male	1990	18	(15 to 22)	98.3	(79.2 to 117.9)
				2013	20	(14 to 26)	71.4	(52.7 to 92.8)
			Female	1990	27	(20 to 31)	94.7	(70.2 to 108.9)
				2013	23	(17 to 33)	58.1	(42.6 to 82.1)
			Both	1990	46	(37 to 52)	97.5	(79.4 to 110.0)
				2013	43	(35 to 54)	64.2	(51.5 to 80.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	9	(7 to 12)	47.1	(36.3 to 62.9)
				2013	13	(8 to 18)	42.7	(28.6 to 58.2)
			Female	1990	13	(10 to 17)	49.8	(36.3 to 62.1)
				2013	14	(10 to 20)	35.5	(25.2 to 50.5)
			Both	1990	23	(19 to 28)	49.4	(41.2 to 59.7)
				2013	27	(22 to 34)	38.9	(30.8 to 48.7)
	DALYs	Ischemic Stroke	Male	1990	328	(267 to 394)	1598.2	(1300.6 to 1907.0)
				2013	354	(259 to 475)	1115.7	(818.6 to 1473.4)
			Female	1990	370	(269 to 427)	1310.5	(959.5 to 1519.5)
				2013	309	(221 to 438)	767.8	(547.1 to 1088.7)
			Both	1990	698	(575 to 785)	1448	(1193.3 to 1629.5)
				2013	664	(533 to 830)	927.1	(745.5 to 1155.7)
		Hemorrhagic						
		Stroke	Male	1990	219	(173 to 285)	1004.9	(794.8 to 1316.0)
				2013	307	(206 to 430)	844.6	(574.2 to 1163.2)
			Female	1990	249	(187 to 305)	921.5	(693.3 to 1132.8)
				2013	265	(184 to 389)	617.3	(431.1 to 904.1)

					Cou	ints, All ages	Rate ner	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Both	1990	468	(400 to 555)	966.8	(830.1 to 1147.9)
				2013	573	(445 to 721)	722.9	(566.3 to 911.0)
	YLDs	Ischemic Stroke	Male	1990	9	(5 to 13)	45.7	(28.9 to 65.5)
				2013	13	(8 to 19)	42.8	(27.3 to 60.5)
			Female	1990	9	(5 to 12)	32.5	(21.5 to 45.8)
				2013	13	(8 to 19)	31.9	(20.6 to 47.1)
			Both	1990	18	(12 to 25)	38.1	(25.7 to 52.3)
				2013	26	(17 to 37)	36.7	(24.0 to 51.5)
		Hemorrhagic						
		Stroke	Male	1990	3	(1 to 4)	14.4	(8.8 to 21.4)
				2013	5	(3 to 8)	14.7	(8.5 to 23.9)
			Female	1990	2	(1 to 4)	10.6	(6.4 to 15.7)
				2013	4	(2 to 6)	10.2	(6.1 to 16.1)
			Both	1990	5	(3 to 8)	12.3	(8.0 to 17.1)
			Both	2013	9	(6 to 14)	12.2	(7.6 to 18.4)
Argentina	Deaths	Ischemic Stroke	Male	1990	8,606	(6,509 to 10,418)	77.1	(58.5 to 93.3)
				2013	7,588	(5,954 to 10,483)	43.6	(34.0 to 59.9)
			Female	1990	10,114	(6,814 to 11,327)	63.3	(42.6 to 71.0)
				2013	8,355	(6,237 to 13,796)	26.8	(20.1 to 44.5)
			Both	1990	18,720	(14,179 to 21,349)	69.6	(52.7 to 79.0)
				2013	15,944	(12,773 to 23,071)	33.5	(26.8 to 48.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,688	(3,661 to 6,923)	37.6	(29.3 to 55.8)
				2013	7,590	(5,087 to 9,436)	41.4	(27.5 to 51.7)
			Female	1990	4,918	(4,167 to 6,953)	30.8	(26.1 to 43.3)

r								
				2013	6,866	(5,127 to 9,453)	24.8	(18.7 to 33.7)
			Both	1990	9,607	(8,261 to 13,206)	34	(29.2 to 46.8)
				2013	14,457	(11,223 to 16,998)	32	(25.1 to 37.4)
						(114,177 to		
	DALYs	Ischemic Stroke	Male	1990	158,895	194,046)	1252.4	(916.9 to 1524.8)
						(88,809 to		
				2013	112,721	157,395)	617.3	(486.8 to 857.8)
						(88,722 to		
			Female	1990	139,883	156,391)	855.3	(540.0 to 956.7)
						(67,274 to		
				2013	86,705	142,378)	312.7	(242.5 to 512.9)
						(218,233 to		
			Both	1990	298,779	345,881)	1035.2	(762.7 to 1194.7)
						(161,285 to		
				2013	199,427	284,333)	441.1	(356.5 to 627.0)
		Hemorrhagic Stroke	Male	1990	119,786	(94,850 to 173,107)	867.6	(686.7 to 1259.3)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		71		,		(118,086 to		
				2013	165,541	202,771)	861.6	(612.2 to 1058.4)
					,	(92,929 to		,
			Female	1990	105,621	143,302)	659.9	(577.6 to 892.5)
						(91,555 to		
				2013	113,906	148,772)	476.6	(384.8 to 619.2)
						(195,073 to		·
			Both	1990	225,407	302,551)	756.4	(655.2 to 1016.5)
						(225,825 to		•
				2013	279,447	321,708)	650.6	(527.4 to 748.3)
						· · · · · · · · · · · · · · · · · · ·		

	YLDs	Ischemic Stroke	Male	1990	5,119	(3,416 to 6,934)	38.9	(26.2 to 52.6)
				2013	5,627	(3,686 to 7,992)	29.8	(19.6 to 42.2)
			Female	1990	3,749	(2,478 to 5,169)	22.8	(15.1 to 31.4)
				2013	4,607	(2,911 to 6,712)	18.7	(12.0 to 27.4)
			Both	1990	8,869	(5,997 to 11,830)	30	(20.4 to 39.9)
				2013	10,234	(6,732 to 14,436)	23.5	(15.5 to 33.2)
		Hemorrhagic						
		Stroke	Male	1990	1,737	(1,108 to 2,425)	12.8	(8.2 to 17.9)
				2013	2,432	(1,575 to 3,425)	12.6	(8.1 to 17.8)
			Female	1990	1,059	(685 to 1,510)	6.5	(4.2 to 9.3)
				2013	1,512	(972 to 2,188)	6.4	(4.1 to 9.2)
			Both	1990	2,797	(1,836 to 3,782)	9.4	(6.1 to 12.6)
			Both	2013	3,944	(2,651 to 5,465)	9.2	(6.2 to 12.7)
Armenia	Deaths	Ischemic Stroke	Male	1990	737	(539 to 918)	94.7	(70.5 to 117.7)
				2013	1,135	(868 to 1,381)	82.2	(62.6 to 100.0)
			Female	1990	1,090	(901 to 1,368)	85.7	(71.1 to 106.3)
				2013	1,418	(1,072 to 1,696)	68.1	(51.9 to 81.2)
			Both	1990	1,827	(1,536 to 2,162)	89.4	(75.7 to 105.5)
				2013	2,554	(2,104 to 2,902)	74.3	(61.3 to 84.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	573	(431 to 769)	59.2	(42.6 to 82.3)
				2013	616	(463 to 790)	43.8	(32.9 to 56.2)
			Female	1990	636	(459 to 821)	45.9	(32.6 to 60.2)
				2013	663	(521 to 878)	34.1	(26.8 to 45.3)
			Both	1990	1,210	(957 to 1,474)	51.3	(39.7 to 63.8)
				2013	1,280	(1,074 to 1,564)	38.5	(32.4 to 46.8)
	DALYs	Ischemic Stroke	Male	1990	12,369	(8,785 to 15,498)	1327.5	(966.9 to 1644.9)
				2013	15,901	(11,968 to 19,299)	1119.2	(839.4 to 1361.2)
			Female	1990	14,680	(11,870 to 18,901)	1079.6	(879.3 to 1375.3)

				2013	15,716	(12,265 to 18,841)	805.4	(629.3 to 968.5)
			Both	1990	27,049	(22,316 to 32,740)	1183.3	(986.0 to 1416.0)
					31,618	(26,298 to 36,160	945.5	
				2013				
)		(787.1 to 1079.5)
					Cou	ints, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	13,649	(10,640 to 17,826)	1152.8	(872.1 to 1515.1)
				2013	12,254	(9,438 to 15,499)	835.7	(635.3 to 1050.5)
			Female	1990	12,519	(9,437 to 15,916)	827.1	(624.1 to 1054.4)
				2013	10,558	(8,316 to 13,743)	587.5	(465.1 to 768.8)
			Both	1990	26,168	(21,704 to 31,014)	966.4	(791.4 to 1159.8)
				2013	22,813	(19,453 to 27,111)	702.3	(600.6 to 832.6)
	YLDs	Ischemic Stroke	Male	1990	96	(60 to 144)	8	(5.1 to 11.7)
				2013	128	(80 to 184)	8.7	(5.4 to 12.5)
			Female	1990	122	(75 to 180)	8.1	(5.0 to 12.0)
				2013	142	(91 to 203)	8	(5.1 to 11.4)
			Both	1990	219	(144 to 304)	8.1	(5.4 to 11.3)
				2013	270	(179 to 377)	8.3	(5.5 to 11.5)
		Hemorrhagic						
		Stroke	Male	1990	102	(63 to 150)	7.6	(4.8 to 11.2)
				2013	118	(71 to 171)	7.8	(4.7 to 11.3)
			Female	1990	72	(44 to 107)	4.5	(2.8 to 6.8)
				2013	84	(52 to 123)	4.9	(3.1 to 7.1)
			Both	1990	174	(113 to 244)	5.9	(3.9 to 8.3)
			Both	2013	202	(131 to 284)	6.2	(4.0 to 8.8)
Australia	Deaths	Ischemic Stroke	Male	1990	3,265	(2,625 to 4,025)	49.2	(39.6 to 60.6)
				2013	3,889	(2,926 to 4,913)	24.1	(18.2 to 30.4)

		Female	1990	5,007	(3,860 to 5,815)	45.3	(34.8 to 52.6)
			2013	6,022	(4,737 to 8,314)	25.9	(20.5 to 36.0)
		Both	1990	8,273	(7,031 to 9,258)	47.4	(40.2 to 53.1)
			2013	9,911	(8,170 to 12,548)	25.3	(21.1 to 31.9)
					(1,374 to 2,391)		
Deaths	Hemorrhagic Stroke	Male	1990	1,736		22.9	(18.0 to 32.6)
			2013	2,088	(1,279 to 2,722)		(8.2 to 17.4)
		Female					(17.1 to 29.6)
							(9.1 to 16.7)
		Both			` '		(19.3 to 28.9)
			2013			13.2	(9.5 to 15.8)
DALYs	Ischemic Stroke	Male	1990	-	` '	614.5	(507.2 to 745.4)
			2013	-		287.6	(222.6 to 349.7)
		Female	1990		(38,357 to 55,777)	451.3	(355.5 to 514.6)
			2013		(39,309 to 64,337)	234	(194.7 to 316.9)
		Both	1990	94,066	(81,914 to 105,700)	524.5	(456.9 to 587.6)
			2013	91.996	1 ' '	260.3	(222.7 to 316.6)
			2010	-			100,000, Age-standardized
measure	Stroke Type	sex	vear				95% UI
	Stroke	Male	1990	36,484	(29,440 to 48,697)	439.5	(356.6 to 592.0)
			2013	33,491	(22,164 to 41,519)	226.1	(151.1 to 280.5)
		Female	1990	36,716	(29,819 to 48,065)	378.7	(304.2 to 495.2)
			2013	33,136	(24,978 to 42,148)	197.1	(147.3 to 251.0)
		Both	1990	73,200	(64,380 to 90,778)	407.4	(358.5 to 506.5)
		Deaths Stroke DALYs Ischemic Stroke measure Stroke Type Hemorrhagic	Deaths Hemorrhagic Stroke Male Female Both DALYS Ischemic Stroke Male Female Both Both Female Female Female Female Female Female Both	Deaths	Both 1990 8,273 2013 9,911	Both 1990 8,273 (7,031 to 9,258)	Deaths

		T			1			
				2013	66,627	(50,267 to 77,565)	211.1	(160.8 to 245.0)
	YLDs	Ischemic Stroke	Male	1990	4,968	(3,365 to 6,924)	61.2	(41.3 to 85.4)
				2013	9,037	(6,032 to 12,413)	60.6	(40.4 to 83.2)
			Female	1990	3,992	(2,704 to 5,657)	40.3	(27.3 to 57.0)
				2013	6,474	(4,282 to 8,842)	38.9	(25.4 to 53.4)
			Both	1990	8,960	(6,249 to 12,189)	49.7	(34.7 to 67.6)
			2013	15,512	(10,827 to 20,711)	49.2	(34.1 to 66.0)	
		Hemorrhagic						
	Stroke	Stroke	Male	1990	1,346	(872 to 1,956)	16.2	(10.4 to 23.4)
				2013	2,333	(1,521 to 3,227)	16.1	(10.4 to 22.4)
			Female	1990	1,094	(729 to 1,553)	11.4	(7.7 to 16.2)
				2013	1,828	(1,150 to 2,628)	11.7	(7.3 to 16.6)
			Both	1990	2,440	(1,652 to 3,428)	13.6	(9.2 to 19.1)
			Both	2013	4,161	(2,824 to 5,612)	13.8	(9.3 to 18.5)
Austria	Deaths	Ischemic Stroke	Male	1990	2,474	(1,971 to 3,025)	65.2	(51.8 to 79.3)
				2013	1,829	(1,335 to 2,305)	27.3	(19.9 to 34.4)
			Female	1990	4,965	(3,521 to 5,607)	59.1	(41.8 to 66.4)
				2013	3,037	(2,219 to 5,004)	25.3	(18.7 to 41.3)
			Both	1990	7,440	(5,908 to 8,303)	61.7	(49.2 to 68.8)
				2013	4,866	(3,851 to 6,905)	26.5	(21.0 to 37.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,202	(935 to 1,582)	30.8	(23.8 to 40.6)
				2013	1,181	(750 to 1,627)	18.2	(11.6 to 24.7)
			Female	1990	2,095	(1,500 to 2,645)	27.6	(20.2 to 34.8)
				2013	1,575	(1,171 to 2,476)	14.9	(11.4 to 22.4)
			Both	1990	3,297	(2,717 to 4,065)	29	(24.2 to 35.8)
				2013	2,756	(2,192 to 3,683)	16.4	(13.3 to 21.6)
	DALYs	Ischemic Stroke	Male	1990	34,786	(28,410 to 41,379)	891.1	(728.3 to 1058.7)
	2.12.3			2013	25,948	(20,130 to 31,777)	395.3	(306.3 to 482.4)

			Female	1990	52,020	(37,854 to 58,086)	678.8	(494.9 to 759.8)
				2013	28,404	(21,874 to 43,552)	280.4	(218.1 to 426.5)
			Both	1990	86,806	(71,470 to 96,451)	763.7	(630.2 to 850.0)
				2013	54,353	(44,322 to 70,600)	333.4	(271.5 to 428.8)
		Hemorrhagic Stroke	Male	1990	22,350	(17,731 to 28,599)	556.7	(441.5 to 714.8)
			'		Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	18,425	(12,550 to 23,563)	297.3	(202.5 to 379.2)
			Female	1990	28,414	(21,480 to 34,819)	447.6	(346.9 to 541.8)
				2013	17,708	(14,141 to 25,700)	215.7	(175.6 to 304.0)
			Both	1990	50,765	(43,818 to 60,746)	494.5	(432.8 to 592.7)
				2013	36,134	(29,889 to 45,325)	253.3	(208.6 to 310.1)
	YLDs	Ischemic Stroke	Male	1990	3,520	(2,237 to 5,003)	88	(56.0 to 125.0)
				2013	6,282	(4,103 to 8,998)	98.3	(64.5 to 140.4)
			Female	1990	4,112	(2,754 to 5,755)	62.6	(41.9 to 87.4)
				2013	5,132	(3,218 to 7,289)	61.6	(39.5 to 86.5)
			Both	1990	7,633	(5,180 to 10,473)	72.4	(48.9 to 99.6)
				2013	11,415	(7,696 to 15,688)	78	(52.7 to 107.1)
		Hemorrhagic						
		Stroke	Male	1990	731	(415 to 1,179)	18.3	(10.4 to 29.5)
				2013	1,170	(588 to 1,935)	19	(10.0 to 31.2)
			Female	1990	801	(395 to 1,460)	13.2	(7.1 to 22.5)
				2013	988	(544 to 1,591)	13.1	(7.4 to 20.9)
			Both	1990	1,532	(935 to 2,273)	15.1	(9.4 to 22.2)
			Both	2013	2,159	(1,241 to 3,077)	15.7	(9.2 to 22.3)
Azerbaijan	Deaths	Ischemic Stroke	Male	1990	730	(535 to 983)	66.7	(48.1 to 89.7)

				2013	1,417	(1,049 to 1,855)	59.7	(43.7 to 77.9)
			Female	1990	999	(818 to 1,351)	55.5	(45.1 to 74.8)
				2013	1,758	(1,130 to 2,253)	52.3	(33.9 to 67.6)
			Both	1990	1,729	(1,421 to 2,215)	60	(49.0 to 76.4)
				2013	3,176	(2,414 to 3,841)	55.6	(42.3 to 67.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,566	(1,268 to 1,891)	115.5	(92.1 to 141.8)
				2013	2,476	(1,875 to 3,126)	94	(71.4 to 118.9)
			Female	1990	1,955	(1,616 to 2,423)	96.1	(77.2 to 119.6)
				2013	2,948	(2,069 to 3,648)	84.3	(59.5 to 104.7)
			Both	1990	3,522	(3,026 to 4,037)	103.8	(87.8 to 121.0)
				2013	5,425	(4,248 to 6,336)	88.8	(68.8 to 104.1)
	DALYs	Ischemic Stroke	Male	1990	15,378	(11,577 to 20,966)	1020.9	(749.2 to 1375.8)
				2013	24,401	(17,857 to 32,841)	852.2	(628.2 to 1130.7)
			Female	1990	16,071	(12,910 to 21,872)	744	(606.2 to 1011.9)
				2013	23,681	(15,417 to 31,016)	646.2	(421.4 to 845.7)
			Both	1990	31,450	(25,680 to 40,924)	857.6	(702.7 to 1099.7)
				2013	48,082	(37,427 to 59,133)	737.9	(571.5 to 896.1)
		Hemorrhagic						
		Stroke	Male	1990	40,404	(33,244 to 49,118)	2175.9	(1762.3 to 2638.3)
				2013	53,732	(40,646 to 68,699)	1649.8	(1243.9 to 2080.9)
			Female	1990	41,282	(34,133 to 49,739)	1678.3	(1388.3 to 2047.0)
					Counts, All ages		Rate per 100,000, Age-standardized	
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	52,799	(37,685 to 65,640)	1344.6	(946.9 to 1661.6)
			Both	1990	81,686	(71,583 to 92,832)	1885.8	(1626.3 to 2154.7)
				2013	106,531	(85,085 to	1482.7	(1171.3 to 1745.2)

	1	1	 		1	T		
						125,554)		
	YLDs	Ischemic Stroke	Male	1990	151	(97 to 224)	7.9	(5.0 to 11.4)
				2013	296	(188 to 433)	8.4	(5.4 to 12.2)
			Female	1990	188	(119 to 271)	7.6	(4.8 to 10.9)
				2013	314	(199 to 453)	7.6	(4.8 to 10.9)
			Both	1990	340	(230 to 474)	7.7	(5.2 to 10.7)
			2013	610	(400 to 848)	8	(5.3 to 11.0)	
		Hemorrhagic						
		Stroke	Male	1990	182	(116 to 260)	8	(5.2 to 11.4)
				2013	326	(202 to 469)	8.3	(5.2 to 11.7)
			Female	1990	125	(78 to 188)	4.6	(2.9 to 6.9)
				2013	211	(133 to 307)	4.8	(3.1 to 7.0)
			Both	1990	308	(204 to 425)	6.1	(4.1 to 8.4)
			Both	2013	537	(358 to 760)	6.4	(4.3 to 9.0)
Bahrain	Deaths	Ischemic Stroke	Male	1990	36	(26 to 60)	60	(44.0 to 93.9)
				2013	52	(34 to 71)	34.4	(23.9 to 46.1)
			Female	1990	29	(22 to 49)	63.8	(49.0 to 104.4)
				2013	58	(42 to 75)	44.3	(32.0 to 56.1)
			Both	1990	66	(51 to 100)	61.6	(48.3 to 90.8)
				2013	111	(85 to 137)	39.6	(31.0 to 48.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	39	(26 to 53)	40.6	(24.3 to 56.3)
				2013	39	(26 to 56)	13.9	(9.0 to 20.2)
			Female	1990	42	(29 to 51)	58	(39.1 to 70.7)
				2013	41	(30 to 56)	21.1	(15.5 to 28.9)
			Both	1990	81	(61 to 98)	48.2	(33.8 to 58.7)
				2013	80	(64 to 100)	17.4	(13.6 to 22.2)
	DALYs	Ischemic Stroke	Male	1990	742	(511 to 1,347)	849.2	(603.5 to 1368.7)
				2013	1,084	(669 to 1,492)	425.1	(286.5 to 575.6)

			Female	1990	530	(395 to 889)	870.4	(668.9 to 1440.2)
				2013	859	(598 to 1,139)	503.3	(359.2 to 649.7)
			Both	1990	1,272	(953 to 2,039)	860.3	(668.3 to 1291.9)
				2013	1,944	(1,391 to 2,456)	466.7	(356.3 to 576.9)
		Hemorrhagic						
		Stroke	Male	1990	1,362	(970 to 1,819)	853.6	(560.8 to 1144.5)
				2013	1,271	(853 to 1,805)	258	(173.0 to 371.0)
			Female	1990	1,393	(992 to 1,709)	1249.2	(873.6 to 1522.0)
				2013	1,116	(812 to 1,526)	375.5	(272.8 to 512.5)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	2,756	(2,159 to 3,296)	1021.1	(763.7 to 1226.1)
				2013	2,388	(1,888 to 2,981)	310.4	(248.8 to 389.3)
	YLDs	Ischemic Stroke	Male	1990	12	(8 to 18)	9.6	(6.5 to 13.5)
				2013	46	(29 to 69)	9.1	(5.8 to 13.3)
			Female	1990	6	(3 to 9)	6.9	(4.3 to 10.1)
				2013	20	(12 to 29)	6.8	(4.3 to 9.9)
			Both	1990	19	(12 to 26)	8.4	(5.7 to 11.7)
				2013	66	(43 to 96)	8.1	(5.4 to 11.6)
		Hemorrhagic						
		Stroke	Male	1990	2	(1 to 3)	1.2	(0.7 to 1.8)
				2013	7	(4 to 12)	1.2	(0.7 to 1.8)
			Female	1990	1	(0 to 2)	1.4	(0.8 to 2.2)
				2013	4	(2 to 6)	1.2	(0.7 to 2.0)
			Both	1990	3	(2 to 5)	1.3	(0.8 to 1.9)
			Both	2013	11	(7 to 18)	1.2	(0.8 to 1.9)
Bangladesh	Deaths	Ischemic Stroke	Male	1990	13,667	(6,053 to 20,412)	68.4	(31.1 to 100.3)

			, ,		1	1		I
				2013	41,762	(22,786 to 57,364)	116.2	(65.7 to 156.9)
			Female	1990	12,062	(6,932 to 17,527)	72.8	(41.2 to 104.0)
				2013	42,286	(14,897 to 62,317)	127.6	(44.8 to 185.0)
			Both	1990	25,730	(14,722 to 33,330)	70.4	(41.0 to 90.5)
						(43,323 to		
				2013	84,049	109,294)	121.7	(62.9 to 156.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	15,730	(8,522 to 22,634)	66.3	(33.6 to 99.8)
				2013	44,782	(28,712 to 75,765)	108.5	(67.6 to 185.5)
			Female	1990	16,301	(11,493 to 25,268)	81.7	(57.1 to 129.5)
				2013	49,521	(26,929 to 70,387)	130.6	(72.7 to 187.4)
			Both	1990	32,031	(23,745 to 43,431)	73.4	(52.7 to 103.1)
						(62,327 to		
				2013	94,304	126,815)	119	(78.9 to 162.5)
						(107,357 to		
	DALYs	Ischemic Stroke	Male	1990	241,301	373,524)	1012	(456.8 to 1531.7)
						(333,132 to		(
				2013	642,761	928,948)	1563.4	(845.2 to 2187.8)
			F	1000	200 200	(123,990 to	1001 5	(C20 0 +- 4552 0)
			Female	1990	209,290	316,273)	1061.5	(628.0 to 1553.0)
				2013	624,572	(220,804 to 963,289)	1675.8	(596.2 to 2518.8)
				2013	024,372	(264,527 to	1075.8	(390.2 to 2318.8)
			Both	1990	450,591	596,285)	1034.3	(605.1 to 1347.7)
			500.1	1000	.55,551	(658,458 to		(303.1 to 13)
				2013	1,267,333	1,688,535)	1617.2	(841.2 to 2129.5)
								,
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		, i						

		Hemorrhagic				(238,275 to		
		Stroke	Male	1990	387,461	533,037)	1335.5	(771.1 to 1887.9)
					,	(664,068 to		,
				2013	993,527	1,585,407)	1979	(1301.9 to 3255.3)
						(272,413 to		
			Female	1990	381,644	583,282)	1598.6	(1137.7 to 2447.9)
						(547,777 to		
				2013	1,042,594	1,479,916)	2337	(1249.6 to 3302.4)
						(596,039 to		(
			Both	1990	769,105	991,699)	1456.3	(1100.2 to 1904.5)
				2012	2.026.422	(1,345,715 to	2140 1	/1/417 2 += 2050 2\
	VI Da	laskania Charles	0.4-1-	2013	2,036,122	2,679,184)	2148.1	(1417.3 to 2858.3)
	YLDs	Ischemic Stroke	Male	1990	8,097	(5,146 to 11,417)	28.5	(17.8 to 40.3)
				2013	15,160	(9,659 to 21,772)	29.6	(19.0 to 42.3)
			Female	1990	5,964	(3,993 to 8,498)	24.1	(16.2 to 34.0)
				2013	12,378	(8,037 to 17,566)	25.9	(17.0 to 36.8)
			Both	1990	14,061	(9,589 to 18,875)	26.5	(17.9 to 35.8)
				2013	27,538	(18,423 to 37,706)	27.8	(18.8 to 38.1)
		Hemorrhagic						
		Stroke	Male	1990	4,422	(2,799 to 6,243)	14.4	(9.1 to 20.3)
				2013	7,880	(4,856 to 11,014)	14.3	(9.0 to 20.2)
			Female	1990	2,922	(1,904 to 4,106)	10.7	(6.9 to 15.0)
				2013	5,692	(3,650 to 8,161)	10.8	(7.0 to 15.5)
			Both	1990	7,344	(4,780 to 9,959)	12.7	(8.3 to 17.2)
			Both	2013	13,573	(8,850 to 18,496)	12.6	(8.3 to 17.2)
Barbados	Deaths	Ischemic Stroke	Male	1990	85	(71 to 102)	88.3	(73.9 to 106.3)
				2013	77	(57 to 100)	61.2	(45.2 to 79.1)
			Female	1990	147	(112 to 173)	89.3	(68.2 to 104.7)
				2013	112	(82 to 166)	58.3	(42.9 to 86.1)
_			Both	1990	232	(195 to 262)	89.8	(75.3 to 101.3)

				2013	189	(153 to 246)	60.1	(48.3 to 78.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	35	(28 to 47)	36.9	(29.8 to 49.6)
				2013	46	(27 to 63)	33.5	(19.2 to 45.7)
			Female	1990	58	(44 to 78)	38.1	(29.9 to 50.8)
				2013	58	(40 to 82)	31.1	(21.4 to 43.2)
			Both	1990	94	(79 to 119)	38	(32.3 to 47.7)
				2013	105	(78 to 134)	32.5	(24.1 to 41.7)
	DALYs	Ischemic Stroke	Male	1990	1,321	(1,119 to 1,573)	1347.9	(1141.8 to 1605.7)
				2013	1,263	(936 to 1,604)	893	(667.3 to 1142.4)
			Female	1990	1,737	(1,334 to 2,029)	1124	(864.8 to 1308.9)
				2013	1,368	(1,017 to 1,949)	738	(548.5 to 1048.8)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Cou Mean	nts, All ages 95% UI	Rate per Mean	100,000, Age-standardized 95% UI
Country	measure	Stroke Type	sex Both	year 1990				
Country	measure	Stroke Type			Mean	95% UI	Mean	95% UI
Country	measure	Stroke Type Hemorrhagic		1990	Mean 3,059	95% UI (2,616 to 3,442)	Mean 1225.7	95% UI (1051.9 to 1379.8)
Country	measure			1990	Mean 3,059	95% UI (2,616 to 3,442)	Mean 1225.7	95% UI (1051.9 to 1379.8)
Country	measure	Hemorrhagic	Both	1990 2013	Mean 3,059 2,632	95% UI (2,616 to 3,442) (2,157 to 3,297)	Mean 1225.7 812.9	95% UI (1051.9 to 1379.8) (666.3 to 1017.3)
Country	measure	Hemorrhagic	Both	1990 2013 1990	Mean 3,059 2,632 801	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000)	Mean 1225.7 812.9 794	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5)
Country	measure	Hemorrhagic	Both Male	1990 2013 1990 2013	Mean 3,059 2,632 801 1,000	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337)	Mean 1225.7 812.9 794 648.1	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1)
Country	measure	Hemorrhagic	Both Male	1990 2013 1990 2013 1990	Mean 3,059 2,632 801 1,000 963	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337) (771 to 1,217)	Mean 1225.7 812.9 794 648.1 699.5	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1) (558.7 to 873.0)
Country	measure	Hemorrhagic	Male Female	1990 2013 1990 2013 1990 2013	Mean 3,059 2,632 801 1,000 963 955	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337) (771 to 1,217) (669 to 1,299)	Mean 1225.7 812.9 794 648.1 699.5 530	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1) (558.7 to 873.0) (375.7 to 722.1)
Country	measure	Hemorrhagic	Male Female	1990 2013 1990 2013 1990 2013 1990	Mean 3,059 2,632 801 1,000 963 955 1,765	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337) (771 to 1,217) (669 to 1,299) (1,556 to 2,102)	Mean 1225.7 812.9 794 648.1 699.5 530 745.1	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1) (558.7 to 873.0) (375.7 to 722.1) (656.6 to 885.2)
Country		Hemorrhagic Stroke	Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013	Mean 3,059 2,632 801 1,000 963 955 1,765 1,955	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337) (771 to 1,217) (669 to 1,299) (1,556 to 2,102) (1,496 to 2,441)	Mean 1225.7 812.9 794 648.1 699.5 530 745.1 588.6	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1) (558.7 to 873.0) (375.7 to 722.1) (656.6 to 885.2) (453.2 to 736.0)
Country		Hemorrhagic Stroke	Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013 1990	Mean 3,059 2,632 801 1,000 963 955 1,765 1,955 44	95% UI (2,616 to 3,442) (2,157 to 3,297) (650 to 1,000) (634 to 1,337) (771 to 1,217) (669 to 1,299) (1,556 to 2,102) (1,496 to 2,441) (29 to 61)	Mean 1225.7 812.9 794 648.1 699.5 530 745.1 588.6 45.1	95% UI (1051.9 to 1379.8) (666.3 to 1017.3) (642.3 to 996.5) (412.2 to 864.1) (558.7 to 873.0) (375.7 to 722.1) (656.6 to 885.2) (453.2 to 736.0) (30.3 to 63.0)

					,			
			Both	1990	92	(63 to 124)	37.6	(25.4 to 50.4)
				2013	122	(84 to 164)	37.5	(25.7 to 50.3)
		Hemorrhagic						
		Stroke	Male	1990	14	(8 to 20)	14.3	(8.8 to 20.6)
			2013	20	(12 to 32)	13.6	(8.3 to 21.0)	
		Female	1990	14	(8 to 20)	9.9	(5.8 to 14.6)	
				2013	19	(12 to 29)	10.9	(6.7 to 16.1)
			Both	1990	28	(18 to 39)	11.7	(7.7 to 16.3)
			Both	2013	40	(26 to 58)	12.1	(7.9 to 17.5)
Belarus	Deaths	Ischemic Stroke	Male	1990	4,160	(3,218 to 5,187)	120.1	(95.1 to 147.8)
				2013	6,008	(4,857 to 7,632)	131.1	(106.5 to 163.7)
			Female	1990	7,908	(6,464 to 9,686)	102.2	(83.3 to 125.3)
				2013	10,632	(8,378 to 12,627)	101.4	(81.3 to 120.9)
			Both	1990	12,069	(10,546 to 14,231)	109	(95.6 to 128.2)
				2013	16,640	(14,442 to 19,185)	114	(98.8 to 131.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,186	(1,341 to 3,083)	52.4	(31.2 to 75.3)
				2013	2,567	(1,788 to 3,445)	50.4	(34.8 to 67.2)
			Female	1990	2,883	(1,639 to 4,001)	39.1	(22.6 to 53.8)
				2013	3,066	(2,209 to 4,057)	34.4	(25.6 to 45.4)
			Both	1990	5,070	(3,198 to 6,388)	44.5	(28.0 to 55.8)
				2013	5,634	(4,502 to 7,000)	41.4	(33.1 to 51.5)
	DALYs	Ischemic Stroke	Male	1990	75,232	(55,640 to 96,493)	1900.2	(1449.5 to 2387.8)
						(87,588 to		
				2013	109,927	143,250)	2217.5	(1778.3 to 2881.6)
						(82,955 to		
			Female	1990	104,828	131,363)	1396.6	(1096.6 to 1752.9)
				2010	100 -00	(104,123 to		(4444.0) (4655.5)
				2013	126,723	153,003)	1341.4	(1111.2 to 1628.9)

					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(153,863 to		
			Both	1990	180,061	219,533)	1590.5	(1369.2 to 1928.3)
						(205,476 to		
				2013	236,651	284,826)	1698.2	(1474.3 to 2049.3)
		Hemorrhagic				,		,
		Stroke	Male	1990	56,576	(36,989 to 77,734)	1219.1	(785.3 to 1692.2)
				2013	67,334	(47,607 to 89,909)	1246.6	(871.1 to 1661.0)
			Female	1990	57,077	(34,395 to 75,667)	821.1	(509.3 to 1073.2)
				2013	57,079	(43,399 to 74,385)	727.4	(567.8 to 936.8)
						(76,868 to		
			Both	1990	113,653	140,641)	992.7	(680.5 to 1225.2)
						(99,448 to		
				2013	124,413	153,556)	957.4	(764.8 to 1182.8)
	YLDs	Ischemic Stroke	Male	1990	1,749	(1,101 to 2,539)	38.7	(24.8 to 55.3)
				2013	2,162	(1,421 to 3,062)	41.9	(27.7 to 58.4)
			Female	1990	1,729	(1,121 to 2,486)	25.2	(16.4 to 36.2)
				2013	1,903	(1,188 to 2,709)	24.3	(15.5 to 34.3)
			Both	1990	3,478	(2,331 to 4,795)	30.7	(20.6 to 42.4)
				2013	4,065	(2,698 to 5,624)	31.5	(20.9 to 43.5)
		Hemorrhagic						,
		Stroke	Male	1990	60	(36 to 90)	1.3	(0.8 to 1.9)
				2013	66	(42 to 97)	1.3	(0.8 to 1.9)
			Female	1990	60	(36 to 94)	0.9	(0.6 to 1.5)
				2013	65	(38 to 102)	0.9	(0.5 to 1.4)
			Both	1990	121	(77 to 172)	1.1	(0.7 to 1.6)
			Both	2013	131	(83 to 192)	1.1	(0.7 to 1.6)
Belgium	Deaths	Ischemic Stroke	Male	1990	2,650	(2,119 to 3,256)	50.4	(39.8 to 62.0)

				2013	2,280	(1,721 to 3,018)	24.9	(18.9 to 32.9)
			Female	1990	4,681	(3,314 to 5,450)	45.8	(32.7 to 53.1)
				2013	3,465	(2,589 to 5,307)	22	(16.7 to 33.7)
			Both	1990	7,332	(5,877 to 8,261)	48.1	(38.7 to 54.1)
				2013	5,745	(4,657 to 7,971)	23.4	(19.2 to 32.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,676	(1,308 to 2,213)	30.6	(23.9 to 40.6)
				2013	2,041	(1,322 to 2,606)	23.2	(15.2 to 29.5)
			Female	1990	2,533	(1,895 to 3,265)	27.5	(20.9 to 35.5)
				2013	2,611	(1,866 to 3,622)	18.9	(13.9 to 25.8)
			Both	1990	4,210	(3,502 to 5,136)	28.9	(24.1 to 35.3)
				2013	4,652	(3,580 to 5,637)	20.8	(16.3 to 25.1)
	DALYs	Ischemic Stroke	Male	1990	38,829	(31,988 to 46,398)	701.6	(578.8 to 836.9)
				2013	30,971	(25,161 to 37,977)	359	(290.2 to 440.8)
			Female	1990	49,158	(36,493 to 55,476)	534.6	(401.6 to 604.6)
				2013	33,084	(26,267 to 48,174)	256.5	(205.2 to 367.3)
			Both	1990	87,987	(73,840 to 98,699)	605.8	(509.4 to 677.9)
					Cou	ints, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	64,055	(53,909 to 82,201)	303	(255.7 to 386.9)
		Hemorrhagic						
		Stroke	Male	1990	30,711	(24,849 to 39,006)	543.8	(439.9 to 691.1)
				2013	29,629	(20,408 to 36,265)	365.7	(256.4 to 447.4)
			Female	1990	34,882	(27,315 to 43,873)	453.3	(356.6 to 562.0)
				2013	28,877	(22,422 to 38,725)	277.2	(222.5 to 372.6)
			Both	1990	65,593	(56,879 to 78,646)	492.7	(432.1 to 587.0)
				2013	58,507	(47,770 to 68,927)	317.2	(260.6 to 371.4)
	YLDs	Ischemic Stroke	Male	1990	4,859	(3,271 to 6,771)	84.9	(57.3 to 118.3)
				2013	6,957	(4,631 to 9,795)	85	(56.3 to 119.8)

			Female	1990	4,769	(3,192 to 6,808)	59.6	(40.0 to 83.4)
				2013	5,937	(3,934 to 8,410)	56.4	(37.7 to 77.2)
			Both	1990	9,628	(6,644 to 13,116)	70.1	(48.5 to 95.6)
				2013	12,894	(8,772 to 17,439)	69.5	(47.2 to 94.0)
	Hemorrhagic Stroke	Hemorrhagic						
		Stroke	Male	1990	1,005	(564 to 1,532)	17.8	(10.1 to 26.9)
				2013	1,365	(773 to 2,482)	17.5	(10.1 to 31.1)
			Female	1990	941	(514 to 1,553)	12.7	(7.2 to 21.1)
				2013	1,167	(673 to 1,867)	12.5	(7.5 to 19.7)
			Both	1990	1,946	(1,205 to 2,942)	14.8	(9.2 to 22.2)
			Both	2013	2,533	(1,622 to 3,743)	14.8	(9.7 to 21.5)
Belize	Deaths	Ischemic Stroke	Male	1990	27	(21 to 33)	74.4	(58.9 to 90.3)
				2013	51	(38 to 65)	79.1	(60.3 to 100.6)
			Female	1990	28	(23 to 34)	68.5	(54.7 to 82.7)
				2013	60	(45 to 78)	78.7	(59.5 to 102.6)
			Both	1990	56	(48 to 64)	71.4	(61.7 to 82.3)
				2013				
					111	(93 to 134)	79.2	(66.7 to 94.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	15	(12 to 19)	37.2	(28.2 to 47.3)
				2013	31	(20 to 41)	41.5	(26.6 to 54.1)
			Female	1990	17	(13 to 21)	39.1	(30.1 to 48.6)
				2013	39	(28 to 53)	46.9	(33.1 to 62.2)
			Both	1990	33	(27 to 39)	38.3	(31.2 to 45.3)
				2013	71	(54 to 87)	44.5	(33.6 to 54.0)
	DALYs	Ischemic Stroke	Male	1990	512	(409 to 618)	1187.8	(947.4 to 1431.6)
				2013	965	(725 to 1,244)	1292.8	(984.6 to 1633.3)
			Female	1990	453	(355 to 546)	1001.2	(789.1 to 1219.1)
				2013	952	(711 to 1,302)	1157.9	(877.7 to 1563.0)

			1 1				1	
			Both	1990	965	(819 to 1,108)	1093.4	(928.1 to 1258.5)
				2013	1,917	(1,578 to 2,380)	1224.8	(1015.3 to 1497.0)
		Hemorrhagic						
		Stroke	Male	1990	445	(352 to 543)	807.2	(621.5 to 1002.5)
					Cou	nts, All ages		100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	838	(568 to 1,118)	880.3	(576.2 to 1150.7)
			Female	1990	443	(355 to 540)	803.5	(633.5 to 971.6)
				2013	928	(646 to 1,263)	933.2	(657.2 to 1269.5)
			Both	1990	889	(749 to 1,025)	805.8	(672.8 to 934.6)
				2013	1,766	(1,385 to 2,207)	909.2	(708.1 to 1127.8)
	YLDs	Ischemic Stroke	Male	1990	19	(12 to 27)	48.2	(30.9 to 69.3)
				2013	33	(21 to 46)	44.7	(28.3 to 62.2)
			Female	1990	13	(8 to 19)	31.2	(19.5 to 46.1)
				2013	26	(17 to 37)	31.8	(20.4 to 45.4)
			Both	1990	32	(21 to 46)	39.3	(26.5 to 56.2)
				2013	59	(39 to 81)	37.9	(25.0 to 52.4)
		Hemorrhagic						
		Stroke	Male	1990	6	(4 to 9)	15.2	(9.5 to 21.9)
				2013	12	(8 to 19)	14.3	(8.9 to 22.5)
			Female	1990	4	(2 to 7)	10	(6.1 to 15.5)
				2013	10	(6 to 15)	10.3	(6.0 to 15.4)
			Both	1990	11	(7 to 16)	12.6	(8.2 to 17.9)
			Both	2013	22	(14 to 33)	12.2	(8.0 to 17.7)
Benin	Deaths	Ischemic Stroke	Male	1990	419	(224 to 547)	65.6	(37.9 to 85.0)
				2013	1,204	(386 to 1,778)	100.4	(35.0 to 144.8)
			Female	1990	565	(275 to 765)	65.7	(33.5 to 91.4)
				2013	1,320	(412 to 1,917)	90.5	(29.6 to 128.5)
			Both	1990	985	(610 to 1,255)	65.9	(41.3 to 85.6)

			2013	2,525	(1,202 to 3,444)	95.2	(45.7 to 127.1)
	Hemorrhagic						
Deaths	Stroke	Male	1990	532	(409 to 722)	62	(45.8 to 87.6)
			2013	994	(570 to 1,829)	60.2	(30.8 to 125.0)
		Female	1990	627	(420 to 919)	59.8	(38.1 to 91.5)
			2013	938	(557 to 1,851)	51.1	(28.1 to 109.2)
		Both	1990	1,160	(897 to 1,551)	61	(44.0 to 85.3)
			2013	1,932	(1,211 to 3,386)	55.4	(32.1 to 107.3)
DALYs	Ischemic Stroke	Male	1990	9,340	(4,682 to 12,389)	1065.4	(570.4 to 1396.5)
			2013	26,165	(7,707 to 39,746)	1599.3	(523.2 to 2343.0)
		Female	1990	10,715	(4,911 to 14,738)	955.1	(456.3 to 1283.3)
			2013	23,502	(6,939 to 35,643)	1273.9	(401.9 to 1860.9)
		Both	1990	20,056	(12,312 to 25,502)	1004.7	(620.4 to 1279.4)
			2013	49,668	(21,384 to 68,636)	1426.3	(683.1 to 1940.5)
	Hemorrhagic						
	Stroke	Male	1990	18,105	(13,604 to 23,959)	1342.3	(1025.4 to 1827.4)
			2013	30,119	(18,819 to 49,888)	1268	(718.1 to 2357.2)
		Female	1990	17,419	(11,864 to 23,901)	1147.5	(772.0 to 1665.4)
			2013	23,668	(15,068 to 41,566)	966	(583.9 to 1860.3)
		Both	1990	35,525	(27,963 to 44,702)	1238	(963.9 to 1648.6)
				Cou	nts, All ages	Rate per	100,000, Age-standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			2013	53,788	(36,857 to 83,507)	1110.2	(701.8 to 1927.0)
YLDs	Ischemic Stroke	Male	1990	155	(96 to 225)	15.7	(9.9 to 23.0)
			2013	337	(211 to 482)	15.5	(9.9 to 22.4)
		Female	1990	143	(90 to 207)	11.1	(7.0 to 16.3)
			2013	294	(182 to 428)	12	(7.5 to 17.1)
		Both	1990	298	(198 to 414)	13	(8.8 to 17.9)
			2013	632	(414 to 889)	13.6	(8.9 to 18.8)
	DALYs	Deaths Stroke DALYS Ischemic Stroke Hemorrhagic Stroke measure Stroke Type	Deaths Stroke Male Female Both DALYS Ischemic Stroke Male Female Both Hemorrhagic Stroke Male Female Both Female Female Female Female Female Female Female	Hemorrhagic Stroke	Deaths	Hemorrhagic Stroke	Deaths Hemorrhagic Stroke Male 1990 532 (409 to 722) 62 Both 2013 994 (570 to 1,829) 60.2 Both 1990 627 (420 to 919) 59.8 Both 1990 1,160 (897 to 1,551) 61 Both 1990 1,160 (897 to 1,551) 61 DALYS Ischemic Stroke Male 1990 9,340 (4,682 to 12,389) 1065.4 DALYS Ischemic Stroke Male 1990 10,715 (4,911 to 14,738) 955.1 Both 1990 10,715 (4,911 to 14,738) 955.1 Both 1990 20,056 (12,312 to 25,502) 1004.7 Both 1990 18,105 (13,604 to 23,959) 1342.3 Hemorrhagic Male 1990 18,105 (13,604 to 23,959) 1342.3 Stroke Male 1990 17,419 (11,864 to 23,901) 1147.5 Both 1990 17,419 (11,864 to 23,901)

			T I					
		Hemorrhagic Stroke	Male	1990	49	(30 to 71)	4.2	(2.5 to 6.2)
		00.000		2013	107	(67 to 160)	4	(2.5 to 6.0)
			Female	1990	40	(24 to 59)	2.8	(1.7 to 4.1)
				2013	78	(48 to 115)	2.7	(1.7 to 4.0)
			Both	1990	90	(57 to 125)	3.4	(2.2 to 4.7)
			Both	2013	186	(120 to 266)	3.3	(2.2 to 4.7)
Bhutan	Deaths	Ischemic Stroke	Male	1990	48	(23 to 76)	64	(32.7 to 97.0)
				2013	146	(88 to 209)	77.8	(47.4 to 109.2)
			Female	1990	46	(27 to 73)	70	(42.6 to 105.3)
				2013	129	(61 to 196)	84.8	(41.2 to 127.4)
			Both	1990	95	(56 to 142)	67	(41.3 to 96.3)
				2013	275	(160 to 392)	81	(48.1 to 113.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	64	(35 to 107)	60.5	(33.1 to 98.2)
				2013	144	(83 to 232)	65.6	(39.2 to 103.8)
			Female	1990	62	(37 to 101)	67.4	(40.6 to 107.5)
				2013	122	(58 to 206)	71.5	(33.6 to 117.6)
			Both	1990	127	(79 to 198)	63.8	(40.5 to 96.2)
				2013	267	(157 to 419)	68.2	(41.3 to 104.9)
	DALYs	Ischemic Stroke	Male	1990	1,014	(506 to 1,670)	959.9	(482.3 to 1515.5)
				2013	2,335	(1,341 to 3,550)	1077.5	(630.3 to 1587.7)
			Female	1990	916	(513 to 1,536)	1010.1	(607.2 to 1583.2)
				2013	1,861	(839 to 2,955)	1106.2	(508.7 to 1735.7)
			Both	1990	1,931	(1,111 to 2,986)	984.8	(591.1 to 1465.6)
				2013	4,196	(2,394 to 6,253)	1090.8	(633.4 to 1597.4)
		Hemorrhagic						
		Stroke	Male	1990	1,936	(1,123 to 3,127)	1265.3	(713.0 to 2095.0)
				2013	3,403	(1,889 to 5,706)	1263.2	(717.2 to 2066.3)

			Female	1990	1,763	(1,051 to 2,819)	1367.7	(818.4 to 2216.2)
				2013	2,669	(1,260 to 4,635)	1316.3	(619.2 to 2283.3)
			Both	1990	3,699	(2,348 to 5,801)	1313.6	(817.6 to 2041.8)
				2013	6,073	(3,357 to 10,092)	1285.6	(728.9 to 2103.1)
	YLDs	Ischemic Stroke	Male	1990	38	(24 to 55)	27.5	(17.1 to 39.8)
				2013	76	(50 to 111)	29.2	(18.8 to 42.0)
			Female	1990	30	(19 to 44)	26	(16.6 to 36.7)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	51	(33 to 74)	25.2	(16.4 to 35.8)
			Both	1990	69	(46 to 96)	26.8	(17.9 to 37.2)
				2013	128	(86 to 178)	27.4	(18.4 to 38.1)
		Hemorrhagic						
		Stroke	Male	1990	22	(13 to 32)	14.2	(8.7 to 20.9)
				2013	41	(26 to 60)	14.4	(9.2 to 20.8)
			Female	1990	14	(9 to 21)	11.3	(7.4 to 15.8)
				2013	24	(15 to 34)	10.8	(7.0 to 15.3)
			Both	1990	36	(24 to 51)	12.9	(8.4 to 17.8)
			Both	2013	65	(43 to 90)	12.8	(8.5 to 17.8)
Bolivia	Deaths	Ischemic Stroke	Male	1990	472	(368 to 600)	43.7	(33.2 to 55.1)
				2013	784	(564 to 1,044)	37.4	(27.2 to 49.8)
			Female	1990	586	(436 to 744)	45.4	(33.5 to 57.4)
				2013	1,088	(762 to 1,539)	40.5	(28.2 to 57.0)
			Both	1990	1,059	(885 to 1,241)	44.7	(36.9 to 52.7)
				2013	1,872	(1,465 to 2,386)	39.3	(30.9 to 49.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	633	(480 to 775)	39.2	(28.9 to 49.2)
				2013	832	(567 to 1,150)	31	(21.1 to 42.4)
			Female	1990	774	(595 to 994)	42.6	(32.3 to 55.3)

				2013	1,023	(661 to 1,413)	32.2	(20.6 to 44.8)
			Both	1990	1,408	(1,163 to 1,677)	41	(33.6 to 48.9)
				2013	1,856	(1,403 to 2,356)	31.7	(23.6 to 40.1)
	DALYs	Ischemic Stroke	Male	1990	11,284	(8,690 to 14,573)	720.9	(564.5 to 909.2)
				2013	14,829	(10,275 to 20,312)	548.2	(388.9 to 739.5)
			Female	1990	12,420	(8,864 to 16,720)	714.2	(530.0 to 913.2)
				2013	17,443	(11,702 to 25,049)	556.8	(383.1 to 797.4)
			Both	1990	23,705	(19,097 to 28,340)	717.9	(596.4 to 842.1)
				2013	32,273	(24,664 to 41,960)	554.4	(428.8 to 707.2)
		Hemorrhagic						
		Stroke	Male	1990	24,777	(18,703 to 31,164)	992.7	(755.0 to 1215.8)
				2013	24,210	(16,473 to 33,406)	676.2	(454.6 to 936.3)
			Female	1990	28,004	(20,326 to 36,465)	1083.2	(839.5 to 1397.1)
				2013	26,632	(17,683 to 36,951)	690.6	(450.1 to 961.4)
			Both	1990	52,782	(42,932 to 63,703)	1040	(859.1 to 1231.6)
				2013	50,842	(38,786 to 65,147)	684.4	(517.1 to 874.6)
	YLDs	Ischemic Stroke	Male	1990	276	(178 to 390)	18.2	(12.1 to 25.6)
				2013	468	(310 to 653)	16	(10.6 to 22.3)
			Female	1990	149	(96 to 219)	8.4	(5.5 to 12.2)
				2013	271	(174 to 393)	7.9	(5.0 to 11.5)
			Both	1990	425	(281 to 594)	12.9	(8.7 to 17.8)
				2013	739	(498 to 1,019)	11.6	(7.9 to 15.9)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	112	(69 to 167)	6.2	(3.8 to 9.4)
				2013	209	(131 to 311)	6	(3.8 to 8.8)
			Female	1990	71	(45 to 106)	3.4	(2.1 to 5.1)

				2013	136	(87 to 202)	3.5	(2.2 to 5.3)
			Both	1990	183	(120 to 265)	4.7	(3.1 to 6.8)
			Both	2013	345	(228 to 493)	4.7	(3.1 to 6.6)
Bosnia and								
Herzegovina	Deaths	Ischemic Stroke	Male	1990	1,129	(940 to 1,430)	105.7	(87.7 to 133.1)
				2013	2,406	(1,616 to 2,976)	109.3	(74.3 to 135.0)
			Female	1990	1,355	(1,147 to 1,668)	83.2	(70.5 to 102.2)
				2013	2,751	(1,545 to 3,483)	87.6	(49.7 to 111.1)
			Both	1990	2,484	(2,225 to 2,988)	91.9	(82.1 to 109.8)
				2013	5,157	(3,522 to 6,256)	96.9	(66.8 to 117.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	906	(728 to 1,114)	67.6	(53.3 to 83.9)
				2013	676	(340 to 1,257)	29.7	(14.8 to 56.0)
			Female	1990	1,086	(822 to 1,250)	59.2	(44.4 to 69.3)
				2013	724	(385 to 1,598)	23.7	(12.7 to 51.1)
			Both	1990	1,993	(1,699 to 2,261)	62.8	(52.8 to 72.1)
				2013	1,401	(762 to 2,654)	26.4	(14.5 to 49.7)
	DALYs	Ischemic Stroke	Male	1990	17,901	(14,797 to 22,792)	1404.6	(1168.9 to 1776.4)
				2013	34,001	(21,856 to 42,683)	1413.7	(916.7 to 1764.6)
			Female	1990	18,084	(15,508 to 22,484)	993	(847.4 to 1231.2)
				2013	31,952	(16,704 to 40,844)	991	(521.8 to 1262.2)
			Both	1990	35,986	(32,007 to 43,584)	1156.8	(1037.4 to 1394.3)
				2013	65,953	(42,838 to 81,208)	1176.1	(766.5 to 1447.3)
		Hemorrhagic						
		Stroke	Male	1990	22,019	(18,179 to 26,865)	1313.2	(1072.6 to 1597.6)
				2013	13,314	(7,579 to 23,159)	566.6	(329.1 to 973.8)
			Female	1990	21,634	(17,119 to 24,506)	1045.5	(816.9 to 1185.2)
				2013	11,798	(7,054 to 23,644)	403.8	(250.2 to 780.1)
			Both	1990	43,654	(38,350 to 49,405)	1165.4	(1020.9 to 1326.9)

				2013	25,112	(15,315 to 43,629)	479.9	(299.7 to 813.4)
	YLDs	Ischemic Stroke	Male	1990	369	(231 to 544)	20.7	(13.2 to 30.1)
				2013	617	(405 to 887)	25.4	(16.7 to 36.5)
			Female	1990	302	(200 to 431)	14.1	(9.4 to 19.9)
				2013	417	(263 to 601)	14.4	(9.3 to 20.6)
			Both	1990	671	(452 to 925)	17	(11.7 to 23.6)
				2013	1,034	(701 to 1,436)	19.4	(13.2 to 27.0)
		Hemorrhagic						
		Stroke	Male	1990	92	(53 to 142)	4.7	(2.6 to 7.0)
				2013	124	(72 to 195)	5.4	(3.2 to 8.3)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	77	(46 to 117)	3.4	(2.1 to 5.2)
				2013	96	(57 to 155)	3.6	(2.2 to 5.8)
			Both	1990	169	(108 to 238)	4	(2.5 to 5.6)
			Both	2013	221	(140 to 332)	4.4	(2.8 to 6.6)
Botswana	Deaths	Ischemic Stroke	Male	1990	57	(28 to 92)	45.2	(23.2 to 68.9)
				2013	130	(77 to 214)	48.5	(29.4 to 76.7)
			Female	1990	93	(49 to 156)	51	(28.4 to 79.9)
				2013	109	(51 to 262)	28.4	(14.0 to 63.1)
			Both	1990	151	(89 to 235)	49.1	(30.3 to 72.3)
				2013	240	(156 to 404)	36.3	(24.0 to 60.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	88	(51 to 139)	54.1	(32.3 to 82.5)
				2013	134	(83 to 207)	38.1	(23.7 to 59.3)
			Female	1990	121	(73 to 185)	56.9	(34.7 to 84.0)
				2013	93	(49 to 200)	19.8	(10.3 to 43.1)
			Both	1990	210	(138 to 287)	56.1	(37.5 to 74.4)
				2013	227	(155 to 348)	27.2	(18.1 to 42.8)

					•			
	DALYs	Ischemic Stroke	Male	1990	1,186	(542 to 1,986)	678.4	(338.6 to 1069.2)
				2013	2,475	(1,420 to 4,106)	693.7	(412.9 to 1153.9)
			Female	1990	1,560	(767 to 2,936)	677.3	(357.1 to 1168.8)
				2013	1,407	(633 to 3,606)	285.4	(127.7 to 732.7)
			Both	1990	2,746	(1,523 to 4,613)	681.5	(404.9 to 1062.5)
				2013	3,883	(2,489 to 6,453)	450.4	(287.1 to 762.4)
		Hemorrhagic						
		Stroke	Male	1990	2,390	(1,384 to 3,793)	980.9	(575.3 to 1551.3)
				2013	3,741	(2,341 to 5,614)	722.5	(446.2 to 1107.6)
			Female	1990	2,650	(1,570 to 4,126)	922.6	(553.1 to 1421.4)
				2013	2,309	(1,296 to 4,430)	327.9	(176.9 to 698.3)
			Both	1990	5,041	(3,319 to 7,030)	952.5	(625.7 to 1303.5)
				2013	6,050	(4,244 to 8,740)	495.9	(336.4 to 749.8)
	YLDs	Ischemic Stroke	Male	1990	48	(31 to 68)	19.8	(13.0 to 27.5)
				2013	92	(59 to 130)	19.5	(12.7 to 28.0)
			Female	1990	55	(35 to 80)	17.8	(11.4 to 25.9)
				2013	104	(67 to 146)	18	(12.0 to 25.3)
			Both	1990	103	(68 to 141)	18.6	(12.5 to 25.4)
				2013	196	(132 to 265)	18.6	(12.6 to 25.8)
		Hemorrhagic						
		Stroke	Male	1990	15	(10 to 23)	5.2	(3.3 to 7.7)
				2013	30	(18 to 43)	5.2	(3.2 to 7.8)
			Female	1990	15	(9 to 23)	4.4	(2.7 to 6.4)
				2013	27	(17 to 39)	4.3	(2.7 to 6.2)
			Both	1990	31	(20 to 45)	4.7	(3.1 to 6.7)
			Both	2013	57	(38 to 80)	4.7	(3.2 to 6.6)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Brazil	Deaths	Ischemic Stroke	Male	1990	24,841	(20,179 to 28,951)	86.3	(70.2 to 101.2)

			2013	36,981	(29,150 to 44,077)	56.9	(45.0 to 68.1)
		Female	1990	24,304	(17,795 to 27,108)	72.4	(53.3 to 80.6)
			2013	37,836	(30,797 to 50,377)	43.1	(35.2 to 57.2)
		Both	1990	49,146	(40,878 to 54,390)	78.9	(65.3 to 87.2)
			2013	74,817	(64,647 to 88,799)	49.2	(42.7 to 58.2)
	Hemorrhagic				, ,		,
Deaths	Stroke	Male	1990	22,706	(18,608 to 27,657)	62	(50.1 to 76.3)
			2013	35,225	(26,486 to 42,496)	47.3	(35.3 to 57.4)
		Female	1990	22,735	(17,864 to 26,112)	54.7	(42.7 to 64.2)
			2013	33,728	(27,983 to 43,770)	36.4	(30.1 to 47.3)
		Both	1990	45,442	(39,138 to 51,192)	58.2	(50.0 to 66.3)
			2013	68,953	(58,773 to 79,266)	41.3	(35.0 to 47.9)
					(369,857 to	1 = 1 5	(00.000)
DALYs	Ischemic Stroke	Male	1990	457,393	538,603)	1323	(1076.0 to 1541.3)
					(455,234 to		·
			2013	578,617	692,263)	805.8	(637.8 to 958.1)
					(267,720 to		
		Female	1990	373,233	420,521)	962.2	(706.8 to 1072.8)
					(387,263 to		
			2013	465,090	620,491)	518.8	(432.8 to 690.9)
					(681,647 to		
		Both	1990	830,627	922,020)	1131.1	(945.3 to 1251.3)
			2012	4 0 4 2 7 0 7	(886,778 to	647.7	(550.6), 764.0)
			2013	1,043,707	1,230,889)	647.7	(552.6 to 761.2)
	Hemorrhagic	Mala	1000	624 572	(528,073 to	1200	(11541+-10041)
	Stroke	Male	1990	634,573	772,437)	1396	(1154.1 to 1684.1)
			2013	831,946	(629,519 to 1,000,392)	981	(739.5 to 1180.1)
			2013	031,340	(475,107 to	301	(733.3 (0 1100.1)
		Female	1990	607,310	674,093)	1205.6	(958.8 to 1354.7)
			2013	709,011	(597,946 to	721.1	(608.9 to 921.0)
	l		2013	/05,011	(337,340 to	/ 41.1	(000.5 to 521.0)

						907,512)		
						(1,077,713 to		
			Both	1990	1,241,884	1,380,959)	1295.2	(1119.5 to 1444.5)
						(1,318,406 to		
				2013	1,540,958	1,764,972)	840.8	(720.7 to 962.7)
	YLDs	Ischemic Stroke	Male	1990	14,146	(8,818 to 20,599)	36.4	(22.6 to 53.3)
				2013	27,025	(17,097 to 38,635)	33.9	(21.6 to 48.7)
			Female	1990	10,255	(6,778 to 14,334)	23.3	(15.3 to 33.5)
				2013	20,500	(13,350 to 28,692)	21.7	(14.2 to 30.4)
			Both	1990	24,401	(15,986 to 33,788)	29.4	(19.4 to 40.9)
				2013	47,526	(32,459 to 64,447)	27.3	(18.7 to 36.8)
		Hemorrhagic						
		Stroke	Male	1990	2,851	(1,764 to 4,247)	6.2	(3.9 to 9.3)
				2013	5,324	(3,315 to 7,601)	6.2	(3.9 to 8.8)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	2,298	(1,445 to 3,401)	4.5	(2.9 to 6.7)
				2013	4,606	(2,956 to 6,590)	4.6	(3.0 to 6.7)
			Both	1990	5,149	(3,352 to 7,229)	5.3	(3.5 to 7.5)
			Both	2013	9,931	(6,734 to 13,706)	5.4	(3.6 to 7.3)
Brunei	Deaths	Ischemic Stroke	Male	1990	25	(18 to 33)	73.5	(55.3 to 96.2)
				2013	34	(25 to 45)	47.9	(35.5 to 62.5)
			Female	1990	29	(20 to 37)	87.5	(63.6 to 109.1)
				2013	45	(35 to 59)	52.4	(41.2 to 70.2)
			Both	1990	54	(43 to 66)	80.5	(65.7 to 96.5)
				2013	80	(66 to 99)	50.6	(41.9 to 62.1)
		Hemorrhagic				, ,		, ,
1	Daatlaa	Stroke	Male	1990	35	(26 to 43)	75.7	(57.2 to 96.1)
	Deaths	Stroke	iviale	1990	33	(20 (0 43)	75.7	(37.2 to 30.1)

		Female	1990	28	(19 to 34)	70.8	(47.6 to 88.5)
			2013	41	(30 to 55)	37.4	(27.7 to 49.9)
		Both	1990	63	(48 to 74)	73.4	(55.9 to 87.3)
			2013	91	(71 to 112)	40	(31.5 to 49.5)
DALYs	Ischemic Stroke	Male	1990	431	(318 to 605)	1043.1	(783.9 to 1398.7)
			2013	641	(495 to 855)	631.5	(492.4 to 811.4)
		Female	1990	500	(323 to 681)	1171.6	(804.5 to 1517.7)
			2013	726	(562 to 937)	621.1	(492.8 to 797.5)
		Both	1990	931	(706 to 1,166)	1104.9	(877.1 to 1342.0)
			2013	1,367	(1,134 to 1,693)	627	(525.2 to 760.3)
	Hemorrhagic			·			
	Stroke	Male	1990	954	(736 to 1,159)	1565.6	(1203.3 to 1921.6)
			2013	1,336	(909 to 1,767)	824.7	(575.7 to 1089.1)
		Female	1990	701	(477 to 896)	1357.4	(939.0 to 1677.7)
			2013	982	(725 to 1,308)	656	(489.3 to 870.2)
		Both	1990	1,655	(1,298 to 1,932)	1466	(1135.1 to 1697.0)
			2013	2,319	(1,822 to 2,806)	741.1	(588.1 to 902.6)
YLDs	Ischemic Stroke	Male	1990	46	(30 to 65)	99.1	(64.6 to 143.5)
			2013	117	(76 to 172)	91.6	(59.3 to 134.2)
		Female	1990	24	(16 to 34)	57.4	(37.9 to 82.0)
			2013	71	(47 to 100)	58	(37.4 to 81.0)
		Both	1990	70	(47 to 98)	79.1	(53.0 to 111.4)
			2013	189	(127 to 263)	74.6	(50.2 to 103.7)
	Hemorrhagic		40.5.5	4.5	(10 :)	00.5	(0.4.0
	Stroke	Male	1990	19	(12 to 27)	32.9	(21.2 to 48.7)
			2013	47	(29 to 69)	32.1	(20.2 to 47.2)
		Female	1990	11	(7 to 15)	20.3	(12.9 to 28.5)
			2013	35	(22 to 50)	23.6	(15.1 to 33.9)

			Both	1990	30	(20 to 42)	26.9	(18.0 to 37.5)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	2013	82	(54 to 115)	27.8	(18.4 to 39.3)
Bulgaria	Deaths	Ischemic Stroke	Male	1990	7,551	(6,430 to 8,864)	171.1	(145.5 to 200.8)
				2013	8,540	(6,810 to 9,937)	156.6	(124.3 to 183.0)
			Female	1990	8,593	(7,237 to 9,876)	146.5	(123.5 to 169.2)
				2013	10,243	(7,710 to 11,773)	115.5	(86.8 to 132.9)
			Both	1990	16,144	(14,401 to 17,923)	157.6	(141.0 to 175.1)
				2013	18,784	(15,320 to 20,808)	132.8	(108.5 to 147.0)
		Hemorrhagic			-			
	Deaths	Stroke	Male	1990	4,910	(3,804 to 5,840)	100.6	(77.6 to 119.8)
				2013	3,691	(2,656 to 5,403)	68.1	(49.0 to 99.0)
			Female	1990	5,513	(4,010 to 6,165)	90.7	(65.4 to 101.5)
				2013	3,771	(2,740 to 6,452)	46.8	(34.7 to 78.9)
			Both	1990	10,423	(8,423 to 11,655)	95.7	(77.3 to 107.2)
				2013	7,462	(5,548 to 11,181)	56.4	(42.4 to 83.2)
						(95,194 to		
	DALYs	Ischemic Stroke	Male	1990	113,546	132,999)	2354.3	(1991.8 to 2757.0)
						(93,335 to		
				2013	116,341	135,754)	2118	(1694.9 to 2474.8)
				1000	100 200	(90,282 to	4000.0	(4504.5 + 2064.2)
			Female	1990	109,308	124,809)	1808.8	(1501.5 to 2064.3)
				2013	109,150	(82,045 to 125,223)	1303	(980.2 to 1494.5)
				2013	103,130	(195,821 to	1303	(300.2 to 1434.3)
			Both	1990	222,855	248,704)	2059.5	(1818.5 to 2290.2)
					,==0	(187,684 to		(
				2013	225,491	250,159)	1658.8	(1378.2 to 1844.6)
		Hemorrhagic	Male	1990	102,354	(80,597 to	1979.4	(1555.0 to 2326.8)

	Stroke				120,725)		
			2013	71,454	(54,174 to 99,043)	1356.6	(1033.7 to 1869.7)
					(70,593 to		
		Female	1990	96,364	105,892)	1579.8	(1152.4 to 1735.5)
			2013	55,602	(42,338 to 91,901)	795	(618.5 to 1278.5)
					(162,235 to		
		Both	1990	198,719	· · · · · · · · · · · · · · · · · · ·	1774.6	(1446.6 to 1975.8)
					• •		(
					,		(829.8 to 1461.6)
YLDs	Ischemic Stroke	Male	1990	1,180	(756 to 1,676)		(14.9 to 32.1)
			2013	1,318	(862 to 1,948)	25.3	(16.5 to 37.1)
		Female	1990	876	(572 to 1,238)	14.8	(9.7 to 20.8)
			2013	969	(621 to 1,422)	14.7	(9.7 to 21.0)
		Both	1990	2,056	(1,397 to 2,871)	18.6	(12.6 to 25.6)
			2013	2,288	(1,550 to 3,208)	19.5	(13.2 to 27.4)
	Hemorrhagic	Male	1990	243	(138 to 382)	4.9	(2.8 to 7.6)
	Stroke						
							100,000, Age-standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			2013	251	(141 to 385)	5.2	(3.0 to 7.8)
		Female	1990	196	(116 to 296)	3.5	(2.2 to 5.2)
			2013	206	(123 to 316)	3.6	(2.2 to 5.4)
		Both	1990	439	(272 to 633)	4.2	(2.6 to 6.0)
		Both	2013	458	(292 to 649)	4.3	(2.8 to 6.2)
Deaths	Ischemic Stroke	Male	1990	364	(145 to 543)	30.2	(12.5 to 45.1)
			2013	1,057	(269 to 1,505)	65.8	(19.1 to 91.2)
		Female	1990	445	(172 to 722)	28.5	(11.6 to 45.5)
			2013	1,258	(327 to 1,860)	56.3	(14.7 to 81.6)
		YLDs Ischemic Stroke Hemorrhagic Stroke measure Stroke Type	Female Both YLDs Ischemic Stroke Male Female Female Both Hemorrhagic Stroke Male Female Both Female Both Both Hemorrhagic Stroke Male Stroke Male	Pemale 1990 2013	Female	2013	Semale

Both 1990 809 (491 to 1,082) 29.2 (17.6 to 39.4)	1		, ,			1	-	
Hemorrhagic Stroke Male 1990 426 (257 to 614) 29 (16.2 to 43.9)			Both	1990	809	(491 to 1,082)	29.2	(17.6 to 39.4)
Deaths Stroke Male 1990 426 (257 to 614) 29 (16.2 to 43.9) 1 2013 804 (447 to 1,516) 37.7 (19.1 to 80.1) 1 Female 1990 520 (294 to 840) 27.5 (15.2 to 45.7) 2013 896 (576 to 1,694) 30.9 (18.9 to 63.7) (661 to 1,285) 28.2 (18.7 to 40.0) 2013 1,700 (1,135 to 2,940) 33.7 (21.2 to 62.2) DALYS Ischemic Stroke Male 1990 8,449 (3,236 to 12,678) 550.7 (223.6 to 817.6) 2013 22,850 (5,652 to 33,576) 1077.3 (290.6 to 1523.8) (29.6 to 1523.8) Female 1990 9,650 (3,441 to 15,590) 487.4 (183.7 to 781.2) 2013 23,627 (5,940 to 37,025) 834.5 (222.2 to 1236.0) Both 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7) Hemorrhagic Stroke Male 1990 12,909 (2013	2,315	(981 to 3,180)	60.2	(25.9 to 81.3)
2013 804 (447 to 1,516) 37.7 (19.1 to 80.1)								
Female 1990 520 (294 to 840) 27.5 (15.2 to 45.7) 2013 896 (576 to 1,694) 30.9 (18.9 to 63.7) Both 1990 947 (661 to 1,285) 28.2 (18.7 to 40.0) 2013 1,700 (1,135 to 2,940) 33.7 (21.2 to 62.2) DALYS Ischemic Stroke Male 1990 8,449 (3,236 to 12,678) 550.7 (223.6 to 817.6) Female 1990 9,650 (3,441 to 15,590) 487.4 (183.7 to 781.2) 2013 23,627 (5,940 to 37,025) 834.5 (222.2 to 1236.0) Both 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7) Both 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) Female 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) Female 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDS Ischemic Stroke Male 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 655) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)	Deaths	Stroke	Male	1990		(257 to 614)	29	(16.2 to 43.9)
Both 1990 947 (661 to 1,694) 30.9 (18.9 to 63.7)				2013	804	(447 to 1,516)	37.7	(19.1 to 80.1)
Both 1990 947 (661 to 1,285) 28.2 (18.7 to 40.0)			Female	1990	520	(294 to 840)	27.5	(15.2 to 45.7)
DALYS Ischemic Stroke Male 1990 8,449 (3,236 to 12,678) 550.7 (223.6 to 817.6)				2013	896	(576 to 1,694)	30.9	(18.9 to 63.7)
DALYS Ischemic Stroke Male 1990 8,449 (3,236 to 12,678) 550.7 (223.6 to 817.6)			Both	1990	947	(661 to 1,285)	28.2	(18.7 to 40.0)
2013 22,850 (5,652 to 33,576) 1077.3 (290.6 to 1523.8) Female 1990 9,650 (3,441 to 15,590) 487.4 (183.7 to 781.2) 2013 23,627 (5,940 to 37,025) 834.5 (222.2 to 1236.0) Both 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7) 2013 46,477 (19,143 to 65,753) 935.6 (397.4 to 1282.2) Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) 2013 23,553 (14,077 to 40,405) 777.3 (426.0 to 1495.0) Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) 2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) 2013 47,586 (34,081 to 74,078) 688 (460.0 to 1196.2) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) 2013 447 (283 to 655) 11.8 (7.7 to 17.4)				2013	1,700	(1,135 to 2,940)	33.7	(21.2 to 62.2)
Female 1990 9,650 (3,441 to 15,590) 487.4 (183.7 to 781.2) 2013 23,627 (5,940 to 37,025) 834.5 (222.2 to 1236.0) 8oth 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7) 2013 46,477 (19,143 to 65,753) 935.6 (397.4 to 1282.2) Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) 2013 23,553 (14,077 to 40,405) 777.3 (426.0 to 1495.0) Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) 2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) 8oth 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) YLDs Ischemic Stroke Male 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) 8oth 1990 516 (340 to 724) 13.3 (8.8 to 18.4)	DALYs	Ischemic Stroke	Male	1990	8,449	(3,236 to 12,678)	550.7	(223.6 to 817.6)
Both 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7)				2013	22,850	(5,652 to 33,576)	1077.3	(290.6 to 1523.8)
Both 1990 18,100 (10,485 to 24,399) 514.3 (311.2 to 683.7) Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) Emale 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) Emale 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) Emale 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) Emale 1990 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Emale 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) YLDs Ischemic Stroke Male 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 364) 11.3 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3			Female	1990	9,650	(3,441 to 15,590)	487.4	(183.7 to 781.2)
Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.4 to 1282.2)				2013	23,627	(5,940 to 37,025)	834.5	(222.2 to 1236.0)
Hemorrhagic Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) 2013 23,553 (14,077 to 40,405) 777.3 (426.0 to 1495.0) Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) 2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) YLDs Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 365) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)			Both	1990	18,100	(10,485 to 24,399)	514.3	(311.2 to 683.7)
Stroke Male 1990 12,909 (8,240 to 17,651) 653.7 (397.2 to 932.5) 2013 23,553 (14,077 to 40,405) 777.3 (426.0 to 1495.0) Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) 2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) YLDs Ischemic Stroke Male 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 364) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)				2013	46,477	(19,143 to 65,753)	935.6	(397.4 to 1282.2)
2013 23,553 (14,077 to 40,405) 777.3 (426.0 to 1495.0)		Hemorrhagic						
Female 1990 14,785 (8,918 to 21,442) 606.1 (346.7 to 942.6) 2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) 2013 47,586 (34,081 to 74,078) 688 (460.0 to 1196.2) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) 2013 446 (285 to 637) 14.9 (9.5 to 21.3) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 516 (340 to 724) 13.3 (8.8 to 18.4)		Stroke	Male	1990	12,909	(8,240 to 17,651)	653.7	(397.2 to 932.5)
2013 24,033 (15,881 to 40,806) 624.8 (403.0 to 1156.2) Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) 2013 47,586 (34,081 to 74,078) 688 (460.0 to 1196.2) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) 2013 446 (285 to 637) 14.9 (9.5 to 21.3) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) 2013 447 (283 to 655) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)				2013	23,553	(14,077 to 40,405)	777.3	(426.0 to 1495.0)
Both 1990 27,694 (19,990 to 35,545) 626.5 (439.8 to 835.4) YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) VLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) VLDs Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Female 1990 249 (158 to 364) 11.3 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)			Female	1990	14,785	(8,918 to 21,442)	606.1	(346.7 to 942.6)
YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) Pemale 1990 249 (158 to 364) 11.3 (7.1 to 16.3) Pemale 1990 249 (158 to 364) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)				2013	24,033	(15,881 to 40,806)	624.8	(403.0 to 1156.2)
YLDs Ischemic Stroke Male 1990 267 (175 to 381) 15.9 (10.2 to 22.5) 2013 446 (285 to 637) 14.9 (9.5 to 21.3) Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) 2013 447 (283 to 655) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)			Both	1990	27,694	(19,990 to 35,545)	626.5	(439.8 to 835.4)
2013 446 (285 to 637) 14.9 (9.5 to 21.3)				2013	47,586	(34,081 to 74,078)	688	(460.0 to 1196.2)
Female 1990 249 (158 to 364) 11.3 (7.1 to 16.3) 2013 447 (283 to 655) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)	YLDs	Ischemic Stroke	Male	1990	267	(175 to 381)	15.9	(10.2 to 22.5)
2013 447 (283 to 655) 11.8 (7.7 to 17.4) Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)				2013	446	(285 to 637)	14.9	(9.5 to 21.3)
Both 1990 516 (340 to 724) 13.3 (8.8 to 18.4)			Female	1990	249	(158 to 364)	11.3	(7.1 to 16.3)
				2013	447	(283 to 655)	11.8	(7.7 to 17.4)
2013 893 (592 to 1,243) 13.1 (8.8 to 18.2)			Both	1990	516	(340 to 724)	13.3	(8.8 to 18.4)
				2013	893	(592 to 1,243)	13.1	(8.8 to 18.2)

		Hemorrhagic						
		Stroke	Male	1990	84	(53 to 123)	4.2	(2.6 to 6.2)
				2013	149	(90 to 217)	3.9	(2.4 to 5.7)
			Female	1990	72	(45 to 106)	2.9	(1.8 to 4.4)
				2013	121	(74 to 178)	2.7	(1.6 to 4.1)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	156	(103 to 218)	3.5	(2.3 to 4.8)
			Both	2013	270	(173 to 381)	3.2	(2.0 to 4.5)
Burundi	Deaths	Ischemic Stroke	Male	1990	614	(42 to 1,086)	90.6	(6.3 to 157.3)
				2013	816	(174 to 1,312)	72.8	(15.9 to 114.4)
			Female	1990	799	(71 to 1,503)	92.2	(8.9 to 171.7)
				2013	1,012	(390 to 1,655)	80.5	(31.5 to 130.2)
			Both	1990	1,413	(171 to 2,316)	91.7	(12.0 to 150.1)
				2013	1,829	(727 to 2,585)	77.3	(32.0 to 108.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,259	(829 to 2,003)	148.8	(94.2 to 245.7)
				2013	1,436	(976 to 2,197)	99.4	(63.9 to 162.3)
			Female	1990	1,643	(967 to 2,778)	157.7	(91.3 to 280.1)
				2013	1,601	(1,023 to 2,899)	105.7	(64.9 to 198.0)
			Both	1990	2,903	(1,994 to 4,315)	154	(103.0 to 240.4)
				2013	3,038	(2,216 to 4,632)	103	(72.5 to 167.2)
	DALYs	Ischemic Stroke	Male	1990	12,176	(888 to 22,088)	1450.3	(110.6 to 2569.4)
				2013	15,514	(3,177 to 25,631)	1090.4	(240.7 to 1748.4)
			Female	1990	14,210	(1,279 to 27,148)	1390.1	(127.8 to 2656.8)
				2013	16,592	(6,004 to 27,423)	1126.1	(420.8 to 1846.5)
			Both	1990	26,386	(3,073 to 42,885)	1420.2	(176.8 to 2327.0)
				2013	32,106	(11,996 to 45,909)	1114.4	(441.2 to 1583.9)

		Hemorrhagic						
		Stroke	Male	1990	33,465	(22,511 to 51,334)	3092.3	(2052.6 to 4877.6)
				2013	37,529	(26,587 to 53,374)	1941.8	(1325.9 to 2955.5)
			Female	1990	39,894	(23,039 to 63,947)	3178.7	(1891.7 to 5263.7)
				2013	36,679	(24,427 to 63,078)	1947.5	(1259.8 to 3464.7)
						(51,954 to		
			Both	1990	73,360	104,544)	3144.9	(2175.9 to 4592.3)
						(56,573 to		(
				2013	74,209	107,511)	1949.5	(1436.4 to 2939.2)
	YLDs	Ischemic Stroke	Male	1990	121	(77 to 181)	10.6	(6.9 to 15.9)
				2013	218	(135 to 324)	10.3	(6.4 to 15.0)
			Female	1990	95	(59 to 139)	7.1	(4.6 to 10.4)
				2013	170	(108 to 247)	7.5	(4.9 to 10.7)
			Both	1990	217	(145 to 305)	8.7	(5.9 to 12.3)
				2013	388	(259 to 541)	8.8	(5.9 to 12.2)
		Hemorrhagic						
		Stroke	Male	1990	41	(26 to 64)	3	(1.9 to 4.4)
				2013	78	(47 to 118)	2.9	(1.7 to 4.3)
			Female	1990	28	(17 to 41)	1.8	(1.1 to 2.6)
				2013	51	(31 to 79)	1.9	(1.1 to 2.8)
			Both	1990	70	(45 to 100)	2.3	(1.6 to 3.3)
			Both	2013	130	(82 to 188)	2.4	(1.5 to 3.4)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Cambodia	Deaths	Ischemic Stroke	Male	1990	1,100	(702 to 1,721)	101.2	(62.9 to 152.6)
Carribodia		isomerms stroke						
Cambodia				2013	3,738	(2,650 to 4,710)	105.8	(74.8 to 133.2)
Camboula			Female	2013 1990	3,738 1,387	(2,650 to 4,710) (928 to 1,945)	105.8 83.4	(74.8 to 133.2) (57.0 to 115.2)
Cambodia			Female		-			, ,

			2013	9,129	(6,804 to 10,695)	107	(80.2 to 125.2)
	Hemorrhagic						
Deaths	Stroke	Male	1990	1,860	(1,215 to 2,423)	112.4	(62.3 to 158.0)
			2013	3,415	(2,461 to 4,745)	83.1	(60.6 to 118.5)
		Female	1990	1,749	(1,106 to 2,430)	85.3	(53.9 to 120.0)
			2013	4,051	(2,878 to 6,381)	75.7	(54.7 to 120.4)
		Both	1990	3,610	(2,711 to 4,389)	97.1	(71.0 to 122.1)
			2013	7,466	(5,863 to 10,474)	79.1	(62.0 to 113.2)
DALYs	Ischemic Stroke	Male	1990	26,084	(17,403 to 41,044)	1637.9	(1048.4 to 2550.6)
			2013	61,879	(43,887 to 80,155)	1607.9	(1149.1 to 2064.2)
		Female	1990	25,009	(14,687 to 37,289)	1232.4	(791.1 to 1743.6)
			2013	74,237	(50,772 to 95,778)	1445.1	(998.8 to 1852.9)
		Both	1990	51,094	(37,018 to 67,482)	1409.7	(1048.9 to 1818.5)
					(100,312 to		
			2013	136,117	163,382)	1517.2	(1128.4 to 1812.6)
	Hemorrhagic						
	Stroke	Male	1990	64,081	(47,386 to 81,495)	2641.4	(1698.8 to 3462.0)
			2012	05.022	(68,572 to	1026.6	(4.402.4 +- 2662.0)
		Famala	2013	95,033	127,997)	1936.6	(1402.1 to 2662.0)
		Female	1990	49,488	(30,006 to 64,786) (64,848 to	1838.5	(1164.0 to 2504.5)
			2013	91,073	133,682)	1560.2	(1106.4 to 2330.7)
			2015	31,073	(85,910 to	1300.2	(110011 to 20017)
		Both	1990	113,569	137,698)	2204.9	(1663.6 to 2669.9)
					(148,691 to		
			2013	186,107	241,297)	1733.6	(1371.3 to 2294.3)
YLDs	Ischemic Stroke	Male	1990	338	(220 to 485)	22.7	(14.7 to 32.2)
			2013	1,021	(664 to 1,426)	26.6	(17.6 to 37.9)
		Female	1990	402	(254 to 576)	20	(12.5 to 28.8)
			2013	1,191	(728 to 1,726)	22.7	(13.9 to 32.8)

			Both	1990	740	(497 to 1,023)	21	(13.9 to 28.9)
				2013	2,213	(1,460 to 3,101)	24.3	(15.9 to 34.3)
		Hemorrhagic						
		Stroke	Male	1990	434	(281 to 605)	25.1	(16.7 to 35.2)
				2013	1,205	(812 to 1,695)	28.7	(19.3 to 40.4)
			Female	1990	438	(283 to 625)	19.9	(12.7 to 28.3)
				2013	1,186	(774 to 1,691)	21.6	(14.0 to 30.5)
			Both	1990	873	(593 to 1,184)	22	(15.1 to 30.0)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	2013	2,391	(1,607 to 3,320)	24.6	(16.7 to 34.3)
Cameroon	Deaths	Ischemic Stroke	Male	1990	1,392	(1,018 to 1,758)	76.8	(57.1 to 96.9)
				2013	3,585	(1,986 to 4,791)	114.6	(65.7 to 150.9)
			Female	1990	2,024	(1,443 to 2,840)	94.5	(67.2 to 134.5)
				2013	3,583	(2,131 to 4,875)	101.3	(61.4 to 138.2)
			Both	1990	3,417	(2,754 to 4,344)	86.9	(70.7 to 110.4)
				2013	7,169	(4,606 to 9,022)	107.7	(70.7 to 135.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,340	(877 to 1,738)	57.7	(37.1 to 76.3)
				2013	2,575	(1,498 to 4,301)	62.3	(33.5 to 113.7)
			Female	1990	1,713	(1,204 to 2,374)	66.5	(45.9 to 92.5)
				2013	2,287	(1,363 to 4,860)	52.1	(30.0 to 117.3)
			Both	1990	3,053	(2,454 to 3,796)	62.6	(49.6 to 78.7)
				2013	4,863	(3,188 to 8,777)	57	(35.3 to 110.9)
	DALYs	Ischemic Stroke	Male	1990	29,132	(21,104 to 38,805)	1230.7	(900.4 to 1549.3)
				2013	71,707	(37,042 to 97,572)	1827.9	(1002.2 to 2466.7)
			Female	1990	37,923	(26,982 to 53,913)	1431.5	(1019.8 to 2005.0)
				2013	60,187	(35,034 to 85,234)	1431.7	(840.6 to 1968.9)

		Both	1990	67,056	(52,193 to 88,002)	1343.3	(1076.8 to 1693.8)
					(80,529 to		
			2013	131,895	167,596)	1622.1	(1017.6 to 2051.5)
	Hemorrhagic						
	Stroke	Male	1990	41,930	(27,938 to 53,939)	1225.4	(815.4 to 1591.3)
					(51,913 to		
			2013	81,471	123,263)	1381.3	(809.3 to 2306.6)
		Female	1990	47,096	(33,518 to 64,633)	1348.6	(963.7 to 1856.1)
					(39,954 to		
			2013	63,193	128,170)	1062.2	(646.4 to 2190.3)
					(70,611 to		
		Both	1990	89,026	110,682)	1293.2	(1043.4 to 1608.4)
					(101,731 to		
			2013	144,665	237,873)	1217.2	(812.5 to 2159.0)
YLDs	Ischemic Stroke	Male	1990	416	(270 to 597)	15.4	(10.2 to 22.1)
			2013	756	(499 to 1,057)	16	(10.7 to 22.4)
		Female	1990	339	(221 to 481)	11.2	(7.2 to 16.0)
			2013	600	(382 to 851)	11.4	(7.3 to 16.3)
		Both	1990	756	(515 to 1,044)	13.1	(9.1 to 18.1)
			2013	1,356	(920 to 1,859)	13.6	(9.3 to 18.4)
	Hemorrhagic						
	Stroke	Male	1990	133	(82 to 195)	4.2	(2.6 to 6.2)
			2013	232	(145 to 332)	4	(2.5 to 5.7)
		Female	1990	96	(57 to 140)	2.9	(1.7 to 4.1)
			2013	166	(104 to 250)	2.7	(1.7 to 4.2)
				Cou			100,000, Age-standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Both	1990	229	(148 to 321)	3.5	(2.3 to 4.9)
		Both	2013	398	(257 to 564)	3.3	(2.2 to 4.7)
		YLDs Ischemic Stroke Hemorrhagic Stroke	Hemorrhagic Stroke Male Female Both YLDs Ischemic Stroke Male Female Hemorrhagic Stroke Both Female Female Female Both	Hemorrhagic Stroke Male 1990	Hemorrhagic Stroke Male 1990 41,930	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 41,930 (27,938 to 53,939) 1225.4

Canada	Deaths	Ischemic Stroke	Male	1990	3,669	(2,918 to 4,595)	32.4	(25.7 to 40.9)
				2013	3,595	(2,493 to 4,636)	14.5	(10.0 to 18.6)
			Female	1990	5,096	(3,946 to 6,297)	27.6	(21.4 to 34.1)
				2013	6,098	(4,498 to 8,497)	16	(11.9 to 22.3)
			Both	1990	8,765	(7,463 to 10,297)	29.8	(25.4 to 35.1)
				2013	9,693	(7,664 to 12,281)	15.6	(12.3 to 19.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,694	(2,071 to 3,460)	22.2	(16.7 to 28.6)
				2013	3,125	(1,950 to 4,087)	12.7	(8.0 to 16.5)
			Female	1990	3,491	(2,726 to 4,626)	19.9	(15.7 to 26.2)
				2013	4,708	(3,195 to 6,133)	13.7	(9.5 to 17.3)
			Both	1990	6,186	(5,302 to 7,752)	21	(18.0 to 26.4)
				2013	7,834	(5,781 to 9,439)	13.3	(10.0 to 15.9)
	DALYs	Ischemic Stroke	Male	1990	63,456	(51,961 to 76,493)	512.5	(419.5 to 621.1)
				2013	64,598	(49,405 to 80,761)	267.6	(205.1 to 334.5)
			Female	1990	62,609	(52,818 to 76,077)	356.5	(299.5 to 432.4)
				2013	72,506	(56,515 to 91,590)	229.7	(179.3 to 290.4)
						(109,888 to		
			Both	1990	126,066	147,102)	424	(369.8 to 494.2)
						(110,955 to		
				2013	137,104	167,209)	248	(200.5 to 302.3)
		Hemorrhagic						
		Stroke	Male	1990	52,275	(42,341 to 66,103)	395.1	(318.3 to 498.8)
				2013	48,388	(33,659 to 60,069)	204	(142.5 to 252.6)
			Female	1990	54,785	(45,174 to 70,701)	337.5	(278.8 to 436.1)
				2013	57,455	(42,626 to 71,042)	203.8	(152.8 to 252.4)
						(94,509 to		
			Both	1990	107,061	128,050)	362.5	(319.6 to 433.7)
				2012	405.011	(83,239 to	200.0	(464.4 : 204.6)
				2013	105,844	122,228)	203.9	(161.1 to 234.1)

	YLDs	Ischemic Stroke	Male	1990	16,602	(10,873 to 23,077)	129.3	(85.0 to 180.1)
				2013	28,884	(19,190 to 41,027)	120.2	(80.0 to 170.2)
			Female	1990	15,758	(9,838 to 23,339)	94.1	(58.3 to 138.9)
				2013	28,122	(17,807 to 40,763)	98	(61.1 to 141.4)
			Both	1990	32,360	(22,091 to 46,012)	109.2	(74.7 to 154.9)
				2013	57,007	(38,122 to 79,385)	108.1	(72.3 to 150.2)
		Hemorrhagic						
		Stroke	Male	1990	2,491	(1,605 to 3,524)	18.8	(12.0 to 26.6)
				2013	4,251	(2,651 to 6,166)	18.3	(11.5 to 26.4)
			Female	1990	2,998	(1,836 to 4,519)	18.5	(11.3 to 27.9)
				2013	5,482	(3,361 to 7,948)	20.6	(12.7 to 30.1)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	5,490	(3,680 to 7,677)	18.5	(12.4 to 26.0)
			Both	2013	9,733	(6,305 to 13,695)	19.4	(12.5 to 27.4)
Cape Verde	Deaths	Ischemic Stroke	Male	1990	88	(70 to 111)	123.8	(99.2 to 155.3)
				2013	173	(117 to 241)	146.6	(99.9 to 203.1)
			Female	1990	101	(70 to 128)	110.5	(77.4 to 139.4)
				2013	127	(77 to 259)	65.1	(39.5 to 134.2)
			Both	1990	189	(150 to 228)	116.4	(92.9 to 140.7)
				2013	300	(218 to 440)	97.5	(70.7 to 143.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	72	(54 to 98)	92.1	(68.0 to 126.7)
				2013	79	(44 to 140)	59.4	(32.4 to 107.6)
			Female	1990	73	(53 to 102)	76.3	(55.5 to 107.5)
<u>-</u>				2013	45	(17 to 144)	24.1	(9.3 to 76.6)
			Both	1990	146	(112 to 191)	83.1	(62.9 to 107.7)

DALYs	Ischemic Stroke	Male	1990	1,591	(1,234 to 2,065)	1973.3	(1542.1 to 2527.2)
			2013	2,853	(1,817 to 4,139)	2247.2	(1453.4 to 3218.0)
		Female	1990	1,389	(948 to 1,836)	1375.7	(950.8 to 1780.3)
			2013	1,339	(802 to 2,953)	719	(415.6 to 1602.8)
		Both	1990	2,980	(2,321 to 3,687)	1630.1	(1271.4 to 2003.5)
			2013	4,192	(2,919 to 6,231)	1355.1	(942.4 to 2009.9)
	Hemorrhagic						
	Stroke	Male	1990	2,000	(1,527 to 2,682)	2011.8	(1548.7 to 2663.2)
			2013	2,161	(1,294 to 3,542)	1314.5	(754.3 to 2274.2)
		Female	1990	1,391	(1,002 to 1,976)	1187.4	(851.8 to 1698.5)
			2013	646	(233 to 2,068)	356.2	(127.2 to 1148.0)
		Both	1990	3,392	(2,660 to 4,477)	1538.8	(1205.8 to 2041.5)
			2013	2,808	(1,610 to 5,134)	792.1	(438.5 to 1506.9)
YLDs	Ischemic Stroke	Male	1990	12	(8 to 18)	15.7	(10.3 to 22.7)
			2013	21	(13 to 32)	15.3	(9.7 to 21.9)
		Female	1990	12	(8 to 17)	11.4	(7.4 to 15.7)
			2013	22	(14 to 31)	11.9	(7.6 to 17.0)
		Both	1990	25	(16 to 33)	13.2	(8.9 to 18.0)
							(8.8 to 18.5)
			2013	43	(29 to 60)	13.4	
	Hemorrhagic						
	Stroke	Male	1990	3	(2 to 5)	4.2	(2.6 to 6.2)
			2013	6	(4 to 9)	3.9	(2.4 to 5.8)
		Female	1990	3	(2 to 4)	2.9	(1.8 to 4.2)
			2013	5	(3 to 7)	2.8	(1.8 to 4.1)
		Both	1990	6	(4 to 9)	3.4	(2.2 to 4.8)
		Both	2013	11	(8 to 16)	3.3	(2.2 to 4.6)
				C	nte Allagae	Doto co	100 000 Age-standardized

Counts, All ages

Rate per 100,000, Age-standardized

Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Central African	Deaths	Ischemic Stroke	Male	1990	437	(236 to 638)	87.5	(48.6 to 121.9)
Republic				2013	359	(113 to 565)	50.8	(16.4 to 78.9)
			Female	1990	713	(431 to 1,008)	121.3	(72.8 to 170.5)
				2013	996	(329 to 1,666)	108.9	(34.9 to 177.6)
			Both	1990	1,151	(726 to 1,474)	107.3	(67.6 to 136.7)
				2013	1,356	(512 to 2,000)	84.9	(32.6 to 124.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	533	(326 to 756)	83.9	(50.3 to 119.6)
				2013	428	(220 to 700)	48.7	(24.1 to 84.2)
			Female	1990	805	(439 to 1,133)	112.3	(61.2 to 160.2)
				2013	1,001	(353 to 1,651)	93	(31.7 to 153.9)
			Both	1990	1,339	(903 to 1,752)	100	(66.5 to 132.3)
				2013	1,429	(755 to 2,071)	73.9	(38.0 to 108.0)
	DALYs	Ischemic Stroke	Male	1990	9,296	(4,958 to 14,782)	1469.6	(788.7 to 2174.5)
				2013	6,980	(2,098 to 11,481)	795.9	(253.4 to 1261.7)
			Female	1990	12,884	(7,638 to 19,650)	1810.7	(1093.4 to 2639.1)
				2013	16,866	(5,644 to 30,334)	1569.7	(531.8 to 2681.6)
			Both	1990	22,181	(13,660 to 29,917)	1667.1	(1045.8 to 2147.7)
				2013	23,847	(8,643 to 36,230)	1238.5	(467.2 to 1830.3)
		Hemorrhagic						
		Stroke	Male	1990	15,615	(9,443 to 22,498)	1948	(1173.7 to 2787.6)
				2013	11,737	(6,201 to 18,471)	1026.6	(535.1 to 1658.4)
			Female	1990	20,992	(11,426 to 29,304)	2381.1	(1297.7 to 3355.2)
				2013	25,050	(9,327 to 40,342)	1890.8	(678.5 to 3083.3)
			Both	1990	36,608	(24,813 to 48,419)	2187.8	(1487.3 to 2880.3)
				2013	36,788	(20,066 to 53,623)	1501.6	(808.9 to 2169.7)
	YLDs	Ischemic Stroke	Male	1990	105	(69 to 147)	14.7	(9.7 to 20.4)
				2013	150	(91 to 215)	13.9	(8.6 to 19.6)

			Female	1990	90	(57 to 127)	10.8	(6.9 to 15.3)
				2013	142	(91 to 203)	11.3	(7.3 to 16.4)
			Both	1990	195	(132 to 266)	12.6	(8.5 to 17.2)
				2013	293	(194 to 404)	12.5	(8.2 to 17.2)
		Hemorrhagic						
		Stroke	Male	1990	31	(20 to 45)	3.8	(2.5 to 5.5)
				2013	47	(29 to 70)	3.7	(2.3 to 5.5)
			Female	1990	24	(15 to 37)	2.7	(1.7 to 4.1)
				2013	36	(22 to 56)	2.6	(1.5 to 4.0)
			Both	1990	56	(37 to 80)	3.2	(2.1 to 4.6)
			Both	2013	84	(54 to 119)	3.1	(2.0 to 4.4)
Chad	Deaths	Ischemic Stroke	Male	1990	343	(171 to 538)	42.6	(21.6 to 64.8)
				2013	775	(210 to 1,253)	56.2	(16.1 to 88.4)
				2013		(210 to 1,253) nts, All ages		(16.1 to 88.4) 100,000, Age-standardized
Country	measure	Stroke Type	sex	2013 year				·
Country	measure	Stroke Type	sex Female		Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type		year	Cou Mean	nts, All ages 95% UI	Rate per Mean	100,000, Age-standardized 95% UI
Country	measure	Stroke Type		year 1990	Cou Mean 519	nts, All ages 95% UI (225 to 733)	Rate per Mean 54.3	100,000, Age-standardized 95% UI (25.3 to 77.4)
Country	measure	Stroke Type	Female	year 1990 2013	Cou Mean 519 1,115	nts, All ages 95% UI (225 to 733) (211 to 1,734)	Rate per Mean 54.3 71.7	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5)
Country	measure	Stroke Type Hemorrhagic	Female	year 1990 2013 1990	Cou Mean 519 1,115 862	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157)	Rate per Mean 54.3 71.7 49.4	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9)
Country	measure		Female	year 1990 2013 1990	Cou Mean 519 1,115 862	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157)	Rate per Mean 54.3 71.7 49.4	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9)
Country		Hemorrhagic	Female Both	year 1990 2013 1990 2013	Cou Mean 519 1,115 862 1,891	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157) (678 to 2,728)	Rate per Mean 54.3 71.7 49.4 65.1	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9) (24.1 to 93.4)
Country		Hemorrhagic	Female Both	year 1990 2013 1990 2013	Cou Mean 519 1,115 862 1,891	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157) (678 to 2,728) (210 to 555)	Rate per Mean 54.3 71.7 49.4 65.1 37.5	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9) (24.1 to 93.4) (16.6 to 54.5)
Country		Hemorrhagic	Both Male	year 1990 2013 1990 2013 1990 2013	Mean 519 1,115 862 1,891 404 618	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157) (678 to 2,728) (210 to 555) (343 to 1,150)	Rate per Mean 54.3 71.7 49.4 65.1 37.5 31.8	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9) (24.1 to 93.4) (16.6 to 54.5) (14.9 to 66.9)
Country		Hemorrhagic	Both Male	year 1990 2013 1990 2013 1990 2013 1990	Cou Mean 519 1,115 862 1,891 404 618 573	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157) (678 to 2,728) (210 to 555) (343 to 1,150) (319 to 808)	Rate per Mean 54.3 71.7 49.4 65.1 37.5 31.8 48.6	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9) (24.1 to 93.4) (16.6 to 54.5) (14.9 to 66.9) (26.3 to 72.3)
Country		Hemorrhagic	Both Male Female	year 1990 2013 1990 2013 1990 2013 1990 2013	Mean 519 1,115 862 1,891 404 618 573 831	nts, All ages 95% UI (225 to 733) (211 to 1,734) (492 to 1,157) (678 to 2,728) (210 to 555) (343 to 1,150) (319 to 808) (478 to 1,582)	Rate per Mean 54.3 71.7 49.4 65.1 37.5 31.8 48.6 40.3	100,000, Age-standardized 95% UI (25.3 to 77.4) (15.1 to 110.5) (28.9 to 66.9) (24.1 to 93.4) (16.6 to 54.5) (14.9 to 66.9) (26.3 to 72.3) (21.8 to 85.3)

				2013	16,270	(4,122 to 27,205)	889	(243.8 to 1442.0)
			Female	1990	10,090	(4,194 to 14,549)	810.2	(346.4 to 1132.9)
				2013	20,195	(3,721 to 32,145)	1025.2	(198.5 to 1611.5)
			Both	1990	17,675	(9,978 to 24,111)	755.5	(432.1 to 1011.9)
				2013	36,466	(12,897 to 53,663)	966.2	(351.7 to 1410.2)
		Hemorrhagic						
		Stroke	Male	1990	13,638	(8,428 to 18,251)	797.9	(410.5 to 1102.8)
				2013	21,207	(13,385 to 34,286)	688	(369.6 to 1290.2)
			Female	1990	16,673	(9,701 to 22,316)	958.9	(542.2 to 1328.2)
				2013	25,253	(15,307 to 42,886)	820.8	(470.7 to 1503.6)
			Both	1990	30,311	(20,936 to 38,019)	886.6	(607.4 to 1132.0)
				2013	46,460	(32,224 to 72,862)	759.1	(491.1 to 1355.0)
	YLDs	Ischemic Stroke	Male	1990	189	(118 to 269)	15.2	(9.6 to 21.8)
				2013	350	(224 to 498)	15.2	(9.7 to 22.4)
			Female	1990	154	(98 to 224)	10.9	(6.9 to 16.1)
				2013	277	(181 to 383)	10.8	(7.1 to 15.2)
			Both	1990	344	(229 to 478)	12.9	(8.7 to 17.9)
				2013	628	(419 to 857)	12.8	(8.6 to 17.8)
		Hemorrhagic Stroke	Male	1990	61	(38 to 90)	4.2	(2.6 to 6.1)
		Stroke	Widic	2013	113	(73 to 160)	3.9	(2.5 to 5.7)
			Female	1990	43	(27 to 64)	2.7	(1.7 to 4.1)
			Terriale	2013	79	(49 to 116)	2.6	(1.6 to 3.7)
			Both	1990	105	(68 to 151)	3.4	(2.2 to 4.9)
			Both	2013	193	(128 to 268)	3.4	(2.1 to 4.4)
Chile	Deaths	Ischemic Stroke	Male	1990	2,559	(2,026 to 3,079)	81.8	(64.7 to 98.9)
Cilie	Deatils	ischeniic stroke	IVIAIC	2013	2,339	(2,246 to 3,758)	40.2	(31.1 to 52.1)
				2013	2,030	(2,240 to 3,730)	70.2	(51.1 (0 52.1)
			Female	1990	2,845	(2,045 to 3,195)	61.2	(44.1 to 68.7)

		7						1
				2013	3,176	(2,465 to 4,811)	28.9	(22.5 to 43.9)
			Both	1990	5,405	(4,304 to 6,095)	69.8	(55.9 to 78.4)
					Cou	ints, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	6,067	(5,014 to 8,191)	33.7	(28.0 to 45.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,319	(993 to 1,960)	36.9	(27.4 to 55.9)
				2013	2,563	(1,629 to 3,281)	32.7	(20.3 to 42.2)
			Female	1990	1,450	(1,216 to 2,198)	29.9	(25.0 to 45.5)
				2013	2,602	(1,753 to 3,254)	24.5	(16.6 to 30.5)
			Both	1990	2,770	(2,285 to 3,958)	32.9	(27.0 to 47.2)
				2013	5,166	(3,723 to 6,101)	28.2	(20.3 to 33.4)
	DALYs	Ischemic Stroke	Male	1990	44,116	(33,855 to 52,897)	1228.8	(954.4 to 1468.2)
				2013	40,609	(31,874 to 54,230)	522.9	(411.5 to 690.0)
			Female	1990	39,281	(27,426 to 43,804)	810.1	(567.0 to 902.3)
				2013	32,714	(25,988 to 49,787)	314.8	(250.1 to 482.3)
			Both	1990	83,398	(65,213 to 94,784)	988.9	(780.2 to 1121.2)
				2013	73,324	(60,610 to 98,472)	406.8	(336.9 to 544.6)
		Hemorrhagic						
		Stroke	Male	1990	31,161	(24,188 to 44,569)	736.4	(563.0 to 1070.9)
				2013	51,850	(34,701 to 64,665)	602.6	(396.7 to 753.3)
			_					
			Female	1990	29,714	(25,639 to 43,617)	574.6	(495.2 to 840.9)
				2013	41,652	(30,732 to 49,890)	411	(304.0 to 490.7)
			Both	1990	60,875	(51,251 to 85,035)	645.6	(539.7 to 905.9)
				2012	00 -00	(71,125 to	400.0	(270 7
				2013	93,503	109,059)	499.8	(378.7 to 583.0)
	YLDs	Ischemic Stroke	Male	1990	1,364	(902 to 1,913)	34.1	(22.4 to 47.1)
				2013	2,582	(1,669 to 3,597)	30.5	(19.8 to 42.6)

	_				1	1		
			Female	1990	1,007	(646 to 1,439)	19.9	(12.7 to 28.5)
				2013	1,837	(1,183 to 2,602)	18.3	(11.8 to 25.9)
			Both	1990	2,371	(1,613 to 3,219)	26.1	(17.7 to 35.6)
				2013	4,419	(2,948 to 5,929)	23.8	(15.9 to 31.9)
		Hemorrhagic						
		Stroke	Male	1990	514	(338 to 714)	11.4	(7.5 to 15.7)
				2013	1,133	(694 to 1,621)	13	(8.0 to 18.6)
			Female	1990	308	(198 to 446)	5.8	(3.7 to 8.4)
				2013	636	(390 to 931)	6.4	(3.9 to 9.3)
			Both	1990	823	(553 to 1,116)	8.3	(5.5 to 11.3)
			Both	2013	1,769	(1,129 to 2,440)	9.4	(6.0 to 13.0)
					,	(132,718 to		,
China	Deaths	Ischemic Stroke	Male	1990	169,156	219,389)	61.8	(48.4 to 80.7)
						(295,377 to		
				2013	414,113	511,239)	74.7	(53.6 to 92.1)
						(119,763 to		
			Female	1990	148,634	210,776)	43.7	(35.5 to 62.1)
						(212,678 to		
				2013	321,862	376,981)	51.9	(34.5 to 60.7)
								400,000
						nts, All ages		100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(273,999 to		(
			Both	1990	317,791	410,202)	51.6	(44.6 to 66.9)
				2012	725.076	(554,470 to	63.0	(47.1 + 2.72.1)
		Homorrhogic		2013	735,976	849,199)	62.9	(47.1 to 72.1)
	Deaths	Hemorrhagic Stroke	Male	1990	533,294	(446,725 to 601,019)	171.7	(141.1 to 193.3)
	Deatils	JUNE	IVIAIC	1990	333,234	(558,315 to	1/1./	(141.1 (0 133.3)
				2013	696,125	897,513)	114.2	(91.8 to 149.8)
			Female	1990	454,968	(394,067 to	127.9	(109.5 to 154.4)
			· ca.c	1000	.5.,500	(33.)307.60	± - /.5	(103.5 to 13 1.1)

					549,346)		
					(413,673 to		
			2013	488,586	634,569)	75.5	(64.1 to 98.9)
			2013	400,300	(869,268 to	75.5	(04.1 to 30.3)
		Both	1990	988,262	1,100,203)	148.1	(130.4 to 164.1)
					(1,024,961 to		
			2013	1,184,712	1,462,699)	94.4	(81.6 to 117.8)
					(2,481,710 to		
DALYs	Ischemic Stroke	Male	1990	3,206,648	4,143,927)	949.5	(748.7 to 1227.9)
					(4,811,217 to		
			2013	6,655,368	8,277,431)	1041	(758.8 to 1279.4)
					(1,913,613 to		
		Female	1990	2,415,466	3,403,162)	645.6	(515.3 to 906.1)
					(2,748,902 to		
			2013	3,997,838	4,801,774)	602.7	(413.3 to 719.5)
					(4,761,928 to		
		Both	1990	5,622,115	7,127,920)	786.9	(672.9 to 1000.4)
					(8,141,338 to		
			2013	10,653,207	12,546,385)	816.5	(619.6 to 954.6)
	Hemorrhagic				(10,629,562 to		
	Stroke	Male	1990	12,603,438	14,334,550)	3199.5	(2684.2 to 3610.3)
					(11,341,678 to		
			2013	14,122,661	17,652,785)	1991.6	(1612.5 to 2510.9)
					(7,811,045 to		
		Female	1990	9,095,120	11,219,763)	2268	(1951.4 to 2751.1)
					(6,615,153 to		
			2013	7,919,171	9,675,333)	1122.3	(942.3 to 1387.0)
					(19,249,450 to		
		Both	1990	21,698,558	24,229,403)	2721.6	(2402.2 to 3018.9)
					(19,127,173 to		
			2013	22,041,833	26,422,172)	1555.2	(1351.2 to 1880.9)
YLDs	Ischemic Stroke	Male	1990	106,328	(73,769 to	31.5	(22.0 to 41.9)

			1		1	T .	l I	
						142,238)		
						(148,523 to		
				2013	213,407	287,707)	32.4	(22.8 to 43.4)
						(53,145 to		
			Female	1990	75,481	100,282)	19.9	(14.0 to 26.4)
						(96,931 to		
				2013	138,263	183,697)	19.9	(13.9 to 26.4)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		,,		, , , , , , , , , , , , , , , , , , ,		(126,595 to		
			Both	1990	181,809	241,932)	25.2	(17.6 to 33.3)
					,	(246,176 to		,
				2013	351,670	466,293)	25.8	(18.2 to 34.2)
		Hemorrhagic			,	(76,025 to		,
		Stroke	Male	1990	112,782	151,680)	31.1	(21.1 to 41.7)
						(146,306 to		
				2013	216,225	288,504)	31.3	(21.4 to 41.8)
						(53,851 to		
			Female	1990	78,590	103,914)	19.9	(13.7 to 26.4)
						(96,479 to		
				2013	139,468	187,873)	19.6	(13.6 to 26.4)
						(130,030 to		
			Both	1990	191,372	255,309)	25	(17.0 to 33.4)
						(246,814 to		
			Both	2013	355,693	473,579)	25.3	(17.6 to 33.8)
Colombia	Deaths	Ischemic Stroke	Male	1990	1,973	(1,409 to 2,396)	33.5	(23.7 to 41.3)
				2013	2,147	(1,512 to 3,609)	17.3	(12.0 to 29.3)
						,		
			Female	1990	2,280	(1,530 to 2,641)	30.8	(20.8 to 35.7)
				2013	2,496	(1,785 to 4,142)	15	(10.7 to 24.9)
			Both	1990	4,253	(3,227 to 4,815)	32.1	(23.9 to 36.4)

				2013	4,643	(3,573 to 7,113)	16	(12.3 to 24.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,201	(3,525 to 5,229)	63.6	(52.8 to 80.9)
				2013	7,612	(5,713 to 9,860)	56.8	(42.5 to 73.4)
			Female	1990	4,923	(4,248 to 6,020)	60.3	(51.9 to 74.7)
				2013	8,516	(6,517 to 10,514)	48.9	(37.0 to 60.1)
			Both	1990	9,124	(8,279 to 10,758)	61.8	(55.7 to 73.4)
				2013	16,128	(13,121 to 19,035)	52.4	(42.6 to 61.8)
	DALYs	Ischemic Stroke	Male	1990	37,807	(27,685 to 45,645)	513.7	(371.0 to 619.0)
				2013	35,913	(25,536 to 57,746)	242	(172.6 to 393.3)
			Female	1990	37,628	(25,481 to 43,712)	431.6	(293.7 to 500.6)
				2013	33,795	(24,259 to 55,345)	185.1	(133.8 to 304.3)
			Both	1990	75,436	(58,368 to 85,191)	469	(365.4 to 529.7)
						(53,291 to		
				2013	69,708	104,293)	210.6	(161.3 to 317.4)
		Hemorrhagic				(86,926 to		
		Stroke	Male	1990	103,353	127,604)	1205.8	(1013.8 to 1493.4)
						(115,070 to		
				2013	153,826	199,768)	937.4	(707.0 to 1213.4)
						(95,401 to		()
			Female	1990	112,285	134,715)	1126.7	(967.0 to 1361.8)
				2042	446.006	(114,035 to	747.0	(575.0). 007.4)
				2013	146,886	184,826)	747.2	(575.3 to 937.1)
					Cou	nts, All ages	Rate ner	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Codifici	measare	otroke rype		year		(195,775 to	Mean	3370 31
			Both	1990	215,638	250,992)	1161.5	(1055.8 to 1360.2)
						(244,769 to		,
				2013	300,712	358,842)	832.8	(679.2 to 988.7)

	-							
	YLDs	Ischemic Stroke	Male	1990	505	(317 to 730)	6.9	(4.2 to 10.1)
				2013	1,148	(703 to 1,692)	7.2	(4.4 to 10.6)
			Female	1990	478	(312 to 705)	5.5	(3.6 to 8.2)
				2013	1,139	(722 to 1,680)	5.9	(3.7 to 8.7)
			Both	1990	983	(646 to 1,380)	6.1	(4.1 to 8.6)
				2013	2,287	(1,490 to 3,251)	6.5	(4.2 to 9.2)
		Hemorrhagic						
		Stroke	Male	1990	1,642	(1,104 to 2,255)	19.4	(12.9 to 26.6)
				2013	3,578	(2,346 to 5,157)	20.4	(13.6 to 29.1)
			Female	1990	1,248	(814 to 1,752)	12.8	(8.3 to 17.9)
				2013	2,605	(1,675 to 3,720)	12.5	(8.1 to 17.7)
			Both	1990	2,891	(1,967 to 3,848)	15.8	(10.7 to 20.9)
			Both	2013	6,183	(4,171 to 8,522)	16	(10.9 to 21.9)
Comoros	Deaths	Ischemic Stroke	Male	1990	58	(17 to 102)	111.1	(34.6 to 184.0)
				2013	90	(26 to 148)	99.7	(33.2 to 158.5)
			Female	1990	71	(22 to 119)	111.2	(34.9 to 180.0)
				2013	96	(33 to 153)	91.6	(33.0 to 144.4)
			Both	1990	129	(48 to 194)	111.7	(42.6 to 163.7)
				2013	186	(72 to 269)	95.6	(38.0 to 134.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	96	(58 to 152)	140.8	(84.9 to 224.4)
				2013	157	(93 to 249)	130.8	(77.0 to 208.6)
			Female	1990	116	(67 to 171)	148.5	(87.7 to 217.6)
				2013	152	(88 to 256)	118.8	(69.9 to 206.6)
			Both	1990	212	(144 to 293)	145.6	(99.3 to 201.5)
				2013	309	(206 to 471)	124.8	(83.8 to 198.1)
	DALYs	Ischemic Stroke	Male	1990	1,162	(326 to 2,103)	1720.3	(509.8 to 3011.6)
				2013	1,784	(517 to 3,047)	1515.8	(458.2 to 2484.9)
			Female	1990	1,223	(361 to 2,161)	1583.4	(487.2 to 2715.3)

2013 1,574 (531 to 2,641) 1260.5 (439.8 to 2,641) 1260.5 (616.0 to 2,641) (616.0 to 2,641)	2500.1) 2003.7) 0 4669.9) 0 4154.2)
2013 3,359 (1,274 to 5,003) 1384.6 (541.1 to Hemorrhagic Stroke Male 1990 2,621 (1,634 to 4,139) 2917.3 (1760.9 to 2013 4,194 (2,447 to 6,521) 2620.4 (1540.0 to Female 1990 2,766 (1,620 to 4,195) 2889.4 (1677.4 to	2003.7) 0 4669.9) 0 4154.2)
Hemorrhagic Stroke Male 1990 2,621 (1,634 to 4,139) 2917.3 (1760.9 to 2013 4,194 (2,447 to 6,521) 2620.4 (1540.0 to Female 1990 2,766 (1,620 to 4,195) 2889.4 (1677.4 to 2013)	o 4669.9) o 4154.2)
Stroke Male 1990 2,621 (1,634 to 4,139) 2917.3 (1760.9 to 4,194) 2013 4,194 (2,447 to 6,521) 2620.4 (1540.0 to 4,195) Female 1990 2,766 (1,620 to 4,195) 2889.4 (1677.4 to 4,195)	4154.2)
2013 4,194 (2,447 to 6,521) 2620.4 (1540.0 to Female 1990 2,766 (1,620 to 4,195) 2889.4 (1677.4 to	4154.2)
Female 1990 2,766 (1,620 to 4,195) 2889.4 (1677.4 to	•
	4303.0)
2013 3,445 (2,031 to 5,672) 2168.3 (1265.8 to	
	3598.7)
Both 1990 5,387 (3,673 to 7,425) 2913.8 (1974.8 to	4045.0)
Counts, All ages Rate per 100,000, Age-s	
Country measure Stroke Type sex year Mean 95% UI Mean 95%	UI
2013 7,640 (5,170 to 11,254) 2392 (1593.4 to	3612.6)
YLDs Ischemic Stroke Male 1990 9 (5 to 14) 10 (6.2 to	14.8)
2013 18 (11 to 27) 10.9 (7.2 to	16.2)
Female 1990 7 (4 to 10) 7.1 (4.5 to	10.2)
2013 14 (8 to 21) 7.7 (4.7 to	11.3)
Both 1990 16 (10 to 23) 8.4 (5.5 to	11.7)
2013 33 (21 to 47) 9.2 (6.0 to	13.2)
Hemorrhagic	
Stroke Male 1990 3 (1 to 4) 2.8 (1.7 to	4.1)
2013 6 (3 to 9) 2.9 (1.8 to	4.4)
Female 1990 2 (1 to 3) 1.8 (1.1 to	2.6)
2013 4 (2 to 6) 1.9 (1.1 to	2.8)
Both 1990 5 (3 to 7) 2.2 (1.5 to	3.2)
Botti 1330 3 (3.07) 2.2 (1.3.0)	
Both 2013 10 (6 to 14) 2.4 (1.5 to	3.4)
Both 2013 10 (6 to 14) 2.4 (1.5 to	185.5)

	1		1 1			T		Т
				2013	932	(682 to 1,219)	127.2	(94.1 to 165.4)
			Both	1990	1,218	(867 to 1,447)	150	(109.3 to 178.7)
				2013	1,813	(1,388 to 2,167)	130.7	(101.7 to 155.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	709	(545 to 942)	148.9	(113.3 to 201.2)
				2013	1,083	(784 to 1,460)	128.1	(93.2 to 172.8)
			Female	1990	823	(615 to 1,151)	160.9	(119.5 to 231.4)
				2013	1,030	(710 to 1,561)	118.3	(80.3 to 181.5)
			Both	1990	1,533	(1,269 to 1,944)	156	(128.1 to 202.9)
				2013	2,113	(1,652 to 2,801)	123.5	(95.4 to 165.5)
	DALYs	Ischemic Stroke	Male	1990	11,399	(7,490 to 14,978)	2432	(1646.7 to 3088.2)
				2013	16,958	(11,346 to 22,897)	2080.2	(1446.2 to 2724.5)
			Female	1990	10,839	(6,223 to 14,744)	2151.7	(1331.4 to 2807.9)
				2013	14,692	(10,288 to 19,813)	1727.1	(1240.0 to 2291.1)
			Both	1990	22,238	(14,950 to 27,115)	2291.1	(1614.9 to 2740.6)
				2013	31,651	(23,725 to 39,117)	1901	(1443.2 to 2295.5)
		Hemorrhagic						
		Stroke	Male	1990	19,484	(14,937 to 25,480)	3299.3	(2507.3 to 4378.0)
				2013	30,156	(21,907 to 40,480)	2762.9	(1987.3 to 3726.4)
			Female	1990	19,718	(14,165 to 26,947)	3219	(2354.0 to 4467.6)
				2013	23,769	(16,242 to 36,010)	2228.7	(1530.3 to 3308.2)
			Both	1990	39,202	(32,280 to 48,751)	3268.2	(2696.0 to 4104.9)
				2013	53,925	(42,583 to 70,213)	2495.6	(1958.2 to 3301.9)
	YLDs	Ischemic Stroke	Male	1990	81	(53 to 114)	14.7	(9.6 to 21.1)
				2013	142	(94 to 202)	13.8	(9.1 to 20.1)
			Female	1990	68	(45 to 97)	11.2	(7.3 to 15.6)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	123	(81 to 179)	11.1	(7.3 to 16.0)

1		1		ı	1		
		Both	1990	150	(101 to 205)	12.8	(8.7 to 17.6)
			2013	266	(180 to 368)	12.4	(8.4 to 17.2)
	Hemorrhagic						
	Stroke	Male					(2.4 to 5.7)
			2013	45	(29 to 69)	3.6	(2.3 to 5.5)
		Female	1990	18	(11 to 27)	2.7	(1.7 to 4.0)
			2013	32	(20 to 48)	2.5	(1.6 to 3.7)
		Both	1990	43	(28 to 60)	3.3	(2.1 to 4.6)
		Both	2013	78	(50 to 112)	3.1	(2.0 to 4.4)
Deaths	Ischemic Stroke	Male	1990	261	(165 to 336)	38.4	(24.3 to 49.8)
			2013	243	(153 to 444)	15.2	(9.6 to 27.6)
		Female	1990	305	(185 to 356)	39.3	(23.8 to 45.9)
			2013	261	(175 to 492)	13.3	(8.9 to 25.0)
		Both	1990	566	(375 to 676)	39	(25.8 to 46.6)
			2013	504	(346 to 856)	14.2	(9.7 to 24.1)
	Hemorrhagic						
Deaths	Stroke	Male	1990	152	(105 to 257)	20.6	(13.9 to 36.4)
			2013	463	(260 to 604)	27.6	(15.2 to 36.2)
		Female	1990	178	(139 to 310)	21.8	(16.7 to 39.0)
			2013	515	(307 to 625)	25.9	(15.3 to 31.4)
		Both	1990	330	(252 to 539)	21.3	(15.9 to 35.7)
			2013	979	(623 to 1,181)	26.7	(16.7 to 32.3)
DALYs	Ischemic Stroke	Male	1990	4,004	(2,509 to 5,081)	506.9	(319.9 to 643.8)
			2013	3,350	(2,223 to 5,942)	188.4	(124.7 to 333.1)
		Female	1990	4,089	(2,409 to 4,739)	482.3	(286.8 to 559.6)
			2013	2,877	(2,016 to 5,374)	143.8	(100.6 to 267.7)
		Both	1990	8,093	(5,300 to 9,704)	494.6	(326.2 to 590.9)
			2013	6,228	(4,456 to 10,611)	165	(118.0 to 280.3)
	Hemorrhagic	Male	1990	3,339	(2,505 to 5,133)	370.9	(271.8 to 582.4)
	Deaths	Deaths Ischemic Stroke Hemorrhagic Stroke DALYS Ischemic Stroke	Hemorrhagic Stroke Male Female Both Both Both Deaths Ischemic Stroke Hemorrhagic Stroke Both Female Both Both Both Both Both Both Both Female Female Female Female Female Both Female Both Female Both Both Both Both	Hemorrhagic Stroke Male 1990 2013	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 24 (15 to 37)	Hemorrhagic Stroke Male 1990 24 (15 to 37) 3.8

	Stroke						
			2013	8,099	(5,160 to 10,273)	416.9	(261.0 to 533.0)
		Female	1990	3,510	(2,962 to 5,417)	375.6	(312.6 to 592.7)
			2013	7,280	(4,993 to 8,457)	348	(235.1 to 404.3)
		Both	1990	6,849	(5,628 to 9,970)	373.2	(300.8 to 558.8)
			2013	15,379	(10,973 to 18,112)	381.3	(267.1 to 452.3)
YLDs	Ischemic Stroke	Male	1990	59	(35 to 87)	7.2	(4.4 to 10.9)
			2013	133	(75 to 200)	7	(4.0 to 10.5)
		Female	1990	48	(30 to 71)	5.7	(3.6 to 8.4)
			2013	119	(72 to 180)	5.8	(3.5 to 8.9)
		Both	1990	107	(70 to 153)	6.5	(4.2 to 9.2)
							(4.1 to 9.0)
			2013	253	(164 to 354)	6.4	
				Cou	nts, All ages	Rate per	100,000, Age-standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
	Hemorrhagic						
	Stroke	Male			,		(13.4 to 28.2)
			2013	431	(284 to 614)	20.8	(13.6 to 29.3)
		Female	1990	118	(78 to 167)	12.8	(8.4 to 18.0)
			2013	275	(177 to 393)	12.8	(8.2 to 18.4)
		Both	1990	300	(203 to 413)	16.4	(11.0 to 22.5)
		Both	2013	707	(477 to 949)	16.7	(11.2 to 22.4)
Deaths	Ischemic Stroke	Male	1990	1.761	(1,305 to 2,335)	96	(72.7 to 121.6)
				, -	, , ,		
			2013	3,994	(2,171 to 5,428)	121.8	(72.2 to 160.5)
		Female		-			(72.2 to 160.5) (59.8 to 128.0)
			2013	3,994	(2,171 to 5,428)	121.8	· · ·
			2013 1990	3,994 1,347	(2,171 to 5,428) (873 to 1,881)	121.8 91.3	(59.8 to 128.0)
		Female	2013 1990 2013	3,994 1,347 3,068	(2,171 to 5,428) (873 to 1,881) (2,041 to 4,111)	121.8 91.3 123.4	(59.8 to 128.0) (85.5 to 161.6)
	measure	YLDs Ischemic Stroke Measure Stroke Type Hemorrhagic Stroke	Female Both YLDs Ischemic Stroke Male Female Both Both Both Male Female Both Female Both Female Both Both Both Both Both Both	2013 Female 1990 2013	Stroke Type Sex Year Mean	Stroke Type Sex Year Mean 95% U Hemorrhagic Stroke Male 1990 118 (78 to 167) (77 to 393) (77 to 393) (77 to 394) (77 to 949) (77 to 194) (77 to 949) (77 to 949) (77 to 1949) (77 to 1949) (77 to 949) (77 to 949	Second

1				1		·	
	Stroke						
			2013	2,665	(1,661 to 4,859)	59.8	(34.7 to 116.8)
		Female	1990	1,325	(893 to 1,776)	66.8	(45.5 to 89.9)
			2013	1,917	(1,256 to 3,370)	53.9	(33.8 to 104.0)
		Both	1990	3,199	(2,605 to 3,893)	70.9	(57.2 to 86.2)
			2013	4,583	(3,266 to 7,510)	57.1	(38.7 to 101.8)
DALYs	Ischemic Stroke	Male	1990	41,788	(29,631 to 59,549)	1622.6	(1210.5 to 2141.4)
					(44,263 to		
			2013	86,966	120,790)	1990.7	(1095.0 to 2697.7)
		Female	1990	27,691	(17,225 to 39,496)	1367.2	(892.2 to 1902.8)
			2013	59,847	(39,357 to 82,343)	1791.4	(1191.2 to 2397.0)
		Both	1990	69,480	(52,933 to 91,627)	1509.8	(1193.7 to 1919.8)
					(94,299 to		
			2013	146,813	187,848)	1900.6	(1292.1 to 2353.7)
	Hemorrhagic						
	Stroke	Male	1990	63,074	(45,536 to 79,673)	1668.1	(1190.1 to 2149.3)
			2010	0.1.0.10	(56,342 to	10000	(051.0)
			2013	84,343	134,503)	1363.9	(861.0 to 2456.5)
		Female	1990	41,108	(26,447 to 54,985)	1353.9	(919.2 to 1816.9)
			2013	58,309	(39,548 to 96,556)	1127.3	(741.2 to 1975.8)
					(82,544 to		
		Both	1990	104,182	126,648)	1527.1	(1247.0 to 1860.3)
			2012	442.653	(107,414 to	1252.6	(005.2 ±= 2020.0)
			2013	142,653	214,309)	1252.6	(895.3 to 2030.8)
YLDs	Ischemic Stroke	Male	1990	463	(299 to 659)	15.4	(10.0 to 21.6)
			2013	777	(486 to 1,159)	14.8	(9.1 to 22.4)
		Female	1990	292	(190 to 407)	11.6	(7.8 to 15.9)
			2013	533	(346 to 767)	11.6	(7.6 to 16.9)
		Both	1990	755	(505 to 1,044)	13.6	(9.3 to 18.5)
			2013	1,310	(857 to 1,860)	13.4	(8.7 to 19.1)

					Cou	ints, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	154	(95 to 224)	4.3	(2.7 to 6.2)
				2013	232	(148 to 346)	3.8	(2.4 to 5.7)
			Female	1990	85	(54 to 125)	2.9	(1.9 to 4.2)
				2013	147	(93 to 218)	2.7	(1.7 to 4.0)
			Both	1990	240	(159 to 334)	3.7	(2.4 to 5.0)
			Both	2013	380	(257 to 537)	3.3	(2.2 to 4.7)
Croatia	Deaths	Ischemic Stroke	Male	1990	2,933	(2,541 to 3,408)	157.3	(136.0 to 182.3)
				2013	2,718	(2,230 to 3,145)	91.2	(75.0 to 105.6)
			Female	1990	4,046	(3,451 to 4,773)	118.8	(100.6 to 139.6)
				2013	3,882	(3,229 to 4,622)	69	(57.4 to 81.8)
			Both	1990	6,980	(6,240 to 7,852)	133.3	(119.3 to 150.1)
				2013	6,601	(5,794 to 7,419)	78.1	(68.5 to 87.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,172	(934 to 1,446)	49.4	(39.1 to 61.8)
				2013	865	(638 to 1,097)	28.7	(21.1 to 36.6)
			Female	1990	1,442	(1,057 to 1,703)	41.5	(30.8 to 49.3)
				2013	991	(776 to 1,456)	21.6	(17.6 to 31.1)
			Both	1990	2,614	(2,130 to 2,986)	45.2	(36.9 to 51.6)
				2013	1,857	(1,578 to 2,320)	24.9	(21.3 to 30.9)
	DALYs	Ischemic Stroke	Male	1990	43,859	(37,568 to 51,340)	2116.7	(1828.0 to 2466.0)
				2013	34,899	(28,803 to 40,196)	1133.7	(934.9 to 1308.9)
			Female	1990	46,537	(39,598 to 54,502)	1365.6	(1163.9 to 1584.2)
				2013	35,759	(30,235 to 42,779)	684.7	(573.2 to 823.5)
						(80,774 to		
			Both	1990	90,396	101,705)	1664.2	(1492.4 to 1871.0)
				2013	70,658	(62,331 to 79,434)	874.4	(765.7 to 984.7)

(931.3 to 1406.1)
(465.7 to 751.5)
(676.4 to 1036.0)
(344.0 to 569.6)
(849.4 to 1144.3)
(438.0 to 613.3)
(16.1 to 37.1)
(17.1 to 38.0)
(10.3 to 22.7)
(9.8 to 22.4)
(13.2 to 28.0)
(13.7 to 28.1)
(3.3 to 8.1)
(3.4 to 8.3)
000, Age-standardized
95% UI
(2.0 to 5.4)
(2.4 to 6.0)
(2.8 to 6.4)
(3.1 to 6.8)
(56.4 to 85.5)
(35.8 to 63.2)
(35.8 to 63.2) (56.4 to 83.9)
•
(56.4 to 83.9)
(56.4 to 83.9) (30.1 to 55.2)

	Stroke						
			2013	1,891	(1,446 to 2,602)	26.8	(20.5 to 37.0)
		Female	1990	1,426	(1,173 to 1,963)	29.3	(24.2 to 40.1)
			2013	1,980	(1,440 to 2,801)	23.9	(17.3 to 33.8)
		Both	1990	2,655	(2,304 to 3,335)	28.4	(24.7 to 35.7)
			2013	3,871	(3,132 to 5,131)	25.4	(20.6 to 33.6)
DALYs	Ischemic Stroke	Male	1990	46,172	(37,424 to 56,010)	1037.2	(842.5 to 1258.1)
			2013	47,306	(33,578 to 58,733)	676.6	(480.8 to 838.9)
		Female	1990	44,965	(37,009 to 54,301)	917.7	(751.4 to 1106.0)
			2013	43,514	(28,943 to 51,647)	525.5	(347.8 to 623.3)
					(79,538 to		
		Both	1990	91,137	104,081)	975.8	(850.5 to 1113.7)
					(66,453 to		
			2013	90,821	105,712)	598.6	(435.5 to 697.3)
	Hemorrhagic						
	Stroke	Male	1990	27,528	(22,449 to 35,655)	599.4	(489.0 to 779.3)
			2013	38,746	(30,804 to 50,492)	538.3	(428.7 to 702.0)
		Female	1990	29,842	(25,488 to 39,691)	623.9	(532.8 to 830.0)
			2013	33,091	(24,772 to 44,293)	425.8	(319.0 to 570.2)
		Both	1990	57,371	(51,028 to 70,708)	611.7	(543.7 to 754.0)
			2013	71,838	(60,468 to 88,476)	481.7	(405.5 to 593.6)
YLDs	Ischemic Stroke	Male	1990	2,022	(1,295 to 2,904)	45.5	(29.0 to 65.4)
			2013	3,009	(2,029 to 4,251)	42.9	(28.8 to 60.7)
		Female	1990	1,540	(980 to 2,222)	31.6	(20.2 to 45.5)
			2013	2,598	(1,662 to 3,807)	32.5	(21.0 to 47.2)
		Both	1990	3,563	(2,408 to 4,956)	38.2	(25.9 to 53.2)
			2013	5,608	(3,774 to 7,742)	37.4	(25.4 to 51.7)
	Hemorrhagic						
	Stroke	Male	1990	624	(388 to 878)	13.7	(8.5 to 19.4)

						1		
				2013	950	(564 to 1,471)	13.6	(8.1 to 21.0)
			Female	1990	487	(307 to 698)	10	(6.3 to 14.3)
				2013	819	(510 to 1,240)	10.6	(6.6 to 15.9)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	1,112	(729 to 1,513)	11.8	(7.7 to 16.0)
			Both	2013	1,769	(1,162 to 2,520)	12	(7.9 to 17.1)
Cyprus	Deaths	Ischemic Stroke	Male	1990	195	(124 to 248)	63.9	(40.9 to 81.1)
				2013	136	(104 to 177)	24.7	(19.0 to 32.1)
			Female	1990	252	(146 to 321)	64.3	(37.6 to 81.6)
				2013	196	(154 to 272)	24.8	(19.7 to 34.3)
			Both	1990	448	(298 to 541)	64.4	(43.2 to 77.7)
				2013	333	(283 to 415)	25	(21.3 to 31.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	156	(106 to 200)	50.9	(34.7 to 65.3)
				2013	103	(79 to 133)	19.1	(14.7 to 24.6)
			Female	1990	167	(91 to 220)	44.7	(25.0 to 58.1)
				2013	123	(96 to 180)	16.9	(13.5 to 24.3)
			Both	1990	323	(219 to 397)	47.7	(32.6 to 58.1)
				2013	227	(191 to 294)	18.1	(15.3 to 23.3)
	DALYs	Ischemic Stroke	Male	1990	2,478	(1,706 to 3,018)	813.2	(561.1 to 991.6)
				2013	1,867	(1,498 to 2,324)	343.7	(276.8 to 427.4)
			Female	1990	2,528	(1,589 to 3,111)	681.6	(432.1 to 838.1)
				2013	1,873	(1,552 to 2,466)	265.9	(220.8 to 347.0)
			Both	1990	5,006	(3,554 to 5,853)	743.9	(528.5 to 870.1)
				2013	3,741	(3,225 to 4,473)	302.6	(260.3 to 361.9)
		Hemorrhagic						
		Stroke	Male	1990	2,894	(2,097 to 3,536)	934.8	(679.0 to 1141.2)
				2013	1,701	(1,370 to 2,133)	332.7	(268.9 to 414.5)

			Female	1990	2,618	(1,677 to 3,180)	752.7	(485.9 to 907.9)
				2013	1,513	(1,246 to 2,050)	252.7	(208.8 to 345.3)
			Both	1990	5,512	(4,026 to 6,448)	839.1	(612.5 to 980.7)
				2013	3,214	(2,780 to 3,938)	290.7	(252.4 to 354.0)
	YLDs	Ischemic Stroke	Male	1990	271	(178 to 374)	88.6	(58.2 to 122.4)
				2013	473	(286 to 682)	87.6	(53.4 to 126.1)
			Female	1990	204	(134 to 287)	58.1	(38.1 to 80.3)
				2013	384	(246 to 560)	60.3	(38.9 to 87.9)
			Both	1990	476	(324 to 649)	72.3	(49.4 to 98.4)
				2013	857	(577 to 1,188)	73.1	(49.1 to 101.8)
		Hemorrhagic						
		Stroke	Male	1990	60	(34 to 93)	19.5	(11.0 to 30.1)
				2013	101	(54 to 191)	19.1	(10.3 to 35.6)
			Female	1990	44	(22 to 70)	12.6	(6.4 to 20.2)
				2013	77	(40 to 133)	12.7	(6.9 to 21.3)
			Both	1990	104	(62 to 154)	15.8	(9.5 to 23.6)
			Both	2013	179	(105 to 278)	15.7	(9.3 to 24.3)
Czech Republic	Deaths	Ischemic Stroke	Male	1990	9,026	(7,937 to 10,577)	214.5	(188.8 to 247.7)
				2013	6,097	(5,158 to 7,256)	93.2	(78.9 to 111.1)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	13,046	(10,451 to 14,340)	161.6	(129.1 to 177.5)
				2013	8,533	(7,179 to 11,950)	71.8	(60.5 to 99.9)
			Both	1990	22,073	(19,579 to 24,075)	181.6	(160.9 to 197.7)
				2013	14,631	(12,787 to 18,635)	80.5	(70.5 to 102.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,036	(1,514 to 2,622)	40.9	(30.3 to 52.7)
				2013	1,345	(875 to 1,769)	19.1	(12.3 to 25.2)
			Female	1990	2,708	(1,883 to 3,218)	35.3	(24.6 to 42.0)

			2013	1,505	(1,151 to 2,143)	14.2	(11.0 to 20.1)
		Both	1990	4,745	(3,770 to 5,495)	38	(30.4 to 44.0)
			2013	2,850	(2,243 to 3,613)	16.4	(13.0 to 20.6)
					(113,373 to		
DALYs	Ischemic Stroke	Male	1990	130,413	157,291)	2840.6	(2495.5 to 3365.1)
			2013	75,583	(63,912 to 89,663)	1101.8	(934.5 to 1303.7)
					(116,064 to		
		Female	1990	146,744	161,535)	1862.6	(1452.5 to 2052.8)
					(65,705 to		
			2013	77,797	106,964)	708.8	(598.0 to 973.5)
			1000		(247,252 to		(2000 0 : 2404 4)
		Both	1990	277,157	305,296)	2253.6	(2008.0 to 2481.4)
			2012	152 201	(134,851 to	076.0	(771 4 to 1000 C)
	Hemorrhagic		2013	153,381	190,095)	876.9	(771.4 to 1080.6)
	Stroke	Male	1990	46,472	(35,048 to 60,385)	873.4	(664.6 to 1132.3)
	Stroke	iviale	2013	26,104	(17,754 to 33,060)	360.7	(244.0 to 456.1)
		Famala	1990	48,507	(34,092 to 54,926)	677.5	(476.6 to 766.4)
		Female		-	, , , , , , , , , , , , , , , , , , , ,		,
			2013	21,733	(17,561 to 30,056) (76,608 to	237.7	(194.6 to 323.0)
		Both	1990	94,980	108,996)	768.9	(622.6 to 880.6)
		Botti	2013	·	. ,	295.1	,
VI D-	lashawia Chualia	0.4-1-		47,837	(38,852 to 58,442)		(238.7 to 358.0)
YLDs	Ischemic Stroke	Male	1990	1,207	(768 to 1,743)	23.3	(14.9 to 33.9)
		_	2013	1,803	(1,187 to 2,534)	25.2	(16.7 to 35.5)
		Female	1990	1,070	(670 to 1,548)	15.5	(9.8 to 22.4)
			2013	1,266	(846 to 1,786)	14.6	(9.6 to 20.6)
		Both	1990	2,278	(1,528 to 3,202)	18.9	(12.6 to 26.5)
			2013	3,069	(2,085 to 4,236)	19.5	(13.3 to 26.9)
	Hemorrhagic						
	Stroke	Male	1990	246	(147 to 380)	4.7	(2.8 to 7.2)

				2013	399	(242 to 603)	5.8	(3.5 to 8.7)
			Female	1990	229	(131 to 361)	3.6	(2.1 to 5.6)
				2013	305	(183 to 468)	3.9	(2.4 to 6.0)
			Both	1990	476	(304 to 697)	4.1	(2.6 to 6.0)
			Both	2013	704	(461 to 1,012)	4.8	(3.1 to 6.9)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Democratic Republic of the								
Congo	Deaths	Ischemic Stroke	Male	1990	3,016	(1,394 to 4,547)	74.7	(32.2 to 110.7)
				2013	6,855	(2,162 to 10,125)	86.7	(27.8 to 124.6)
			Female	1990	5,422	(3,432 to 7,358)	105.2	(65.1 to 141.8)
				2013	8,643	(3,936 to 12,557)	93.8	(42.0 to 135.2)
			Both	1990	8,438	(5,394 to 10,919)	93.5	(60.4 to 122.6)
				2013	15,499	(7,495 to 21,141)	91.4	(44.4 to 125.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,687	(3,139 to 6,230)	78.9	(44.9 to 114.4)
				2013	11,054	(7,937 to 15,282)	99.9	(69.8 to 147.5)
			Female	1990	7,683	(5,148 to 10,761)	111.7	(72.0 to 159.6)
				2013	12,733	(8,284 to 19,259)	106	(67.7 to 166.4)
			Both	1990	12,371	(9,018 to 15,702)	98.4	(66.4 to 132.3)
				2013	23,788	(17,939 to 31,599)	103.9	(75.4 to 145.0)
	DALYs	Ischemic Stroke	Male	1990	69,694	(35,579 to 103,009)	1164.3	(529.8 to 1765.8)
				2013	147,726	(47,763 to 229,899)	1357.8	(430.3 to 1995.4)
			Female	1990	101,869	(63,109 to 145,536)	1503.1	(951.2 to 2048.7)

			1		1	/ . .	1	
						(73,162 to		/ · · · · · · · · · · · · · · · · ·
				2013	153,483	228,224)	1306.8	(603.9 to 1895.8)
						(112,125 to		
			Both	1990	171,564	223,633)	1370.2	(873.5 to 1773.7)
						(144,768 to		
				2013	301,210	414,728)	1339.2	(648.6 to 1831.2)
		Hemorrhagic				(116,465 to		
		Stroke	Male	1990	156,150	202,811)	1749.1	(1153.1 to 2334.0)
						(242,686 to		
				2013	338,272	454,253)	2164.3	(1550.7 to 2996.3)
						(144,420 to		
			Female	1990	214,784	299,322)	2288.3	(1539.9 to 3195.5)
						(224,755 to		
				2013	337,142	506,084)	2085.3	(1367.8 to 3139.5)
						(279,198 to		
			Both	1990	370,934	460,551)	2059.5	(1505.9 to 2601.0)
						(519,337 to		
				2013	675,415	864,066)	2132.5	(1611.8 to 2815.2)
	YLDs	Ischemic Stroke	Male	1990	1,018	(640 to 1,468)	14.2	(9.1 to 20.1)
				2013	1,946	(1,228 to 2,789)	14	(8.8 to 19.5)
			Female	1990	892	(550 to 1,256)	10.5	(6.6 to 14.9)
				2013	1,690	(1,079 to 2,472)	10.9	(7.0 to 16.2)
			Both	1990	1,910	(1,247 to 2,655)	12.1	(7.9 to 16.8)
				2013	3,637	(2,388 to 5,056)	12.3	(8.1 to 17.3)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		Hemorrhagic						
		Stroke	Male	1990	322	(192 to 493)	3.7	(2.2 to 5.8)
				2013	647	(417 to 970)	3.8	(2.5 to 5.6)
			Female	1990	253	(154 to 379)	2.6	(1.6 to 3.9)

				2013	460	(293 to 664)	2.5	(1.6 to 3.7)
			Both	1990	575	(364 to 830)	3.1	(2.0 to 4.4)
			Both	2013	1,108	(755 to 1,570)	3.1	(2.1 to 4.4)
Denmark	Deaths	Ischemic Stroke	Male	1990	1,495	(1,176 to 2,002)	48.4	(38.1 to 65.0)
				2013	1,251	(757 to 1,561)	30.7	(18.7 to 38.3)
			Female	1990	2,224	(1,707 to 2,725)	41.7	(32.1 to 51.6)
				2013	1,594	(1,214 to 2,076)	25.9	(19.6 to 33.4)
			Both	1990	3,720	(3,204 to 4,511)	44.8	(38.7 to 54.5)
				2013	2,845	(2,190 to 3,370)	28.2	(21.6 to 33.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,099	(825 to 1,469)	35.6	(26.6 to 47.7)
				2013	1,035	(715 to 1,303)	25.2	(17.6 to 31.7)
			Female	1990	1,565	(1,080 to 1,872)	33.4	(23.8 to 39.6)
				2013	1,164	(874 to 1,652)	20.6	(15.6 to 28.8)
			Both	1990	2,665	(2,094 to 3,178)	34.5	(27.6 to 41.0)
				2013	2,199	(1,786 to 2,722)	22.7	(18.5 to 27.8)
	DALYs	Ischemic Stroke	Male	1990	22,768	(18,684 to 30,312)	726.7	(595.9 to 965.2)
				2013	19,088	(12,793 to 23,300)	458.8	(307.6 to 559.5)
			Female	1990	24,784	(20,114 to 30,311)	524.7	(431.2 to 651.3)
				2013	17,026	(13,814 to 20,945)	313	(257.5 to 382.7)
			Both	1990	47,552	(41,883 to 57,546)	613.6	(540.3 to 744.6)
				2013	36,114	(28,750 to 41,824)	380.9	(302.2 to 441.0)
		Hemorrhagic						
		Stroke	Male	1990	19,363	(15,251 to 26,116)	631.4	(499.2 to 851.7)
				2013	16,233	(11,096 to 20,066)	398	(275.2 to 488.5)
			Female	1990	22,045	(16,494 to 26,261)	564.1	(435.7 to 683.8)
				2013	14,409	(11,261 to 19,198)	300.9	(233.0 to 394.5)
			Both	1990	41,408	(34,734 to 48,490)	593.6	(501.1 to 695.3)
				2013	30,643	(24,923 to 35,730)	346.3	(278.8 to 400.1)

	YLDs	Ischemic Stroke	Male	1990	3,542	(2,414 to 4,759)	113.7	(77.5 to 152.6)
				2013	4,724	(3,072 to 6,824)	113.5	(73.9 to 163.3)
			Female	1990	3,230	(2,183 to 4,519)	78.1	(52.7 to 109.6)
				2013	3,708	(2,384 to 5,290)	76.4	(49.8 to 107.7)
			Both	1990	6,773	(4,724 to 9,158)	93.5	(64.8 to 126.0)
				2013	8,432	(5,786 to 11,341)	94	(64.3 to 126.3)
		Hemorrhagic						
		Stroke	Male	1990	212	(130 to 308)	7	(4.3 to 10.1)
				2013	349	(209 to 522)	8.8	(5.3 to 13.1)
			Female	1990	204	(129 to 298)	5.5	(3.5 to 7.9)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	305	(187 to 452)	6.9	(4.3 to 10.1)
			Both	1990	417	(275 to 581)	6.1	(4.0 to 8.5)
			Both	2013	655	(410 to 934)	7.8	(4.9 to 11.0)
Djibouti	Deaths	Ischemic Stroke	Male	1990	66	(31 to 110)	105.8	(52.7 to 167.3)
				2013	122	(68 to 199)	85.3	(48.4 to 136.0)
			Female	1990	101	(56 to 158)	136.8	(80.1 to 202.7)
				2013	181	(110 to 296)	111	(70.4 to 177.7)
			Both	1990	168	(106 to 258)	123.4	(82.1 to 179.8)
				2013	303	(204 to 476)	99.5	(68.6 to 150.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	81	(45 to 128)	92	(48.5 to 152.2)
				2013	129	(82 to 204)	71.5	(44.8 to 112.0)
			Female	1990	118	(55 to 190)	119.5	(57.7 to 190.0)
				2013	177	(102 to 296)	90.3	(51.0 to 144.5)
			Both	1990	199	(116 to 301)	107.1	(62.2 to 160.6)
				2013	307	(204 to 475)	81.7	(54.9 to 121.7)
	DALYs	Ischemic Stroke	Male	1990	1,353	(620 to 2,355)	1569	(751.0 to 2613.6)

				2013	2,232	(1,204 to 3,770)	1248.1	(698.9 to 2043.7)
			Female	1990	1,878	(990 to 3,113)	1972.1	(1087.6 to 3072.5)
				2013	2,977	(1,744 to 5,200)	1546.3	(933.3 to 2599.2)
			Both	1990	3,232	(1,945 to 5,239)	1794.9	(1137.5 to 2762.7)
				2013	5,209	(3,357 to 8,524)	1411.6	(942.3 to 2243.3)
		Hemorrhagic						
		Stroke	Male	1990	2,373	(1,422 to 3,707)	1875.2	(1053.7 to 2973.5)
				2013	3,293	(2,110 to 5,236)	1403.1	(889.7 to 2203.3)
			Female	1990	3,151	(1,449 to 5,140)	2419	(1143.3 to 3881.5)
				2013	4,074	(2,288 to 6,900)	1706.4	(972.3 to 2864.2)
			Both	1990	5,524	(3,217 to 8,348)	2167	(1255.6 to 3285.2)
				2013	7,368	(4,807 to 11,579)	1564.5	(1033.6 to 2442.8)
	YLDs	Ischemic Stroke	Male	1990	13	(8 to 19)	10.3	(6.6 to 15.1)
				2013	25	(15 to 41)	10.4	(6.4 to 16.2)
			Female	1990	9	(5 to 14)	6.9	(4.3 to 10.1)
				2013	20	(12 to 30)	7.8	(5.0 to 11.4)
			Both	1990	22	(14 to 31)	8.5	(5.6 to 11.8)
				2013	46	(29 to 67)	9.1	(5.8 to 12.9)
		Hemorrhagic						
		Stroke	Male	1990	4	(2 to 6)	2.8	(1.7 to 4.3)
				2013	8	(5 to 13)	2.9	(1.8 to 4.5)
			Female	1990	2	(1 to 4)	1.7	(1.0 to 2.6)
				2013	5	(3 to 8)	1.9	(1.2 to 2.7)
			Both	1990	7	(4 to 10)	2.2	(1.4 to 3.2)
			Both	2013	14	(9 to 20)	2.4	(1.5 to 3.5)
Dominica	Deaths	Ischemic Stroke	Male	1990	16	(12 to 19)	79.9	(64.3 to 98.5)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	18	(13 to 24)	65.7	(46.8 to 88.1)

		Female	1990	20	(17 to 26)	90.5	(74.2 to 113.7)
			2013	20	(13 to 27)	57.1	(38.7 to 76.6)
		Both	1990	36	(32 to 43)	85.8	(74.7 to 101.2)
			2013	38	(29 to 47)	61.6	(47.4 to 76.2)
	Hemorrhagic						
Deaths	Stroke	Male	1990	9	(7 to 12)	44.7	(34.5 to 57.1)
			2013	11	(7 to 16)	39.7	(26.2 to 56.0)
		Female	1990	14	(11 to 17)	60.3	(48.3 to 76.1)
			2013	11	(8 to 15)	33.4	(23.5 to 45.1)
		Both	1990	23	(20 to 27)	53.2	(44.8 to 63.0)
			2013	23	(18 to 29)	36.6	(28.5 to 47.0)
DALYs	Ischemic Stroke	Male	1990	285	(229 to 348)	1307.2	(1051.8 to 1592.7)
			2013	315	(224 to 434)	1063.3	(753.9 to 1455.5)
		Female	1990	324	(264 to 400)	1354.2	(1099.1 to 1679.0)
			2013	246	(171 to 338)	715.3	(498.6 to 982.0)
		Both	1990	610	(530 to 713)	1335.4	(1159.6 to 1565.3)
			2013	562	(445 to 706)	883.1	(699.5 to 1111.8)
	Hemorrhagic						
	Stroke	Male	1990	225	(178 to 280)	930	(735.6 to 1157.6)
			2013	257	(169 to 362)	819.4	(541.3 to 1151.1)
		Female	1990	285	(228 to 348)	1128.1	(906.3 to 1378.1)
			2013	192	(135 to 261)	554.8	(389.7 to 756.5)
		Both	1990	510	(437 to 594)	1035.7	(880.7 to 1206.1)
			2013	450	(338 to 576)	683.8	(514.4 to 874.9)
YLDs	Ischemic Stroke	Male	1990	9	(5 to 13)	42.2	(26.1 to 62.5)
			2013	12	(8 to 18)	43.3	(28.8 to 62.6)
		Female	1990	7	(4 to 10)	32.1	(20.3 to 46.0)
			2013	10	(7 to 14)	31.3	(21.3 to 43.3)
		Both	1990	16	(11 to 23)	36.9	(24.3 to 51.5)

				2013	23	(16 to 32)	36.9	(25.2 to 50.4)
		Hemorrhagic						
		Stroke	Male	1990	3	(1 to 4)	13.5	(8.0 to 21.4)
				2013	4	(2 to 6)	13.5	(7.8 to 20.1)
			Female	1990	2	(1 to 3)	10.1	(6.2 to 15.1)
				2013	3	(2 to 5)	11.1	(7.0 to 16.7)
			Both	1990	5	(3 to 8)	11.7	(7.6 to 17.2)
			Both	2013	8	(5 to 11)	12.3	(7.9 to 17.9)
Dominican								
Republic	Deaths	Ischemic Stroke	Male	1990	946	(713 to 1,137)	69.6	(52.8 to 83.8)
				2013	2,421	(1,794 to 3,467)	72.8	(53.9 to 104.3)
			Female	1990	795	(637 to 943)	61.6	(49.2 to 73.5)
				2013	2,320	(1,697 to 3,146)	63.2	(46.4 to 85.6)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	1,741	(1,429 to 1,958)	65.7	(54.2 to 73.9)
				2013	4,741	(3,843 to 6,168)	67.9	(55.1 to 88.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	556	(447 to 719)	34.2	(27.1 to 45.6)
				2013	1,622	(1,093 to 2,299)	45.6	(30.6 to 64.8)
			Female	1990	533	(453 to 745)	33.8	(28.4 to 48.2)
				2013	1,514	(990 to 2,095)	39.8	(25.9 to 55.3)
			Both	1990	1,090	(944 to 1,372)	34.1	(29.0 to 43.4)
				2013	3,136	(2,289 to 4,084)	42.7	(31.0 to 56.0)
	DALYs	Ischemic Stroke	Male	1990	18,828	(13,909 to 22,594)	1101.6	(828.8 to 1318.4)
				2013	39,310	(28,562 to 57,548)	1117.1	(818.2 to 1620.4)
			<u> </u>					
			Female	1990	14,342	(11,266 to 16,661)	890.9	(714.5 to 1032.0)
			Female		-		890.9 849.4	(714.5 to 1032.0) (641.5 to 1206.6)
			Female Both	1990	14,342	(11,266 to 16,661)		,

				2013	70,998	(56,651 to 96,213)	980.5	(781.5 to 1314.9)
		Hemorrhagic						
		Stroke	Male	1990	15,244	(12,417 to 19,300)	714.4	(581.5 to 911.0)
				2013	36,581	(25,170 to 51,309)	932.6	(633.6 to 1317.5)
			Female	1990	14,530	(12,593 to 19,713)	698	(599.4 to 950.9)
				2013	29,939	(20,994 to 41,521)	736.1	(504.2 to 1025.4)
			Both	1990	29,775	(26,056 to 36,675)	706.8	(615.3 to 876.8)
				2013	66,521	(49,933 to 85,159)	833	(619.6 to 1076.6)
	YLDs	Ischemic Stroke	Male	1990	806	(530 to 1,120)	48.9	(32.0 to 67.3)
				2013	1,633	(1,075 to 2,340)	46.2	(30.6 to 66.8)
			Female	1990	533	(362 to 750)	34.3	(23.3 to 48.1)
				2013	1,288	(834 to 1,846)	34.3	(22.1 to 49.0)
			Both	1990	1,340	(922 to 1,832)	41.7	(28.4 to 56.5)
				2013	2,922	(1,968 to 4,069)	40.1	(27.1 to 55.8)
		Hemorrhagic						
		Stroke	Male	1990	310	(180 to 512)	16.3	(9.3 to 27.5)
				2013	558	(327 to 894)	14.7	(8.4 to 23.9)
			Female	1990	186	(117 to 272)	10.3	(6.5 to 15.0)
				2013	469	(294 to 692)	11.8	(7.4 to 17.4)
			Both	1990	496	(321 to 773)	13.4	(8.5 to 20.9)
			Both	2013	1,028	(663 to 1,482)	13.2	(8.5 to 19.2)
Ecuador	Deaths	Ischemic Stroke	Male	1990	941	(703 to 1,121)	46.7	(35.6 to 55.8)
				2013	1,731	(1,305 to 2,293)	35.1	(26.5 to 46.1)
			Female	1990	892	(618 to 994)	38.5	(27.2 to 43.4)
				2013	1,601	(1,256 to 2,367)	27.6	(21.6 to 40.7)
			Both	1990	1,833	(1,435 to 2,060)	42.4	(33.4 to 47.7)
				2013	3,333	(2,708 to 4,482)	31.2	(25.4 to 41.6)
		Hemorrhagic						•
	Deaths	Stroke	Male	1990	572	(466 to 747)	23.3	(18.6 to 31.3)

					Cou	ints, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
-				2013	1,414	(951 to 1,862)	25.7	(17.1 to 33.9)
			Female	1990	603	(512 to 815)	22.5	(18.7 to 30.7)
				2013	1,268	(854 to 1,681)	20.7	(14.0 to 27.4)
			Both	1990	1,175	(1,041 to 1,479)	22.9	(20.1 to 29.5)
				2013	2,682	(1,997 to 3,261)	23.1	(17.1 to 28.1)
	DALYs	Ischemic Stroke	Male	1990	18,651	(13,398 to 22,257)	746.1	(550.1 to 888.3)
				2013	28,119	(20,850 to 39,099)	516.5	(385.6 to 696.6)
			Female	1990	15,215	(9,998 to 17,088)	560.7	(374.6 to 627.4)
				2013	20,646	(15,506 to 33,237)	342.8	(258.1 to 550.7)
			Both	1990	33,867	(25,571 to 38,420)	649	(498.2 to 733.7)
				2013	48,766	(38,146 to 69,375)	425.8	(335.7 to 595.9)
		Hemorrhagic						
		Stroke	Male	1990	17,059	(14,150 to 21,469)	534.7	(440.8 to 685.8)
				2013	36,498	(25,491 to 48,779)	577.2	(398.3 to 768.9)
			Female	1990	16,720	(14,551 to 22,109)	508.1	(439.4 to 674.0)
				2013	29,355	(20,644 to 39,750)	441.1	(307.2 to 592.7)
			Both	1990	33,780	(30,069 to 41,380)	520.6	(463.4 to 645.5)
				2013	65,853	(51,009 to 81,078)	507.2	(389.1 to 622.4)
	YLDs	Ischemic Stroke	Male	1990	404	(262 to 574)	15.8	(10.3 to 22.4)
				2013	929	(597 to 1,334)	16.6	(10.8 to 23.8)
			Female	1990	229	(145 to 328)	8.3	(5.2 to 11.9)
				2013	495	(325 to 721)	8.1	(5.3 to 11.7)
			Both	1990	634	(417 to 872)	11.9	(7.8 to 16.4)
				2013	1,424	(962 to 2,001)	12.1	(8.2 to 17.2)
		Hemorrhagic						
		Stroke	Male	1990	170	(107 to 260)	5.6	(3.5 to 8.8)
				2013	389	(244 to 593)	6.3	(3.9 to 9.6)

			Female	1990	101	(64 to 147)	3.2	(2.0 to 4.7)
				2013	244	(154 to 362)	3.7	(2.3 to 5.6)
			Both	1990	272	(179 to 389)	4.4	(2.8 to 6.3)
			Both	2013	633	(420 to 908)	4.9	(3.3 to 7.2)
Egypt	Deaths	Ischemic Stroke	Male	1990	13,213	(10,325 to 16,360)	132.9	(102.7 to 163.9)
				2013	19,831	(16,209 to 24,879)	111	(91.2 to 138.4)
			Female	1990	15,084	(11,439 to 17,873)	104.7	(79.0 to 124.2)
				2013	21,009	(17,019 to 26,387)	84.3	(68.2 to 105.7)
			Both	1990	28,297	(23,647 to 32,048)	115.7	(95.7 to 130.9)
				2013	40,840	(35,648 to 48,869)	95.1	(83.4 to 113.2)
		Hemorrhagic						, ,
	Deaths	Stroke	Male	1990	17,419	(13,188 to 20,785)	121	(85.2 to 148.0)
				2013	23,019	(18,819 to 27,921)	103.3	(83.6 to 125.4)
			Female	1990	17,526	(13,447 to 20,609)	105.8	(79.7 to 126.2)
				2013	22,085	(18,004 to 27,661)	84	(68.5 to 106.3)
			Both	1990	34,946	(28,480 to 40,169)	113.2	(89.4 to 131.3)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	45,104	(39,361 to 52,519)	93.3	(80.9 to 109.8)
						(206,609 to		
	DALYs	Ischemic Stroke	Male	1990	271,604	346,190)	1992.4	(1566.1 to 2476.6)
						(289,175 to		
				2013	361,085	463,692)	1607.7	(1307.1 to 2004.3)
			5 -	1000	272.027	(205,531 to	45644	/4470 0 to 4054 4)
			Female	1990	273,037	339,201) (263,397 to	1564.4	(1179.8 to 1854.4)
				2013	325,944	414,696)	1171.1	(950.0 to 1482.5)
				2013	323,377	(449,485 to	11/1.1	(550.0 to 1402.5)
			Both	1990	544,641	635,442)	1740.6	(1462.1 to 1983.8)
				2013	687,030	(597,221 to	1359.6	(1184.9 to 1641.5)
1								

						834,869)		
		Hemorrhagic				(540,038 to		
		Stroke	Male	1990	692,056	836,254)	2923.9	(2252.3 to 3488.8)
						(543,763 to		
				2013	659,363	790,403)	2154.6	(1766.8 to 2585.0)
						(378,922 to		
			Female	1990	519,378	667,917)	2200.3	(1697.6 to 2612.1)
						(368,691 to		
				2013	448,261	546,499)	1441.6	(1187.8 to 1766.6)
				1000		(966,552 to		(2000 4) 200 7 4)
			Both	1990	1,211,435	1,392,435)	2557.8	(2088.1 to 2927.1)
				2012	1 107 624	(964,914 to	1700 1	/15C0 0 to 20C2 0\
	\"."			2013	1,107,624	1,265,227)	1790.1	(1568.9 to 2063.9)
	YLDs	Ischemic Stroke	Male	1990	1,491	(949 to 2,113)	9.6	(6.1 to 13.5)
				2013	2,678	(1,779 to 3,844)	9.7	(6.4 to 13.7)
			Female	1990	1,384	(889 to 2,030)	7.7	(4.9 to 11.2)
				2013	2,259	(1,421 to 3,254)	7.3	(4.6 to 10.6)
			Both	1990	2,875	(1,905 to 3,962)	8.5	(5.7 to 11.7)
				2013	4,938	(3,289 to 6,777)	8.4	(5.6 to 11.7)
		Hemorrhagic						
		Stroke	Male	1990	210	(125 to 314)	1.2	(0.7 to 1.7)
				2013	372	(217 to 597)	1.2	(0.7 to 1.9)
			Female	1990	261	(145 to 412)	1.4	(0.8 to 2.1)
				2013	413	(221 to 658)	1.3	(0.7 to 2.0)
			Both	1990	472	(301 to 701)	1.3	(0.8 to 1.9)
			Both	2013	785	(499 to 1,144)	1.3	(0.8 to 1.8)
El Salvador	Deaths	Ischemic Stroke	Male	1990	421	(248 to 549)	37.4	(22.0 to 49.2)
				2013	325	(222 to 521)	16.3	(11.2 to 26.0)
			Female	1990	474	(251 to 568)	34.1	(18.2 to 41.1)
				2013	391	(279 to 635)	14.2	(10.1 to 23.0)

			Both	1990	895	(546 to 1,090)	35.6	(21.6 to 43.6)
				2013	717	(545 to 1,076)	15.1	(11.5 to 22.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	265	(188 to 436)	20.9	(14.1 to 36.4)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	568	(395 to 746)	28.1	(19.5 to 36.9)
			Female	1990	303	(246 to 495)	19.9	(15.7 to 33.8)
				2013	714	(500 to 940)	26	(18.3 to 34.2)
			Both	1990	569	(447 to 878)	20.4	(15.4 to 32.9)
				2013	1,283	(976 to 1,557)	26.9	(20.4 to 32.6)
	DALYs	Ischemic Stroke	Male	1990	8,969	(5,489 to 11,407)	623.4	(372.4 to 801.7)
				2013	5,358	(3,801 to 8,374)	258.7	(183.4 to 405.1)
			Female	1990	8,796	(4,711 to 10,435)	528.3	(280.9 to 627.3)
				2013	5,370	(3,909 to 8,714)	197.2	(143.5 to 319.0)
			Both	1990	17,766	(11,063 to 21,491)	571.7	(351.8 to 694.4)
				2013	10,728	(8,325 to 15,983)	223.9	(173.5 to 333.2)
		Hemorrhagic						
		Stroke	Male	1990	8,008	(6,244 to 11,718)	476.1	(359.8 to 743.9)
				2013	11,579	(8,550 to 15,073)	538.7	(392.6 to 700.0)
			Female	1990	8,289	(7,185 to 11,696)	442.8	(381.0 to 650.7)
				2013	12,454	(9,228 to 17,009)	451	(334.1 to 613.9)
			Both	1990	16,298	(13,933 to 22,139)	458.1	(380.6 to 656.5)
				2013	24,033	(19,439 to 29,386)	489.6	(395.5 to 598.9)
	YLDs	Ischemic Stroke	Male	1990	84	(53 to 122)	6.9	(4.4 to 10.1)
				2013	135	(83 to 200)	6.9	(4.2 to 10.2)
			Female	1990	83	(53 to 119)	5.7	(3.7 to 8.2)
				2013	149	(93 to 237)	5.7	(3.5 to 9.0)
			Both	1990	167	(113 to 234)	6.2	(4.2 to 8.7)

				2013	284	(187 to 409)	6.2	(4.1 to 9.0)
		Hemorrhagic		2013	204	(10/ (0 403)	0.2	(4.1 (0 3.0)
		Stroke	Male	1990	261	(171 to 359)	19.1	(12.5 to 26.4)
				2013	409	(256 to 583)	19.9	(12.3 to 28.5)
			Female	1990	206	(136 to 282)	13	(8.7 to 17.8)
				2013	336	(212 to 486)	12.5	(7.8 to 18.2)
			Both	1990	467	(313 to 628)	15.8	(10.5 to 21.1)
			Both	2013	746	(490 to 1,031)	15.7	(10.4 to 21.8)
Equatorial Guinea	Deaths	Ischemic Stroke	Male	1990	87	(48 to 128)	124	(67.0 to 177.3)
				2013	122	(82 to 188)	103.3	(72.3 to 148.3)
			Female	1990	93	(57 to 134)	118.8	(74.5 to 168.9)
				2013	108	(73 to 178)	101	(68.2 to 161.8)
			Both	1990	181	(114 to 250)	122	(76.3 to 166.9)
				2013	230	(165 to 349)	103	(77.2 to 152.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	116	(74 to 167)	133	(85.3 to 190.7)
				2013	142	(82 to 227)	81.8	(48.9 to 124.6)
			Female	1990	121	(76 to 182)	133.2	(82.9 to 201.1)
				2013	113	(63 to 193)	78.2	(43.8 to 131.6)
			Both	1990	237	(158 to 339)	134.3	(90.1 to 191.5)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	255	(163 to 404)	80.6	(53.7 to 125.4)
	DALYs	Ischemic Stroke	Male	1990	1,820	(1,009 to 2,763)	2092.9	(1152.9 to 3106.0)
		-		2013	2,658	(1,703 to 4,287)	1592.5	(1085.3 to 2450.4)
		-	Female	1990	1,610	(909 to 2,445)	1737.6	(1034.9 to 2571.8)
				2013	1,862	(1,187 to 3,252)	1351.4	(901.7 to 2256.3)
			Both	1990	3,430	(2,075 to 4,914)	1918.5	(1206.9 to 2715.6)
				2013	4,521	(2,995 to 7,214)	1485.9	(1059.6 to 2276.9)

			1			1		
		Hemorrhagic	Mala	1000	2.400	(2.242 to 4.82C)	2111.0	(1000 1 to 1522 0)
		Stroke	Male	1990	3,400	(2,242 to 4,836)	3111.9	(1989.1 to 4522.0)
				2013	4,418	(2,506 to 7,090)	1816.9	(1063.8 to 2894.5)
			Female	1990	2,997	(1,846 to 4,514)	2768.9	(1696.1 to 4156.7)
				2013	3,192	(1,683 to 5,723)	1560	(859.2 to 2703.2)
			Both	1990	6,398	(4,319 to 9,004)	2959.7	(1963.4 to 4228.6)
				2013	7,610	(4,644 to 12,191)	1700.5	(1083.1 to 2724.0)
	YLDs	Ischemic Stroke	Male	1990	14	(9 to 20)	14.2	(9.3 to 20.5)
				2013	31	(19 to 47)	14.7	(9.2 to 21.7)
			Female	1990	10	(7 to 15)	10.6	(6.9 to 15.0)
				2013	21	(13 to 30)	11.2	(7.1 to 16.0)
			Both	1990	25	(17 to 34)	12.4	(8.5 to 17.0)
	Hemorrhagic Stroke		2013	52	(34 to 75)	13	(8.5 to 18.5)	
		Male	1990	4	(2 to 6)	3.8	(2.4 to 5.8)	
				2013	9	(5 to 14)	3.8	(2.4 to 5.8)
			Female	1990	2	(1 to 4)	2.6	(1.7 to 3.8)
				2013	5	(3 to 8)	2.6	(1.6 to 3.9)
			Both	1990	7	(4 to 10)	3.2	(2.1 to 4.5)
			Both	2013	15	(9 to 21)	3.2	(2.1 to 4.7)
Eritrea	Deaths	Ischemic Stroke	Male	1990	153	(55 to 269)	72.5	(27.0 to 121.3)
				2013	375	(169 to 631)	73.3	(34.3 to 116.9)
			Female	1990	284	(86 to 433)	94.5	(29.1 to 143.1)
				2013	736	(338 to 1,268)	98.8	(46.6 to 163.2)
			Both	1990	438	(197 to 644)	88	(37.2 to 128.3)
				2013	1,111	(625 to 1,694)	90.5	(50.9 to 134.8)
		Hemorrhagic				,		
	Deaths	Stroke	Male	1990	277	(144 to 395)	79.7	(32.1 to 123.5)
				2013	556	(330 to 833)	73.2	(38.7 to 117.0)

			Female	1990	447	(170 to 662)	102.1	(39.2 to 157.8)
				2013	855	(445 to 1,315)	88.5	(43.9 to 140.0)
			Both	1990	724	(424 to 954)	94.3	(52.2 to 131.6)
				2013	1,411	(940 to 1,903)	83.1	(51.7 to 117.2)
	DALYs	Ischemic Stroke	Male	1990	3,775	(1,365 to 6,816)	1162.2	(432.0 to 1998.4)
				2013	8,200	(3,630 to 14,026)	1124.8	(516.8 to 1855.4)
			Female	1990	5,716	(1,733 to 9,063)	1402.6	(431.9 to 2118.3)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	13,076	(5,848 to 23,793)	1406	(650.2 to 2433.4)
			Both	1990	9,491	(4,431 to 13,957)	1328.6	(587.3 to 1919.1)
				2013	21,277	(11,989 to 33,022)	1310.9	(739.9 to 1991.6)
		Hemorrhagic						
		Stroke	Male	1990	8,778	(5,085 to 12,456)	1661.1	(790.7 to 2436.6)
				2013	15,989	(9,855 to 23,097)	1469.9	(841.4 to 2255.1)
			Female	1990	12,418	(4,538 to 18,367)	2045.1	(784.1 to 3041.6)
				2013	20,729	(10,840 to 32,706)	1675.3	(875.6 to 2566.8)
			Both	1990	21,197	(12,471 to 28,008)	1895.1	(1097.0 to 2529.1)
				2013	36,718	(25,460 to 49,497)	1599.4	(1059.6 to 2158.8)
	YLDs	Ischemic Stroke	Male	1990	60	(37 to 90)	10.2	(6.7 to 14.5)
				2013	131	(83 to 185)	10.9	(7.0 to 15.3)
			Female	1990	45	(27 to 66)	6.6	(4.1 to 9.8)
				2013	102	(65 to 148)	6.9	(4.5 to 9.9)
			Both	1990	105	(68 to 146)	8.1	(5.4 to 11.1)
				2013	233	(156 to 324)	8.6	(5.8 to 11.7)
		Hemorrhagic						
		Stroke	Male	1990	23	(14 to 35)	2.9	(1.8 to 4.2)
				2013	48	(30 to 71)	3	(1.9 to 4.4)
			Female	1990	14	(8 to 21)	1.7	(1.1 to 2.4)

				2013	30	(18 to 47)	1.7	(1.0 to 2.6)
			Both	1990	37	(23 to 53)	2.2	(1.4 to 3.2)
			Both	2013	79	(50 to 114)	2.3	(1.4 to 3.2)
Estonia	Deaths	Ischemic Stroke	Male	1990	746	(616 to 901)	136.6	(112.6 to 164.6)
				2013	670	(541 to 812)	87.1	(70.3 to 105.8)
			Female	1990	1,723	(1,306 to 1,897)	129.6	(97.9 to 143.1)
				2013	956	(707 to 1,573)	50.2	(37.4 to 82.8)
			Both	1990	2,470	(2,023 to 2,718)	133.7	(109.6 to 147.0)
				2013	1,627	(1,313 to 2,277)	63	(51.2 to 87.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	250	(154 to 357)	38.4	(23.6 to 55.7)
				2013	221	(130 to 307)	28.5	(16.7 to 39.4)
			Female	1990	426	(267 to 527)	34.8	(22.2 to 42.8)
				2013	200	(146 to 339)	12.8	(9.5 to 20.6)
			Both	1990	677	(450 to 840)	36.8	(24.6 to 45.6)
				2013	421	(332 to 581)	18.9	(15.0 to 25.4)
	DALYs	Ischemic Stroke	Male	1990	12,539	(10,191 to 15,262)	2036.4	(1662.7 to 2444.2)
				2013	9,893	(8,113 to 11,867)	1261.1	(1034.5 to 1513.9)
			Female	1990	20,805	(15,463 to 23,223)	1637.1	(1214.9 to 1840.2)
				2013	9,469	(7,159 to 15,467)	574.7	(436.4 to 935.4)
			Both	1990	33,344	(27,405 to 37,272)	1799.5	(1480.7 to 2011.5)
				2013	19,362	(16,052 to 25,733)	831.4	(690.3 to 1099.7)
					Cou	nts, All ages	Rate per	100,000, Age-standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	6,394	(4,159 to 8,988)	879.2	(572.1 to 1232.9)
				2013	4,740	(2,903 to 6,290)	610.7	(369.8 to 806.9)
			Female	1990	7,750	(5,116 to 9,212)	691.7	(461.0 to 816.6)

						,		
				2013	2,920	(2,200 to 4,600)	233	(179.1 to 358.4)
			Both	1990	14,144	(9,786 to 17,380)	777.6	(540.4 to 953.1)
				2013	7,660	(6,130 to 9,829)	392.1	(314.9 to 492.6)
	YLDs	Ischemic Stroke	Male	1990	331	(212 to 471)	46.2	(29.7 to 65.8)
				2013	354	(218 to 517)	46.2	(28.8 to 67.2)
			Female	1990	350	(233 to 493)	31.6	(20.9 to 44.8)
	Hemorrhagic Stroke			2013	362	(235 to 502)	29.8	(19.7 to 41.4)
			Both	1990	682	(459 to 931)	37.6	(25.4 to 51.4)
				2013	717	(484 to 985)	36.8	(24.8 to 50.6)
		Male	1990	20	(11 to 33)	2.8	(1.6 to 4.6)	
			2013	19	(9 to 35)	2.6	(1.4 to 4.6)	
			Female	1990	22	(12 to 37)	2.2	(1.2 to 3.6)
				2013	23	(12 to 40)	2.1	(1.2 to 3.5)
			Both	1990	43	(24 to 65)	2.4	(1.4 to 3.7)
			Both	2013	42	(24 to 67)	2.4	(1.4 to 3.7)
Ethiopia	Deaths	Ischemic Stroke	Male	1990	6,236	(2,547 to 10,302)	88.2	(36.9 to 140.8)
				2013	10,186	(3,248 to 17,650)	75.1	(24.4 to 127.0)
			Female	1990	6,147	(3,206 to 8,395)	89.7	(46.8 to 122.3)
				2013	13,077	(4,788 to 18,927)	88.6	(33.3 to 127.8)
			Both	1990	12,384	(6,860 to 16,483)	89	(50.3 to 119.7)
				2013	23,264	(9,603 to 33,336)	82.3	(34.5 to 118.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	8,577	(4,056 to 11,721)	98.9	(40.1 to 140.7)
				2013	13,488	(6,316 to 20,961)	85.3	(36.2 to 137.0)
			Female	1990	9,098	(6,270 to 13,425)	105.4	(68.9 to 158.5)
				2013	17,024	(11,414 to 24,784)	99.6	(64.7 to 149.4)
			Both	1990	17,675	(12,680 to 22,798)	102.2	(68.6 to 137.2)
				2013	30,512	(20,565 to 42,486)	92.9	(59.8 to 135.1)

						(50,255 to		
	DALYs	Ischemic Stroke	Male	1990	123,339	212,765)	1423	(573.7 to 2372.8)
						(54,193 to		
				2013	172,140	310,439)	1075.2	(341.3 to 1891.0)
						(56,010 to		
			Female	1990	112,227	160,990)	1342.5	(698.1 to 1851.9)
						(72,079 to		
				2013	202,362	297,417)	1193.8	(437.7 to 1735.3)
						(129,218 to		
			Both	1990	235,567	320,545)	1381.6	(764.8 to 1848.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(149,866 to		
				2013	374,502	542,326)	1138.8	(464.0 to 1638.6)
		Hemorrhagic				(122,959 to		
		Stroke	Male	1990	220,713	292,175)	2052.1	(992.5 to 2773.5)
						(162,552 to		
				2013	301,583	441,827)	1564.5	(766.5 to 2389.4)
						(161,906 to		
			Female	1990	224,426	317,175)	2089.3	(1442.9 to 3060.6)
						(236,158 to		
				2013	350,653	489,419)	1752.1	(1174.7 to 2498.8)
						(336,355 to		
			Both	1990	445,140	547,103)	2070.4	(1496.1 to 2649.2)
						(470,602 to		(,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
				2013	652,236	863,899)	1664.6	(1154.7 to 2273.9)
	YLDs	Ischemic Stroke	Male	1990	602	(406 to 833)	5.4	(3.7 to 7.6)
				2013	1,340	(904 to 1,870)	6.3	(4.2 to 8.8)
			Female	1990	515	(339 to 723)	4.6	(3.0 to 6.3)
				2013	1,008	(671 to 1,432)	4.4	(2.9 to 6.2)
E								

			Both	1990	1,117	(756 to 1,534)	5	(3.4 to 6.8)
				2013	2,348	(1,582 to 3,230)	5.3	(3.6 to 7.2)
		Hemorrhagic						
		Stroke	Male	1990	188	(123 to 267)	1.4	(0.9 to 2.1)
				2013	440	(283 to 633)	1.7	(1.1 to 2.5)
			Female	1990	146	(96 to 206)	1.1	(0.7 to 1.6)
				2013	281	(185 to 401)	1	(0.7 to 1.5)
			Both	1990	334	(225 to 463)	1.3	(0.8 to 1.8)
			Both	2013	722	(484 to 1,008)	1.4	(0.9 to 1.9)
Federated States								
of Micronesia	Deaths	Ischemic Stroke	Male	1990	14	(8 to 21)	93	(55.0 to 131.4)
				2013	17	(11 to 26)	85.5	(56.3 to 128.2)
			Female	1990	9	(5 to 14)	57	(36.7 to 88.7)
				2013	13	(8 to 21)	58.9	(35.5 to 88.8)
			Both	1990	24	(15 to 34)	74.2	(49.7 to 102.1)
				2013	31	(20 to 45)	70.8	(48.3 to 101.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	25	(15 to 37)	130.7	(78.4 to 186.4)
				2013	29	(18 to 45)	116.2	(71.7 to 173.0)
			Female	1990	15	(9 to 22)	79	(52.0 to 120.0)
				2013	20	(12 to 32)	76.8	(46.2 to 118.3)
			Both	1990	40	(26 to 57)	104.3	(68.9 to 146.2)
				2013	50	(31 to 76)	95.5	(61.4 to 139.2)
	DALYs	Ischemic Stroke	Male	1990	315	(174 to 487)	1615.3	(918.2 to 2407.6)
_				2013	361	(216 to 588)	1426.2	(914.7 to 2232.7)
			Female	1990	184	(111 to 286)	925.6	(577.8 to 1439.8)
					Counts, All	Ages	Rate	Per 100,000

Country

measure

Stroke Type

sex

Counts, All Ages Rate Per 100,000

Age- standardized

year Mean 95% UI Mean 95% UI

				2013	244	(127 to 398)	903.5	(497.5 to 1450.9)
			Both	1990	499	(303 to 743)	1259.5	(791.1 to 1804.2)
				2013	605	(362 to 961)	1147.2	(715.7 to 1746.4)
		Hemorrhagic				,		,
		Stroke	Male	1990	764	(465 to 1,163)	3083.7	(1814.7 to 4631.0)
				2013	857	(509 to 1,367)	2675.2	(1614.3 to 4156.1)
			Female	1990	394	(247 to 613)	1654.2	(1045.5 to 2603.5)
				2013	474	(271 to 767)	1522.7	(878.7 to 2437.9)
			Both	1990	1,158	(737 to 1,744)	2363.2	(1511.1 to 3510.1)
				2013	1,331	(816 to 2,121)	2083.5	(1290.1 to 3237.2)
	YLDs	Ischemic Stroke	Male	1990	4	(2 to 6)	25	(16.1 to 36.0)
				2013	6	(4 to 9)	25.6	(16.5 to 36.9)
			Female	1990	3	(2 to 5)	18.8	(12.2 to 27.2)
				2013	5	(3 to 8)	21	(13.9 to 28.9)
			Both	1990	8	(5 to 11)	21.8	(14.5 to 31.0)
				2013	12	(7 to 16)	23.1	(15.3 to 31.7)
		Hemorrhagic						
		Stroke	Male	1990	5	(3 to 8)	29.4	(19.0 to 41.9)
				2013	8	(5 to 11)	30.5	(20.4 to 43.5)
			Female	1990	4	(2 to 6)	20.2	(13.2 to 29.0)
				2013	6	(4 to 9)	22.6	(14.7 to 31.5)
			Both	1990	10	(6 to 14)	24.7	(16.4 to 34.6)
			Both	2013	14	(10 to 20)	26.2	(17.7 to 36.3)
Fiji	Deaths	Ischemic Stroke	Male	1990	94	(60 to 142)	87.7	(58.2 to 125.4)
				2013	95	(67 to 136)	50.2	(35.7 to 69.6)
			Female	1990	108	(51 to 168)	97.9	(49.3 to 143.8)
				2013	98	(72 to 149)	44.5	(32.6 to 66.2)

			Both	1990	202	(126 to 295)	93.1	(58.9 to 128.6)
				2013	194	(152 to 263)	47.1	(37.2 to 63.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	152	(98 to 228)	99.7	(62.1 to 147.7)
				2013	228	(169 to 301)	88.9	(66.7 to 114.6)
			Female	1990	198	(116 to 271)	142.1	(86.6 to 189.3)
				2013	231	(174 to 335)	88.1	(67.9 to 124.5)
			Both	1990	351	(258 to 451)	121.5	(89.4 to 153.7)
				2013	460	(372 to 575)	89.1	(73.8 to 110.7)
	DALYs	Ischemic Stroke	Male	1990	2,212	(1,378 to 3,627)	1458.3	(948.0 to 2202.7)
				2013	2,066	(1,427 to 2,995)	790.2	(568.5 to 1109.5)
			Female	1990	2,351	(1,069 to 3,809)	1592.3	(751.9 to 2492.1)
				2013	1,922	(1,404 to 2,931)	674.3	(501.2 to 1018.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	4,564	(2,804 to 7,047)	1526.5	(956.6 to 2227.7)
				2013	3,988	(3,059 to 5,475)	728.2	(572.5 to 985.8)
		Hemorrhagic						
		Stroke	Male	1990	5,328	(3,543 to 7,889)	2350.7	(1516.9 to 3503.9)
				2013	6,841	(5,054 to 9,171)	1970.3	(1473.6 to 2593.2)
			Female	1990	5,704	(3,370 to 7,893)	3000.2	(1768.9 to 4120.4)
				2013	5,825	(4,268 to 8,521)	1724.9	(1299.5 to 2485.2)
			Both	1990	11,033	(8,225 to 14,151)	2679.2	(1972.1 to 3435.1)
,				2013	12,667	(10,069 to 15,951)	1853.7	(1497.2 to 2300.1)
	YLDs	Ischemic Stroke	Male	1990	34	(22 to 48)	25.3	(16.6 to 34.9)
				2013	64	(40 to 94)	24	(14.9 to 35.3)
			F	1000	20	(10 to 41)	20.6	(13.1 to 30.2)
			Female	1990	28	(18 to 41)	20.6	(13.1 (0.30.2)

	1	ı	, ,		1			
			Both	1990	62	(42 to 84)	22.8	(15.6 to 30.7)
				2013	125	(81 to 177)	22.4	(14.6 to 31.6)
		Hemorrhagic						
		Stroke	Male	1990	47	(31 to 66)	29.9	(19.8 to 41.1)
				2013	86	(57 to 122)	28.9	(18.8 to 41.5)
			Female	1990	35	(22 to 48)	21.9	(14.0 to 30.6)
				2013	70	(44 to 102)	22.5	(14.4 to 32.7)
			Both	1990	82	(56 to 110)	25.8	(17.7 to 34.4)
			Both	2013	156	(103 to 217)	25.5	(16.9 to 35.8)
Finland	Deaths	Ischemic Stroke	Male	1990	1,487	(1,209 to 1,856)	69.7	(56.9 to 87.4)
				2013	1,609	(1,218 to 2,000)	37.8	(28.7 to 46.8)
			Female	1990	2,798	(2,102 to 3,202)	61.7	(46.3 to 70.8)
				2013	2,162	(1,715 to 3,142)	27.9	(22.5 to 40.7)
			Both	1990	4,285	(3,593 to 4,819)	65.4	(54.8 to 73.6)
				2013	3,771	(3,220 to 4,850)	32.3	(27.6 to 41.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	730	(574 to 976)	30.2	(23.4 to 40.6)
				2013	841	(587 to 1,074)	20.3	(14.2 to 25.8)
			Female	1990	1,062	(808 to 1,325)	26.3	(20.2 to 32.7)
				2013	832	(637 to 1,255)	13.3	(10.6 to 19.8)
			Both	1990	1,792	(1,496 to 2,195)	28.2	(23.6 to 34.4)
				2013	1,674	(1,375 to 2,146)	16.5	(13.7 to 20.9)
	DALYs	Ischemic Stroke	Male	1990	23,667	(19,835 to 29,105)	1011.6	(844.3 to 1248.1)
				2013	21,488	(17,297 to 25,747)	508.3	(409.0 to 609.4)
			Female	1990	31,172	(24,193 to 35,228)	724.3	(556.9 to 817.2)
				2013	19,009	(15,708 to 26,448)	291.1	(242.2 to 398.7)
			Both	1990	54,840	(47,501 to 61,422)	843.9	(732.1 to 946.4)
				2013	40,497	(34,913 to 48,853)	388.1	(333.8 to 464.1)
		Hemorrhagic	Male	1990	17,681	(14,427 to 23,265)	675.2	(549.8 to 887.2)

					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	16,166	(11,704 to 19,968)	411.6	(300.8 to 507.4)
			Female	1990	18,429	(14,733 to 22,720)	515.2	(413.3 to 630.8)
				2013	11,351	(9,290 to 16,392)	236.2	(195.9 to 341.9)
			Both	1990	36,111	(31,555 to 42,931)	590.8	(516.7 to 695.3)
				2013	27,518	(23,197 to 33,432)	319.6	(269.2 to 383.0)
	YLDs	Ischemic Stroke	Male	1990	2,594	(1,753 to 3,617)	103.8	(70.5 to 144.5)
				2013	3,256	(2,065 to 4,647)	78.5	(50.0 to 111.9)
			Female	1990	2,564	(1,694 to 3,519)	66	(43.4 to 90.1)
				2013	2,723	(1,775 to 3,851)	50.6	(33.0 to 71.2)
			Both	1990	5,158	(3,526 to 6,965)	81.4	(55.3 to 109.7)
				2013	5,979	(4,055 to 8,175)	63.3	(42.8 to 86.6)
		Hemorrhagic						
		Stroke	Male	1990	722	(467 to 1,033)	28.7	(18.7 to 41.0)
				2013	892	(573 to 1,350)	22.6	(14.6 to 33.8)
			Female	1990	710	(466 to 985)	19.4	(12.8 to 27.1)
				2013	739	(471 to 1,033)	15.8	(10.3 to 21.9)
			Both	1990	1,433	(950 to 1,976)	23.1	(15.3 to 31.7)
			Both	2013	1,632	(1,117 to 2,257)	18.8	(12.8 to 26.1)
France	Deaths	Ischemic Stroke	Male	1990	11,562	(9,234 to 14,185)	38.5	(30.7 to 47.2)
				2013	10,914	(7,876 to 13,895)	20.1	(14.6 to 25.5)
			Female	1990	19,479	(13,669 to 22,745)	32.2	(22.8 to 37.2)
				2013	16,566	(12,723 to 24,286)	16.4	(12.7 to 23.8)
			Both	1990	31,041	(25,040 to 34,953)	35	(28.6 to 39.3)
				2013	27,480	(22,372 to 36,083)	18.1	(14.8 to 23.3)
	Deaths	Hemorrhagic Stroke	Male	1990	8,962	(7,070 to 11,355)	29.5	(23.3 to 37.5)

				2013	9,519	(6,647 to 12,322)	18.7	(13.4 to 23.9)
			Female	1990	12,380	(8,748 to 16,174)	23.1	(16.9 to 29.3)
				2013	12,081	(9,101 to 16,877)	13.9	(10.9 to 19.2)
			Both	1990	21,342	(17,812 to 26,206)	26.1	(22.1 to 31.5)
				2013	21,601	(17,143 to 26,533)	16	(13.0 to 19.2)
						(135,944 to		
	DALYs	Ischemic Stroke	Male	1990	163,087	192,251)	537.9	(447.3 to 633.9)
						(114,086 to		
				2013	145,962	178,742)	298.7	(233.9 to 366.0)
						(141,468 to		
			Female	1990	187,008	213,227)	358.1	(275.0 to 407.2)
				2010		(118,710 to		(470.0)
				2013	147,223	197,032)	190.6	(153.0 to 250.8)
			Both	1990	350.006	(301,967 to 392,056)	434.1	(277.0 to 496.7)
			DOUI	1990	350,096	(244,491 to	434.1	(377.0 to 486.7)
				2013	293,186	354,625)	238.6	(197.0 to 285.7)
				2013	Counts, All	. ,	Rate	Per 100,000
						0	Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
•		Hemorrhagic		•		(134,033 to		
		Stroke	Male	1990	164,931	200,675)	543.7	(444.2 to 658.5)
						(99,967 to		
				2013	133,164	165,152)	298.3	(226.2 to 365.6)
						(115,931 to		
			Female	1990	150,957	187,711)	355.7	(277.9 to 438.9)
				2042	447.040	(94,274 to	402.2	(460.01252.5)
				2013	117,018	155,391)	192.2	(160.0 to 252.5)
			Both	1990	315,888	(274,220 to 368,738)	441.7	(386.0 to 510.1)
			BULII					·
				2013	250,183	(209,169 to	240.7	(202.0 to 278.0)

						293,380)		
	YLDs	Ischemic Stroke	Male	1990	24,896	(16,017 to 35,588)	81.4	(52.2 to 116.0)
				2013	38,863	(25,381 to 54,822)	84.8	(55.4 to 120.0)
			Female	1990	25,010	(16,220 to 35,536)	57.6	(38.1 to 80.8)
				2013	35,395	(21,983 to 51,440)	57.5	(37.0 to 81.2)
			Both	1990	49,906	(34,732 to 68,905)	67.4	(47.2 to 92.1)
						(49,579 to		
				2013	74,258	100,000)	69.9	(47.0 to 94.3)
		Hemorrhagic						
		Stroke	Male	1990	3,325	(1,858 to 5,507)	10.9	(6.1 to 17.9)
				2013	2,397	(1,472 to 3,642)	5.4	(3.4 to 8.2)
			Female	1990	1,500	(890 to 2,255)	3.7	(2.2 to 5.6)
				2013	2,540	(1,452 to 4,149)	4.5	(2.6 to 7.2)
			Both	1990	4,825	(2,866 to 7,356)	6.8	(4.1 to 10.2)
			Both	2013	4,937	(3,117 to 7,281)	4.9	(3.1 to 7.2)
Gabon	Deaths	Ischemic Stroke	Male	1990	329	(199 to 444)	145.9	(89.8 to 192.6)
				2013	574	(385 to 776)	154.5	(105.8 to 204.9)
			Female	1990	314	(182 to 417)	114.6	(68.8 to 152.7)
				2013	545	(361 to 750)	117.7	(77.9 to 161.5)
			Both	1990	643	(412 to 786)	128.4	(83.5 to 156.7)
				2013	1,120	(831 to 1,390)	135	(101.1 to 167.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	366	(266 to 491)	141.6	(102.1 to 196.6)
				2013	605	(417 to 835)	141.6	(99.8 to 196.9)
			Female	1990	356	(246 to 534)	115.3	(79.5 to 178.2)
				2013	513	(310 to 826)	107	(64.1 to 172.7)
			Both	1990	722	(590 to 951)	127.4	(104.2 to 170.7)
				2013	1,119	(867 to 1,516)	124	(96.0 to 168.1)
	DALYs	Ischemic Stroke	Male	1990	6,116	(3,574 to 8,538)	2345.5	(1388.9 to 3209.2)

Female 1990									
Country Measure Stroke Type Sex Year Mean 95% Ul Mean 95% Ul					2013	10,548	(6,380 to 15,866)	2510.1	(1609.0 to 3607.4)
Both 1990 10,967 (6,862 to 13,570) 1905.7 (1205.5 to 2337.				Female	1990	4,850	(2,621 to 6,532)	1551.2	(874.6 to 2064.7)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139.1 Female 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139.1 Female 1990 7,604 (5,259 to 10,761) 2179.3 (1521.0 to 3108.1 Both 1990 16,904 (13,872 to 21,118) 2012.9 (1228.2 to 3238.1 Both 1990 16,904 (13,872 to 21,118) 2619.5 (2154.1 to 3294.1 YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) Female 1990 38 (24 to 54) 11.1 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) </td <td></td> <td></td> <td></td> <td></td> <td>2013</td> <td>7,613</td> <td>(4,735 to 10,814)</td> <td>1581</td> <td>(995.5 to 2242.7)</td>					2013	7,613	(4,735 to 10,814)	1581	(995.5 to 2242.7)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139.10 to				Both	1990	10,967	(6,862 to 13,570)	1905.7	(1205.5 to 2337.3)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139. 1 2013 16,190 (10,808 to 22,560) 3214.2 (2171.7 to 4479. 1 5 2013 16,190 (10,808 to 22,560) 3214.2 (2171.7 to 4479. 2 6 1990 7,604 (5,259 to 10,761) 2179.3 (1521.0 to 3108. 3 2013 10,361 (6,398 to 16,801) 2012.9 (1228.2 to 3238. 4 80 to 1990 16,904 (13,872 to 21,118) 2619.5 (2154.1 to 3294. 5 2013 26,551 (20,420 to 35,371) 2607.1 (2010.5 to 3487. 6 2013 26,551 (20,420 to 35,371) 2607.1 (2010.5 to 3487. 7 4 2013 65 (41 to 94) 13.9 (8.6 to 19.9) 8 2013 58 (36 to 83) 11.2						Counts, All	Ages	Rate	Per 100,000
Hemorrhagic Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139.1								Age-	standardized
Hemorrhagic Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139	Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Stroke Male 1990 9,299 (6,881 to 12,135) 3131.4 (2311.2 to 4139. 2013 16,190 (10,808 to 22,560) 3214.2 (2171.7 to 4479. Female 1990 7,604 (5,259 to 10,761) 2179.3 (1521.0 to 3108. 2013 10,361 (6,398 to 16,801) 2012.9 (1228.2 to 3238. Both 1990 16,904 (13,872 to 21,118) 2619.5 (2154.1 to 3294. YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) 2013 65 (41 to 94) 13.9 (8.6 to 19.9) Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) 2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)					2013	18,162	(12,440 to 24,046)	2023.3	(1429.8 to 2632.4)
2013 16,190 (10,808 to 22,560) 3214.2 (2171.7 to 4479.1 to 4479.1 to 1479.1 to 1479.1			•						
Female 1990 7,604 (5,259 to 10,761) 2179.3 (1521.0 to 3108. 2013 10,361 (6,398 to 16,801) 2012.9 (1228.2 to 3238.4 Both 1990 16,904 (13,872 to 21,118) 2619.5 (2154.1 to 3294.4 YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) YLDs Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) YLDs Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) YLDs Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) Both 1990 80 (53 to 17.1) 12.5 (8.1 to 17.2)			Stroke	Male	1990	9,299			(2311.2 to 4139.2)
2013 10,361 (6,398 to 16,801) 2012.9 (1228.2 to 3238.4					2013	16,190	(10,808 to 22,560)	3214.2	(2171.7 to 4479.3)
Both 1990 16,904 (13,872 to 21,118) 2619.5 (2154.1 to 3294.3) YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) YLDs Female 1990 38 (24 to 54) 13.9 (8.6 to 19.9) YLDs Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) YLDs Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) YLDs Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6)				Female	1990	7,604	(5,259 to 10,761)	2179.3	(1521.0 to 3108.1)
YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) 2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)					2013	10,361	(6,398 to 16,801)	2012.9	(1228.2 to 3238.8)
YLDs Ischemic Stroke Male 1990 42 (27 to 59) 14.9 (9.7 to 21.0) 2013 65 (41 to 94) 13.9 (8.6 to 19.9) Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) 2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)				Both	1990	16,904	(13,872 to 21,118)	2619.5	(2154.1 to 3294.8)
2013 65 (41 to 94) 13.9 (8.6 to 19.9) Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) 2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)					2013	26,551	(20,420 to 35,371)	2607.1	(2010.5 to 3487.0)
Female 1990 38 (24 to 54) 11.1 (7.1 to 15.8) 2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)		YLDs	Ischemic Stroke	Male	1990	42	(27 to 59)	14.9	(9.7 to 21.0)
2013 58 (36 to 83) 11.2 (7.1 to 16.3) Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)					2013	65	(41 to 94)	13.9	(8.6 to 19.9)
Both 1990 80 (53 to 110) 12.8 (8.5 to 17.6) 2013 124 (80 to 171) 12.5 (8.1 to 17.2)				Female	1990	38	(24 to 54)	11.1	(7.1 to 15.8)
2013 124 (80 to 171) 12.5 (8.1 to 17.2)					2013	58	(36 to 83)	11.2	(7.1 to 16.3)
				Both	1990	80	(53 to 110)	12.8	(8.5 to 17.6)
					2013	124	(80 to 171)	12.5	(8.1 to 17.2)
Hemorrnagic Hemorrnagic Hemorrnagic Hemorrnagic Hemorrnagic Hemorrnagic Hemorrnagic Hemorrnagic			Hemorrhagic						
Stroke Male 1990 11 (7 to 16) 3.8 (2.3 to 5.5)			Stroke	Male	1990		(7 to 16)		(2.3 to 5.5)
2013 19 (11 to 29) 3.7 (2.2 to 5.5)					2013	19	(11 to 29)	3.7	(2.2 to 5.5)
Female 1990 9 (5 to 14) 2.7 (1.7 to 4.0)				Female	1990	9	(5 to 14)	2.7	(1.7 to 4.0)
2013 14 (9 to 21) 2.6 (1.6 to 3.8)					2013	14	(9 to 21)	2.6	(1.6 to 3.8)
Both 1990 21 (13 to 29) 3.2 (2.1 to 4.5)				Both	1990	21	(13 to 29)	3.2	(2.1 to 4.5)
Both 2013 34 (22 to 48) 3.1 (2.0 to 4.4)				Both	2013	34	(22 to 48)	3.1	(2.0 to 4.4)
Georgia Deaths Ischemic Stroke Male 1990 2,257 (1,601 to 2,939) 129.3 (92.1 to 167.8)	Georgia	Deaths	Ischemic Stroke	Male	1990	2,257	(1,601 to 2,939)	129.3	(92.1 to 167.8)

F						•		
				2013	2,413	(1,868 to 3,379)	100.2	(77.4 to 140.7)
			Female	1990	3,178	(2,356 to 4,024)	96.8	(71.3 to 122.4)
				2013	2,436	(1,913 to 3,195)	56.1	(44.2 to 73.1)
			Both	1990	5,436	(4,209 to 6,434)	109	(84.4 to 128.7)
				2013	4,849	(4,107 to 6,266)	73.3	(62.1 to 94.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,681	(2,105 to 3,455)	135.2	(104.5 to 179.7)
				2013	3,887	(2,776 to 4,628)	162.8	(117.5 to 194.5)
			Female	1990	3,255	(2,442 to 4,458)	97.3	(72.9 to 133.4)
				2013	4,083	(2,669 to 4,974)	99.7	(66.6 to 121.2)
			Both	1990	5,936	(4,947 to 7,586)	111.8	(92.6 to 144.1)
				2013	7,970	(5,855 to 9,124)	125.4	(93.0 to 143.3)
	DALYs	Ischemic Stroke	Male	1990	39,826	(28,419 to 52,420)	1949.5	(1398.2 to 2522.9)
				2013	37,346	(28,754 to 52,534)	1512	(1168.9 to 2125.2)
			Female	1990	40,857	(29,946 to 52,913)	1219.1	(894.1 to 1579.9)
				2013	26,951	(21,280 to 34,669)	673.1	(531.1 to 869.5)
			Both	1990	80,684	(62,574 to 97,983)	1507.7	(1170.5 to 1825.5)
				2013	64,297	(53,961 to 83,303)	1016.1	(849.2 to 1318.3)
		Hemorrhagic						
		Stroke	Male	1990	61,212	(48,704 to 77,284)	2649.5	(2117.7 to 3367.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	71,664	(53,335 to 85,772)	2964	(2225.4 to 3542.8)
			Female	1990	57,875	(45,843 to 74,722)	1720.8	(1371.6 to 2234.9)
				2013	53,779	(37,887 to 65,288)	1490.4	(1070.9 to 1796.9)
						(101,166 to		
			Both	1990	119,087	147,502)	2108.5	(1782.4 to 2630.1)
				2010		(97,929 to	2440	(46746, 04405)
				2013	125,444	143,650)	2118	(1674.6 to 2418.8)

	_	T			1	,		
	YLDs	Ischemic Stroke	Male	1990	198	(126 to 297)	8.5	(5.5 to 12.6)
				2013	210	(126 to 311)	8.6	(5.2 to 12.8)
			Female	1990	273	(167 to 403)	8.1	(5.0 to 11.9)
				2013	271	(172 to 396)	7.9	(5.0 to 11.4)
			Both	1990	472	(316 to 671)	8.3	(5.6 to 11.8)
				2013	481	(313 to 661)	8.2	(5.3 to 11.3)
		Hemorrhagic						
		Stroke	Male	1990	197	(123 to 293)	8	(5.0 to 12.0)
				2013	190	(116 to 287)	8	(4.9 to 12.1)
			Female	1990	156	(97 to 234)	4.8	(3.0 to 7.1)
				2013	159	(98 to 234)	5	(3.1 to 7.2)
			Both	1990	353	(229 to 497)	6.2	(4.0 to 8.7)
			Both	2013	350	(235 to 493)	6.3	(4.2 to 8.9)
Germany	Deaths	Ischemic Stroke	Male	1990	27,606	(22,086 to 33,383)	70.1	(55.8 to 84.7)
				2013	21,498	(16,124 to 27,296)	28.7	(21.5 to 36.5)
			Female	1990	55,502	(39,143 to 62,849)	61.2	(43.8 to 69.2)
				2013	31,488	(24,204 to 46,501)	24.9	(19.3 to 36.6)
			Both	1990	83,109	(65,982 to 92,617)	64.7	(51.6 to 72.1)
				2013	52,986	(43,056 to 68,992)	26.8	(21.9 to 34.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	13,851	(10,570 to 17,824)	33.7	(25.7 to 43.5)
				2013	13,320	(8,838 to 17,994)	18.4	(12.3 to 24.9)
			Female	1990	25,176	(17,597 to 30,750)	30.6	(21.9 to 37.4)
				2013	15,677	(11,994 to 24,147)	13.9	(11.1 to 21.0)
			Both	1990	39,027	(31,085 to 46,522)	32.1	(25.9 to 38.1)
				2013	28,998	(23,561 to 38,360)	16	(13.1 to 20.8)
						(321,909 to		
	DALYs	Ischemic Stroke	Male	1990	391,750	463,650)	957.6	(787.3 to 1129.6)
				2013	319,296	(255,061 to	435.8	(347.6 to 528.2)

						207 500)		
						387,568)		
						(424,305 to		
			Female	1990	577,330	644,195)	699.2	(516.4 to 781.1)
						(268,044 to		
				2013	325,205	436,797)	304	(249.4 to 406.0)
						(804,862 to		
			Both	1990	969,080	1,070,095)	799.8	(668.5 to 885.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		,		•		(545,740 to		
						775,506)		
				2013	644,502	, ,	365.2	(309.4 to 436.9)
		Hemorrhagic				(199,582 to		·
		Stroke	Male	1990	254,039	316,902)	590.3	(464.9 to 737.8)
						(148,935 to		·
				2013	202,975	257,423)	299	(223.8 to 377.3)
						(255,175 to		·
			Female	1990	334,785	406,305)	485.1	(379.8 to 588.4)
						(153,147 to		
				2013	184,433	258,615)	213.4	(179.6 to 288.9)
						(498,739 to		
			Both	1990	588,825	680,147)	530.8	(452.5 to 609.2)
						(328,395 to		
				2013	387,408	479,657)	252.7	(216.2 to 305.8)
	YLDs	Ischemic Stroke	Male	1990	42,948	(29,440 to 59,483)	100.8	(69.2 to 139.5)
						(53,765 to		
				2013	80,541	109,201)	114.4	(76.1 to 155.3)
			Female	1990	46,731	(30,945 to 66,506)	65.3	(43.1 to 93.7)
					-	(51,480 to		. ,
				2013	75,335	101,255)	83.3	(57.1 to 111.7)

	1				1	T		
						(62,290 to		
			Both	1990	89,680	120,738)	79.3	(55.0 to 107.1)
						(109,649 to		
				2013	155,876	207,427)	97.7	(68.6 to 130.8)
		Hemorrhagic						
		Stroke	Male	1990	6,510	(4,052 to 9,370)	15.3	(9.5 to 22.0)
				2013	15,561	(9,871 to 22,564)	23.2	(15.0 to 33.1)
			Female	1990	11,974	(7,383 to 17,507)	18.5	(11.4 to 27.4)
				2013	19,283	(12,709 to 27,629)	24.1	(16.0 to 34.2)
			Both	1990	18,485	(12,008 to 25,643)	17	(11.1 to 23.6)
			Both	2013	34,845	(23,733 to 47,259)	23.6	(15.9 to 32.0)
Ghana	Deaths	Ischemic Stroke	Male	1990	1,195	(889 to 1,604)	66.6	(50.6 to 86.6)
				2013	5,093	(4,008 to 6,155)	130.6	(105.2 to 157.0)
			Female	1990	1,096	(606 to 2,554)	59.3	(33.0 to 134.2)
				2013	6,341	(3,364 to 8,581)	132.2	(70.6 to 177.5)
			Both	1990	2,292	(1,591 to 3,778)	62.9	(43.9 to 104.4)
				2013	11,434	(8,183 to 13,911)	133	(95.3 to 162.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	721	(490 to 983)	24.4	(15.5 to 34.5)
				2013	1,965	(1,418 to 2,685)	33.8	(23.7 to 47.4)
			Female	1990	696	(447 to 1,365)	24.6	(15.2 to 49.1)
				2013	2,434	(1,380 to 3,305)	38.6	(22.3 to 52.2)
			Both	1990	1,418	(1,056 to 2,051)	24.6	(17.1 to 36.6)
				2013	4,399	(3,299 to 5,483)	36.7	(27.8 to 46.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
	DALYs	Ischemic Stroke	Male	1990	26,875	(19,287 to 36,807)	1070.7	(801.5 to 1417.4)
						(82,282 to		
				2013	108,020	132,761)	2136.1	(1675.2 to 2578.8)

			Female	1990	21,801	(12,187 to 50,169)	868.5	(483.7 to 2000.3)
						(54,784 to		
				2013	104,330	145,123)	1863.1	(998.2 to 2553.4)
			Both	1990	48,676	(33,507 to 79,517)	967.6	(679.1 to 1581.7)
						(151,949 to		
				2013	212,350	259,951)	2004.6	(1438.7 to 2439.4)
		Hemorrhagic						
		Stroke	Male	1990	27,082	(19,460 to 35,248)	606.5	(410.0 to 826.3)
				2013	68,821	(49,035 to 91,893)	855.2	(617.7 to 1169.5)
			Female	1990	23,802	(15,918 to 45,819)	566.4	(362.5 to 1126.4)
				2013	71,146	(40,486 to 96,990)	880.2	(496.4 to 1200.2)
			Both	1990	50,884	(39,914 to 70,802)	587.1	(436.2 to 853.1)
						(103,307 to		
				2013	139,967	173,989)	872.6	(649.9 to 1086.2)
	YLDs	Ischemic Stroke	Male	1990	481	(305 to 705)	15.4	(9.6 to 22.4)
				2013	948	(581 to 1,381)	15.6	(9.7 to 22.5)
			Female	1990	379	(234 to 558)	11.6	(7.0 to 17.2)
				2013	854	(538 to 1,242)	12.1	(7.5 to 18.0)
			Both	1990	861	(565 to 1,221)	13.4	(8.7 to 18.9)
				2013	1,802	(1,162 to 2,488)	13.7	(8.8 to 18.9)
		Hemorrhagic						
		Stroke	Male	1990	156	(98 to 230)	4.1	(2.5 to 6.1)
				2013	288	(177 to 421)	3.9	(2.5 to 5.8)
			Female	1990	111	(67 to 163)	2.9	(1.8 to 4.3)
				2013	225	(143 to 336)	2.8	(1.7 to 4.2)
			Both	1990	268	(176 to 378)	3.5	(2.3 to 5.0)
			Both	2013	513	(336 to 727)	3.3	(2.1 to 4.7)
Greece	Deaths	Ischemic Stroke	Male	1990	4,786	(3,867 to 5,623)	79.6	(64.2 to 93.9)
				2013	5,211	(4,067 to 6,534)	47.9	(37.4 to 59.8)
			Female	1990	7,095	(5,347 to 8,008)	87	(65.5 to 98.2)

				2013	8,456	(6,772 to 11,754)	52.6	(42.1 to 73.5)
			Both	1990	11,882	(9,883 to 13,099)	84.1	(70.0 to 92.7)
				2013	13,667	(11,513 to 17,393)	50.9	(42.8 to 64.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	3,095	(2,456 to 3,886)	51.1	(40.4 to 64.2)
				2013	3,615	(2,534 to 4,766)	35.8	(25.5 to 46.4)
			Female	1990	4,284	(3,234 to 5,267)	54.1	(40.8 to 66.1)
				2013	4,928	(3,854 to 7,092)	33.4	(26.7 to 47.1)
			Both	1990	7,379	(6,192 to 8,708)	53.1	(44.7 to 62.6)
				2013	8,543	(6,997 to 11,029)	34.7	(28.9 to 44.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
	DALYs	Ischemic Stroke	Male	1990	61,359	(51,739 to 71,585)	1001.4	(844.8 to 1167.8)
				2013	59,181	(47,601 to 72,643)	589.8	(478.7 to 721.0)
			Female	1990	75,926	(58,020 to 84,806)	950.7	(726.0 to 1063.0)
						(62,205 to		
				2013	76,671	104,084)	526.8	(424.8 to 716.9)
						(117,212 to		
			Both	1990	137,286	151,148)	977.6	(834.5 to 1074.8)
				2042	425.052	(114,006 to	550 4	(467.5) (606.4)
		I I a wa a wwh a a i a		2013	135,853	166,832)	559.1	(467.5 to 686.1)
		Hemorrhagic Stroke	Male	1990	50,898	(42,167 to 61,320)	841.2	(696.6 to 1017.3)
		Stioke	IVIAIC	2013	49,004	(37,017 to 60,812)	556.4	(423.4 to 685.7)
			Famala		 	, , ,		,
			Female	1990	56,215	(43,542 to 67,136)	760	(593.4 to 901.1)
				2013	50,760	(41,855 to 70,496)	423.7	(355.1 to 582.7)
			Poth	1990	107 114	(93,235 to	802.5	(702.0 to 017.5)
			Both		107,114	122,859)		(702.9 to 917.5)
				2013	99,765	(84,874 to	488.7	(418.4 to 591.4)

						123,356)		
	YLDs	Ischemic Stroke	Male	1990	5,780	(3,783 to 8,190)	93.9	(61.6 to 132.6)
				2013	8,333	(5,352 to 12,088)	91.4	(59.2 to 132.6)
			Female	1990	4,417	(2,948 to 6,159)	58.4	(38.9 to 81.2)
				2013	7,027	(4,436 to 10,115)	60	(38.3 to 85.9)
			Both	1990	10,197	(6,924 to 13,769)	74.5	(50.5 to 100.6)
			50111	2013	15,361	(10,251 to 21,823)	74.3	(50.0 to 104.4)
		Hemorrhagic		2010	13,301	(10)231 (0 21)023)	7 1.3	(30.0 to 10 11 1)
		Stroke	Male	1990	801	(501 to 1,162)	13.3	(8.3 to 19.1)
				2013	1,296	(763 to 1,920)	14.8	(8.9 to 21.8)
			Female	1990	776	(488 to 1,145)	11.1	(7.0 to 16.2)
				2013	982	(591 to 1,459)	9.6	(5.8 to 14.1)
			Both	1990	1,578	(1,065 to 2,177)	12	(8.1 to 16.6)
			Both	2013	2,278	(1,478 to 3,226)	11.9	(7.8 to 16.8)
Grenada	Deaths	Ischemic Stroke	Male	1990	27	(22 to 32)	102.3	(83.4 to 121.6)
				2013	29	(23 to 35)	90.2	(73.4 to 109.9)
			Female	1990	43	(33 to 49)	109	(84.7 to 124.3)
				2013	41	(34 to 53)	83.4	(68.3 to 106.8)
			Both	1990	71	(58 to 79)	106.8	(87.8 to 118.9)
				2013	71	(61 to 85)	87.1	(75.4 to 105.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	14	(11 to 18)	49.4	(38.3 to 66.1)
				2013	16	(12 to 21)	48.4	(35.0 to 60.9)
			Female	1990	23	(18 to 29)	59.1	(46.5 to 74.3)
				2013	22	(17 to 31)	46.9	(36.4 to 64.6)
			Both	1990	37	(32 to 44)	55.3	(47.5 to 66.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

				2013	39	(31 to 48)	48.1	(39.0 to 59.4)
	DALYs	Ischemic Stroke	Male	1990	499	(406 to 588)	1660.8	(1353.5 to 1950.3)
				2013	505	(407 to 626)	1457.9	(1175.3 to 1786.0)
			Female	1990	673	(515 to 761)	1619.7	(1248.5 to 1837.6)
				2013	549	(447 to 706)	1177.6	(953.6 to 1517.7)
			Both	1990	1,172	(967 to 1,305)	1644.6	(1367.4 to 1829.3)
				2013	1,055	(909 to 1,259)	1312.7	(1126.5 to 1565.3)
		Hemorrhagic						
		Stroke	Male	1990	345	(277 to 436)	1069.3	(856.9 to 1378.0)
				2013	398	(295 to 497)	1026.6	(750.2 to 1283.0)
			Female	1990	474	(377 to 578)	1187.1	(941.3 to 1432.7)
				2013	406	(319 to 551)	896.4	(704.6 to 1215.3)
			Both	1990	819	(709 to 946)	1139.2	(989.7 to 1323.3)
				2013	804	(665 to 961)	963	(795.0 to 1150.1)
	YLDs	Ischemic Stroke	Male	1990	13	(9 to 19)	48.1	(32.1 to 66.7)
				2013	15	(10 to 22)	45.3	(29.3 to 64.8)
			Female	1990	12	(7 to 18)	30.9	(19.8 to 45.4)
				2013	14	(9 to 20)	31.3	(20.2 to 44.0)
			Both	1990	26	(18 to 36)	38	(26.3 to 52.3)
				2013	30	(20 to 41)	37.4	(25.1 to 51.8)
		Hemorrhagic						
		Stroke	Male	1990	4	(2 to 6)	14.5	(9.0 to 22.2)
				2013	5	(3 to 7)	14.4	(8.9 to 21.0)
			Female	1990	3	(2 to 6)	9.9	(6.1 to 15.3)
				2013	4	(3 to 7)	10.7	(6.6 to 15.9)
			Both	1990	8	(5 to 11)	11.9	(7.7 to 17.0)
			Both	2013	10	(6 to 14)	12.3	(7.8 to 17.3)
Guatemala	Deaths	Ischemic Stroke	Male	1990	480	(348 to 593)	34.1	(24.4 to 42.3)
				2013	624	(446 to 892)	19.1	(13.6 to 27.6)

			Female	1990	427	(323 to 595)	29.1	(22.0 to 40.6)
				2013	811	(588 to 1,124)	20.9	(15.1 to 28.9)
			Both	1990	907	(734 to 1,077)	31.5	(25.4 to 37.5)
				2013	1,435	(1,143 to 1,880)	20.1	(16.0 to 26.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	416	(329 to 570)	24.7	(19.0 to 34.6)
				2013	835	(569 to 1,108)	24	(16.1 to 32.0)
			Female	1990	420	(336 to 711)	23.7	(18.6 to 40.5)
				2013	1,184	(690 to 1,462)	28.6	(16.5 to 35.6)
			Both	1990	836	(686 to 1,213)	24.2	(19.4 to 35.4)
				2013	2,019	(1,372 to 2,431)	26.5	(17.8 to 32.2)
	DALYs	Ischemic Stroke	Male	1990	10,524	(7,600 to 13,097)	541.5	(392.9 to 668.6)
				2013	11,255	(8,273 to 15,506)	295.2	(216.7 to 405.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	8,853	(6,678 to 12,651)	444.4	(336.4 to 617.2)
				2013	13,047	(9,313 to 17,992)	299.7	(216.5 to 408.8)
			Both	1990	19,377	(15,507 to 23,221)	492.2	(399.6 to 584.6)
				2013	24,302	(19,622 to 31,740)	297.8	(240.9 to 383.6)
		Hemorrhagic						
		Stroke	Male	1990	12,374	(10,097 to 16,155)	511.9	(413.3 to 687.9)
				2013	19,796	(14,490 to 25,029)	458.1	(329.4 to 586.8)
			Female	1990	12,224	(9,870 to 20,899)	496.5	(403.5 to 834.9)
				2013	26,491	(15,947 to 31,865)	536.6	(325.9 to 643.9)
			Both	1990	24,599	(20,521 to 34,664)	504	(417.3 to 718.5)
				2013	46,287	(33,504 to 54,466)	500.6	(362.1 to 590.1)
	YLDs	Ischemic Stroke	Male	1990	110	(70 to 160)	5.8	(3.7 to 8.4)
		isoliciillo sti site	IVIGIC	1000	110	(, 0 to ±00)	5.0	(3.7 to 3)

	-							
			Female	1990	96	(64 to 139)	5.3	(3.5 to 7.6)
				2013	232	(137 to 360)	5.6	(3.3 to 8.8)
			Both	1990	207	(142 to 283)	5.6	(3.9 to 7.5)
				2013	480	(308 to 698)	6.2	(4.0 to 9.2)
		Hemorrhagic						
		Stroke	Male	1990	359	(244 to 503)	16.6	(11.1 to 23.4)
			2013	765	(485 to 1,089)	19.6	(12.5 to 27.9)	
			Female	1990	265	(175 to 378)	12.4	(8.1 to 17.8)
				2013	544	(351 to 762)	11.9	(7.5 to 16.9)
			Both	1990	624	(430 to 840)	14.5	(10.0 to 19.5)
			Both	2013	1,310	(853 to 1,806)	15.5	(10.2 to 21.4)
Guinea	Deaths	Ischemic Stroke	Male	1990	516	(303 to 704)	59.5	(35.8 to 79.6)
				2013	1,772	(1,017 to 2,401)	108.8	(64.2 to 144.4)
			Female	1990	747	(285 to 1,118)	74.2	(30.3 to 110.2)
				2013	2,058	(1,120 to 2,850)	116	(66.8 to 159.2)
			Both	1990	1,263	(820 to 1,606)	67.7	(44.1 to 86.8)
				2013	3,830	(2,576 to 4,868)	113	(76.6 to 141.3)
	Deaths	Hemorrhagic Stroke	Male	1990	599	(376 to 780)	48.7	(27.6 to 66.7)
				2013	1,125	(721 to 1,702)	50.5	(29.8 to 83.7)
			Female	1990	811	(406 to 1,140)	62.4	(31.9 to 92.6)
				2013	1,304	(768 to 2,437)	56.3	(32.5 to 114.1)
			Both	1990	1,411	(871 to 1,788)	56.3	(33.7 to 73.5)
				2013	2,430	(1,695 to 3,809)	53.7	(35.5 to 91.5)
	DALYs	Ischemic Stroke	Male	1990	12,185	(7,485 to 16,896)	960.4	(564.8 to 1309.6)
				2013	37,773	(21,476 to 52,834)	1735.9	(996.1 to 2361.0)
			Female	1990	15,895	(5,598 to 25,049)	1148.7	(444.3 to 1707.2)
				2013	39,051	(20,481 to 55,706)	1719.6	(936.6 to 2389.0)
			Both	1990	28,080	(17,729 to 36,585)	1065.3	(692.9 to 1349.2)

					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	76,824	(51,005 to 99,731)	1732.6	(1173.4 to 2202.1)
		Hemorrhagic				((
		Stroke	Male	1990	21,835	(14,079 to 28,361)	1080.6	(671.5 to 1411.0)
				2013	35,716	(24,413 to 50,494)	1107.8	(712.4 to 1669.6)
			Female	1990	24,828	(11,421 to 35,213)	1287.7	(645.9 to 1804.8)
				2013	36,489	(21,796 to 62,702)	1162.4	(697.3 to 2132.6)
			Both	1990	46,663	(28,276 to 59,373)	1194.7	(744.1 to 1514.4)
				2013	72.205	(51,926 to	1120.0	/700 7 to 1760 f\
	YLDs	Ischemic Stroke	Male	1990	72,205 215	104,757)	1138.8 15.1	(799.7 to 1768.5)
	YLDS	ischemic Stroke	iviale			(127 to 314)		(8.9 to 22.1)
			F	2013	414	(267 to 588)	15.5	(10.1 to 21.9)
			Female	1990	179	(112 to 262)	11.2	(7.0 to 16.1)
				2013	337	(212 to 499)	11.4	(7.2 to 17.1)
			Both	1990	394	(256 to 552)	13	(8.5 to 18.3)
		11		2013	751	(507 to 1,042)	13.3	(9.0 to 18.3)
		Hemorrhagic Stroke	Male	1990	68	(40 to 104)	4.1	(2.5 to 6.3)
		C . C . C		2013	124	(75 to 181)	3.9	(2.4 to 5.6)
			Female	1990	50	(31 to 74)	2.8	(1.7 to 4.2)
				2013	92	(57 to 139)	2.7	(1.7 to 4.2)
			Both	1990	119	(75 to 172)	3.4	(2.2 to 4.9)
			Both	2013	217	(139 to 306)	3.3	(2.1 to 4.6)
Guinea-Bissau	Deaths	Ischemic Stroke	Male	1990	97	(52 to 164)	66.8	(38.7 to 106.6)
				2013	335	(201 to 512)	136.4	(86.2 to 196.4)
			Female	1990	179	(102 to 312)	104.4	(62.5 to 176.4)
				2013	412	(243 to 692)	156.3	(97.9 to 250.5)

_					1	1		
			Both	1990	277	(174 to 447)	88.2	(58.3 to 133.6)
				2013	747	(487 to 1,111)	147.7	(100.8 to 211.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	97	(44 to 165)	49.3	(20.0 to 83.2)
				2013	210	(112 to 364)	60	(31.3 to 108.5)
			Female	1990	147	(65 to 249)	67.8	(27.9 to 116.3)
				2013	224	(126 to 459)	61.7	(33.7 to 132.9)
			Both	1990	244	(137 to 386)	59.6	(31.1 to 93.0)
				2013	434	(268 to 754)	61	(36.0 to 108.8)
	DALYs	Ischemic Stroke	Male	1990	2,296	(1,171 to 3,937)	1128.6	(611.9 to 1887.4)
				2013	7,772	(4,576 to 12,195)	2319.7	(1401.7 to 3500.2)
			Female	1990	3,798	(2,102 to 6,944)	1674.9	(947.0 to 2928.6)
				2013	8,631	(4,879 to 15,982)	2478.5	(1463.2 to 4194.1)
			Both	1990	6,094	(3,711 to 10,137)	1431.8	(908.9 to 2293.1)
				2013	16,403	(10,355 to 25,243)	2411.9	(1583.1 to 3570.8)
		Hemorrhagic						
		Stroke	Male	1990	3,367	(1,612 to 5,381)	1108.6	(502.1 to 1899.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	7,260	(3,982 to 11,629)	1448.2	(769.4 to 2518.7)
			Female	1990	4,488	(2,198 to 7,221)	1454	(656.9 to 2456.2)
				2013	7,072	(4,158 to 13,515)	1418.9	(818.3 to 2849.3)
			Both	1990	7,856	(4,636 to 11,863)	1295.3	(723.8 to 2052.4)
					·			
				2013	14,332	(9,445 to 23,437)	1435.2	(891.0 to 2476.7)
	YLDs	Ischemic Stroke	Male	2013 1990	14,332 35	(9,445 to 23,437) (22 to 52)	1435.2 15.6	(891.0 to 2476.7) (10.0 to 22.8)
	YLDs	Ischemic Stroke	Male			, , , , , , , , , , , , , , , , , , , ,		•
	YLDs	Ischemic Stroke	Male Female	1990	35	(22 to 52)	15.6	(10.0 to 22.8)

			1 1		I	1	1	
			Both	1990	64	(43 to 88)	13.3	(9.1 to 18.2)
				2013	110	(73 to 151)	13.4	(9.1 to 18.3)
		Hemorrhagic						
		Stroke	Male	1990	11	(6 to 16)	4.2	(2.5 to 6.1)
				2013	18	(11 to 27)	3.9	(2.4 to 5.7)
			Female	1990	8	(5 to 11)	2.9	(1.9 to 4.2)
				2013	13	(8 to 20)	2.8	(1.7 to 4.1)
			Both	1990	19	(13 to 27)	3.5	(2.3 to 4.8)
			Both	2013	32	(21 to 45)	3.3	(2.2 to 4.7)
Guyana	Deaths	Ischemic Stroke	Male	1990	288	(212 to 340)	172	(127.6 to 202.9)
				2013	164	(118 to 228)	147.5	(112.0 to 190.9)
			Female	1990	281	(204 to 319)	144.6	(105.7 to 164.0)
				2013	231	(167 to 317)	130.5	(96.7 to 173.7)
			Both	1990	569	(441 to 639)	157.7	(122.5 to 177.0)
				2013	396	(314 to 511)	138.9	(112.1 to 172.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	171	(136 to 223)	94.4	(74.5 to 125.1)
				2013	136	(92 to 187)	84.5	(55.9 to 112.2)
			Female	1990	179	(141 to 230)	89.1	(70.3 to 113.8)
				2013	162	(109 to 215)	78.1	(52.2 to 103.9)
			Both	1990	350	(301 to 433)	91.7	(78.9 to 113.9)
				2013	298	(222 to 374)	81.9	(59.7 to 101.8)
	DALYs	Ischemic Stroke	Male	1990	5,565	(3,951 to 6,565)	3082.6	(2201.4 to 3637.7)
				2013	4,068	(2,866 to 5,910)	2601.3	(1920.7 to 3500.9)
			Female	1990	4,722	(3,317 to 5,435)	2361.7	(1671.4 to 2710.3)
				2013	4,186	(2,919 to 6,147)	2037.8	(1445.9 to 2894.4)
			Both	1990	10,288	(7,797 to 11,614)	2709.3	(2052.2 to 3061.1)
				2013	8,255	(6,355 to 11,090)	2293.2	(1817.9 to 2980.4)
		Hemorrhagic	Male	1990	4,546	(3,661 to 5,760)	2174.4	(1747.0 to 2790.0)

	l l		I I		I			I
		Stroke						
				2013	4,300	(2,974 to 5,983)	1913	(1310.2 to 2591.6)
			Female	1990	4,213	(3,399 to 5,142)	1930.5	(1533.1 to 2371.0)
				2013	4,133	(2,813 to 5,651)	1637.6	(1118.9 to 2205.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	8,760	(7,637 to 10,401)	2049.8	(1773.0 to 2459.3)
				2013	8,433	(6,412 to 10,663)	1773.5	(1330.8 to 2218.5)
	YLDs	Ischemic Stroke	Male	1990	83	(53 to 118)	47.4	(29.5 to 67.5)
				2013	68	(43 to 99)	48.2	(31.0 to 68.5)
			Female	1990	65	(42 to 95)	32.7	(21.4 to 47.4)
				2013	67	(44 to 93)	33	(21.9 to 46.2)
			Both	1990	149	(101 to 208)	39.6	(26.8 to 55.3)
				2013	136	(90 to 188)	38.6	(26.1 to 53.0)
		Hemorrhagic						
		Stroke	Male	1990	29	(17 to 44)	14.8	(8.7 to 23.2)
				2013	27	(16 to 40)	13.9	(8.2 to 21.1)
			Female	1990	23	(14 to 38)	11	(6.5 to 17.7)
				2013	26	(15 to 39)	10.9	(6.5 to 16.6)
			Both	1990	52	(33 to 78)	12.8	(8.1 to 19.3)
			Both	2013	53	(35 to 76)	12	(8.0 to 17.3)
Haiti	Deaths	Ischemic Stroke	Male	1990	1,337	(976 to 1,642)	102.2	(75.1 to 123.7)
				2013	2,032	(1,596 to 2,547)	99.5	(78.3 to 126.0)
			Female	1990	2,261	(1,688 to 2,768)	150.9	(111.7 to 183.5)
				2013	3,071	(2,239 to 4,131)	122.8	(90.7 to 163.3)
			Both	1990	3,598	(2,800 to 4,250)	128.6	(101.7 to 150.2)
				2013	5,104	(4,135 to 6,299)	112.5	(92.1 to 138.4)
	Deaths	Hemorrhagic	Male	1990	1,278	(1,043 to 1,630)	72.1	(57.4 to 94.1)

		Stroke						
				2013	1,621	(1,241 to 2,043)	65.8	(49.8 to 82.7)
			Female	1990	2,091	(1,658 to 2,918)	113.9	(89.5 to 161.8)
				2013	2,891	(2,109 to 3,639)	104.8	(76.4 to 132.2)
			Both	1990	3,369	(2,809 to 4,254)	94.6	(78.2 to 123.9)
				2013	4,513	(3,692 to 5,315)	87.4	(70.5 to 103.3)
	DALYs	Ischemic Stroke	Male	1990	31,458	(22,567 to 39,226)	1851.2	(1359.9 to 2283.8)
				2013	41,916	(32,032 to 52,738)	1656.9	(1300.1 to 2079.4)
			Female	1990	49,548	(36,160 to 62,530)	2665.2	(1987.1 to 3302.3)
				2013	58,698	(41,532 to 80,945)	2011.3	(1456.8 to 2744.5)
			Both	1990	81,006	(62,186 to 96,366)	2287.4	(1787.1 to 2700.5)
					,	(80,331 to		
				2013	100,615	126,329)	1850.1	(1491.2 to 2292.0)
		Hemorrhagic						
		Stroke	Male	1990	50,986	(40,412 to 64,777)	1826.7	(1489.4 to 2313.7)
				2013	49,773	(38,078 to 63,098)	1450.1	(1100.6 to 1827.0)
			Female	1990	70,155	(54,841 to 93,597)	2685.5	(2119.4 to 3687.9)
				2013	74,530	(54,920 to 95,113)	2143.1	(1560.1 to 2720.5)
						(101,622 to		
			Both	1990	121,141	146,940)	2283.4	(1902.5 to 2853.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(101,740 to		
				2013	124,304	146,575)	1823.7	(1483.7 to 2154.4)
	YLDs	Ischemic Stroke	Male	1990	675	(423 to 966)	45.9	(28.6 to 65.6)
				2013	1,091	(690 to 1,550)	45.8	(29.2 to 65.1)
			Female	1990	536	(358 to 753)	33	(22.0 to 46.3)
				2013	912	(582 to 1,269)	32.8	(20.7 to 46.0)

			Both	1990	1,212	(831 to 1,683)	39	(26.6 to 54.1)
				2013	2,004	(1,353 to 2,777)	38.7	(26.3 to 53.6)
		Hemorrhagic						
		Stroke	Male	1990	255	(150 to 405)	15	(8.6 to 24.1)
				2013	397	(243 to 586)	14.2	(8.8 to 21.4)
			Female	1990	182	(120 to 261)	9.8	(6.3 to 14.6)
				2013	321	(200 to 459)	10.2	(6.3 to 14.9)
			Both	1990	437	(285 to 636)	12.2	(8.0 to 18.0)
			Both	2013	719	(479 to 1,000)	12	(8.0 to 16.8)
Honduras	Deaths	Ischemic Stroke	Male	1990	498	(386 to 731)	63.5	(49.4 to 93.8)
				2013	1,169	(784 to 1,811)	66.6	(45.1 to 101.8)
			Female	1990	575	(443 to 729)	64.3	(49.6 to 80.9)
				2013	1,360	(849 to 1,900)	65.4	(40.9 to 91.0)
			Both	1990	1,073	(914 to 1,382)	64.1	(54.6 to 81.5)
				2013	2,529	(1,866 to 3,339)	66.3	(49.2 to 87.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	245	(173 to 415)	22	(15.2 to 37.3)
				2013	466	(287 to 863)	22.5	(13.8 to 41.1)
			Female	1990	308	(220 to 407)	26.3	(18.7 to 35.2)
				2013	606	(361 to 965)	26.7	(16.0 to 42.8)
			Both	1990	554	(436 to 776)	24.3	(19.1 to 33.7)
				2013	1,073	(743 to 1,607)	24.8	(17.2 to 36.5)
	DALYs	Ischemic Stroke	Male	1990	11,430	(8,585 to 17,356)	1040.3	(800.7 to 1520.9)
				2013	21,067	(13,681 to 33,784)	1034.7	(672.4 to 1617.0)
			Female	1990	11,879	(8,880 to 15,456)	999.3	(769.7 to 1286.1)
				2013	20,563	(12,740 to 29,634)	917.4	(574.4 to 1314.1)
			Both	1990	23,309	(19,179 to 31,431)	1020.4	(858.2 to 1338.5)
				2013	41,631	(30,074 to 57,664)	976.9	(708.0 to 1332.0)
		Hemorrhagic	Male	1990	9,988	(7,049 to 17,182)	573.6	(404.9 to 965.7)

		Stroke						
				2013	13,827	(8,698 to 26,716)	522.8	(327.6 to 961.6)
			Female	1990	11,450	(7,837 to 15,297)	678.3	(480.1 to 889.1)
				2013	15,570	(9,283 to 24,743)	578.5	(342.9 to 912.0)
			Both	1990	21,438	(16,505 to 30,677)	626.8	(490.1 to 871.4)
				2013	29,397	(19,880 to 45,330)	552.4	(378.3 to 836.9)
	YLDs	Ischemic Stroke	Male	1990	69	(43 to 102)	7.4	(4.6 to 10.9)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	161	(102 to 234)	8	(5.1 to 11.5)
			Female	1990	58	(36 to 84)	5.9	(3.6 to 8.7)
				2013	147	(85 to 218)	6.7	(3.9 to 10.0)
			Both	1990	128	(83 to 183)	6.6	(4.3 to 9.4)
				2013	308	(201 to 424)	7.4	(4.8 to 10.1)
		Hemorrhagic						
		Stroke	Male	1990	207	(136 to 293)	19.1	(12.5 to 27.6)
				2013	457	(299 to 636)	20.1	(13.0 to 27.8)
			Female	1990	142	(91 to 205)	12.6	(8.0 to 18.3)
				2013	302	(196 to 430)	12.3	(8.1 to 17.8)
			Both	1990	350	(232 to 475)	15.7	(10.4 to 21.7)
			Both	2013	759	(514 to 1,035)	16.1	(10.9 to 21.7)
Hungary	Deaths	Ischemic Stroke	Male	1990	7,053	(5,977 to 8,279)	145.4	(123.3 to 171.2)
				2013	5,938	(4,829 to 7,119)	96.4	(78.5 to 115.6)
			Female	1990	9,189	(7,506 to 10,820)	109.5	(89.2 to 129.0)
				2013	7,870	(6,314 to 10,727)	62.8	(50.2 to 85.2)
			Both	1990	16,242	(14,649 to 18,411)	124	(111.5 to 140.8)
				2013	13,809	(11,818 to 16,752)	75.5	(64.5 to 91.3)
	Deaths	Hemorrhagic	Male	1990	3,186	(2,333 to 3,894)	61	(44.1 to 74.4)

		Stroke						
				2013	2,030	(1,522 to 2,925)	32.3	(24.2 to 46.7)
			Female	1990	3,936	(2,475 to 4,530)	49.1	(31.3 to 56.2)
				2013	2,113	(1,452 to 3,952)	19.6	(13.9 to 36.1)
			Both	1990	7,123	(5,324 to 8,134)	54.5	(40.6 to 62.3)
				2013	4,143	(3,129 to 6,251)	24.9	(19.1 to 37.0)
						(97,691 to		·
	DALYs	Ischemic Stroke	Male	1990	114,967	135,970)	2186.7	(1852.3 to 2578.7)
						(67,556 to		
				2013	83,968	100,490)	1331.3	(1077.7 to 1591.5)
						(93,363 to		
			Female	1990	115,682	133,764)	1413.1	(1137.4 to 1643.9)
						(63,215 to		
				2013	78,970	107,424)	705.9	(561.7 to 962.1)
						(205,632 to		
			Both	1990	230,650	261,246)	1737.4	(1551.3 to 1970.9)
				2042	462.000	(139,151 to	0505	(045.7)
		11		2013	162,938	195,136)	958.5	(815.7 to 1142.2)
		Hemorrhagic	Mala	1000	71.046	/F 4 72 4 + 6 07 F 70\	1200 7	(002.0 to 1570.2)
		Stroke	Male	1990	71,946	(54,734 to 87,579)	1298.7	(982.0 to 1578.3)
				2013	42,348	(33,055 to 57,071)	671.6	(525.4 to 902.1)
			Female	1990	68,690	(45,945 to 76,925)	923.5	(629.0 to 1027.6)
				2013	31,659	(23,660 to 56,253)	357.6	(275.1 to 612.4)
						(110,916 to		(
			Both	1990	140,637	159,718)	1093.9	(868.1 to 1243.2)
					Counts, All	Ages	Rate	Per 100,000
		C: 1 =				050(11)	Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2012	74.000	(59,357 to	406.4	(404 4 to CC4 7)
				2013	74,008	101,744)	496.4	(404.4 to 664.7)
	YLDs	Ischemic Stroke	Male	1990	1,406	(890 to 1,972)	25.5	(16.3 to 35.8)

				2013	1,730	(1,140 to 2,440)	27.5	(18.2 to 38.7)
			Female	1990	1,377	(867 to 1,957)	18.4	(11.7 to 26.2)
				2013	1,504	(973 to 2,146)	16.8	(10.9 to 23.7)
			Both	1990	2,784	(1,836 to 3,752)	21.6	(14.2 to 29.1)
				2013	3,235	(2,215 to 4,478)	21.5	(14.7 to 29.7)
		Hemorrhagic						
		Stroke	Male	1990	296	(173 to 435)	5.4	(3.2 to 7.9)
				2013	368	(220 to 569)	6.1	(3.6 to 9.2)
			Female	1990	297	(183 to 446)	4.3	(2.7 to 6.3)
				2013	352	(210 to 532)	4.5	(2.7 to 6.7)
			Both	1990	594	(387 to 841)	4.8	(3.2 to 6.8)
			Both	2013	720	(454 to 1,037)	5.2	(3.3 to 7.4)
Iceland	Deaths	Ischemic Stroke	Male	1990	44	(35 to 57)	38.5	(30.5 to 50.0)
				2013	47	(33 to 62)	23.1	(16.2 to 30.2)
			Female	1990	56	(45 to 73)	34.2	(27.0 to 44.0)
				2013	58	(42 to 76)	19.7	(14.3 to 25.5)
			Both	1990	101	(87 to 122)	36.3	(31.4 to 43.7)
				2013	105	(82 to 126)	21.4	(16.7 to 25.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	29	(22 to 39)	25.5	(19.2 to 34.1)
				2013	32	(21 to 43)	16	(10.7 to 21.2)
			Female	1990	42	(31 to 56)	27.7	(20.4 to 35.9)
				2013	37	(27 to 52)	13.9	(10.1 to 19.2)
			Both	1990	72	(59 to 89)	26.7	(22.1 to 32.9)
				2013	69	(52 to 87)	14.9	(11.5 to 18.6)
	DALYs	Ischemic Stroke	Male	1990	632	(522 to 778)	538.5	(445.0 to 663.7)
				2013	650	(493 to 823)	331.2	(252.8 to 418.2)
			Female	1990	617	(506 to 745)	402	(331.3 to 487.3)
				2013	551	(424 to 682)	219.6	(169.3 to 269.5)

			Both	1990	1,249	(1,095 to 1,451)	465.9	(408.8 to 540.7)
				2013	1,202	(978 to 1,428)	273	(221.3 to 325.0)
		Hemorrhagic						
		Stroke	Male	1990	516	(411 to 653)	429.3	(340.3 to 544.0)
				2013	467	(321 to 605)	242.9	(166.9 to 313.9)
			Female	1990	625	(481 to 780)	452.8	(348.3 to 563.3)
				2013	422	(318 to 549)	189.1	(143.5 to 244.5)
			Both	1990	1,142	(974 to 1,347)	441	(376.5 to 518.2)
				2013	889	(704 to 1,073)	214.7	(170.1 to 258.1)
	YLDs	Ischemic Stroke	Male	1990	101	(65 to 144)	85.6	(54.8 to 121.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	170	(111 to 241)	88	(57.7 to 124.4)
			Female	1990	93	(60 to 130)	66.6	(43.3 to 92.1)
				2013	131	(84 to 194)	60.1	(38.7 to 88.8)
			Both	1990	195	(134 to 264)	75.3	(51.5 to 102.4)
				2013	302	(203 to 412)	73.4	(49.5 to 99.8)
		Hemorrhagic						
		Stroke	Male	1990	19	(10 to 29)	15.9	(8.9 to 24.3)
				2013	34	(21 to 52)	18.4	(11.4 to 27.2)
			Female	1990	15	(9 to 22)	11.1	(6.9 to 16.5)
				2013	23	(14 to 35)	11.3	(7.0 to 17.3)
			Both	1990	34	(22 to 49)	13.3	(8.6 to 19.2)
			Both	2013	58	(38 to 84)	14.6	(9.6 to 21.1)
						(57,503 to		
India	Deaths	Ischemic Stroke	Male	1990	89,777	120,555)	55	(35.2 to 72.8)
						(113,526 to		,
				2013	167,039	225,568)	56.4	(39.0 to 75.1)
			Female	1990	89,957	(53,931 to	57.9	(35.5 to 77.4)

			T			125 476)		
						125,476)		
						(104,964 to		
				2013	165,730	221,959)	51	(32.6 to 67.8)
						(131,089 to		
			Both	1990	179,734	223,609)	56.6	(41.6 to 69.6)
						(249,102 to		
				2013	332,769	409,620)	53.7	(40.2 to 65.2)
		Hemorrhagic				(82,615 to		
	Deaths	Stroke	Male	1990	115,363	155,523)	54.4	(36.8 to 78.0)
						(141,893 to		
				2013	198,257	289,499)	53.4	(37.9 to 79.2)
						(73,812 to		
			Female	1990	118,482	155,048)	58.9	(36.4 to 79.8)
						(119,426 to		
				2013	183,187	253,800)	48.1	(31.3 to 67.6)
						(179,841 to		
			Both	1990	233,845	289,583)	56.7	(42.9 to 72.1)
						(294,382 to		
				2013	381,445	487,330)	50.8	(39.1 to 65.7)
						(1,190,035 to		
	DALYs	Ischemic Stroke	Male	1990	1,887,041	2,592,045)	903.5	(583.2 to 1205.7)
						(2,167,198 to		
				2013	3,217,138	4,437,095)	880.8	(599.7 to 1187.7)
						(981,536 to		
			Female	1990	1,716,614	2,655,931)	870.6	(524.9 to 1237.9)
						(1,700,216 to		
				2013	2,680,670	3,640,302)	708.8	(454.8 to 945.2)
						(2,639,063 to		
			Both	1990	3,603,655	4,599,668)	888.5	(649.3 to 1106.7)
			·		Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
,		, , , , , ,						

				•			
					(4,426,760 to		
			2013	5,897,809	7,399,023)	793.4	(600.3 to 981.6)
	Hemorrhagic				(2,576,937 to		
	Stroke	Male	1990	3,416,647	4,466,161)	1225.5	(891.0 to 1623.8)
					(3,867,429 to		
			2013	5,305,921	7,568,556)	1142.5	(827.3 to 1658.2)
					(2,146,906 to		
		Female	1990	3,310,584	4,242,407)	1259.2	(794.9 to 1626.8)
					(2,738,256 to		
			2013	4,200,156	5,734,171)	930.3	(608.7 to 1269.2)
					(5,311,589 to		
		Both	1990	6,727,232	8,072,971)	1243	(961.2 to 1525.7)
					(7,431,973 to		
			2013	9,506,078	12,042,460)	1038	(809.3 to 1319.0)
YLDs	Ischemic Stroke	Male	1990	70,213	(47,030 to 97,677)	27.8	(18.7 to 38.4)
					(80,386 to		
			2013	126,656	176,987)	28.2	(17.9 to 39.5)
		Female	1990	57,816	(38,130 to 80,842)	23.4	(15.4 to 32.8)
					(68,805 to		•
			2013	111,072	157,682)	24.1	(15.1 to 34.2)
					(88,055 to		
		Both	1990	128,029	173,850)	25.6	(17.8 to 34.7)
					(158,111 to		
			2013	237,729	323,368)	26.1	(17.3 to 35.3)
	Hemorrhagic						
	Stroke	Male	1990	39,810	(25,980 to 55,947)	14.5	(9.4 to 20.4)
					(46,557 to		
			2013	71,803	101,671)	15.1	(9.8 to 21.2)
		Female	1990	29,115	(18,813 to 40,699)	10.9	(7.0 to 15.2)
			2013	55,638	(34,843 to 81,003)	11.5	(7.3 to 16.7)
	YLDs	Stroke Stroke Hemorrhagic	Stroke Male Female Both YLDs Ischemic Stroke Male Female Hemorrhagic Stroke Male	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 3,416,647 2013 5,305,921 2013 5,305,921 Female 1990 3,310,584 2013 4,200,156 Both 1990 6,727,232 YLDs Ischemic Stroke Male 1990 70,213 YLDs Ischemic Stroke Male 1990 57,816 Female 1990 57,816 111,072 Both 1990 128,029 Hemorrhagic Stroke Male 1990 39,810 Hemorrhagic Stroke Male 1990 29,115	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 3,416,647 4,466,161 1225.5 (2,576,937 to 4,466,161 1225.5 (3,867,429 to 7,568,556 1142.5 (2,146,906 to 4,242,407 1259.2 (2,738,256 to 4,200,156 5,734,171 930.3 (2,738,256 to 5,734,171 930.3 (2,738,256 to 6,727,232 8,072,971 1243 (7,431,973 to 12,042,460) 1038 (7,431,973 to 12,042,460) 1038 (7,431,973 to 12,042,460) 1038 (80,386 to 12,042,460) 1038 (80,386 to 12,042,460) 1038 (80,386 to 176,987) 28.2 (68,805 to 111,072 157,682) 24.1 (158,111 to 2013 237,729 323,368) 26.1 (158,111 to 2013 71,803 101,671) 15.1 (46,557 to 10,987) 14.5 (46,557 to 10,987) 15.1 (18,813 to 40,699) 10.9

						(83,822 to		
			Both	2013	127,442	177,996)	13.3	(8.8 to 18.3)
Indonesia	Deaths	Ischemic Stroke	Male	1990	28,053	(19,459 to 37,159)	101	(71.1 to 131.0)
				2013	68,619	(53,251 to 83,285)	125.3	(97.3 to 150.6)
			Female	1990	29,351	(22,506 to 36,438)	88.5	(66.4 to 109.2)
				2013	84,222	(60,366 to 99,862)	127.1	(91.1 to 150.3)
			Both	1990	57,404	(47,052 to 69,454)	94.2	(76.8 to 113.2)
						(123,946 to		,
				2013	152,841	174,085)	127	(102.3 to 144.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	34,417	(24,458 to 43,843)	93.2	(62.6 to 122.1)
				2013	65,441	(50,479 to 83,627)	88.8	(67.7 to 117.4)
			Female	1990	36,731	(29,443 to 47,173)	88.5	(67.0 to 113.0)
				2013	71,634	(55,254 to 93,242)	89.6	(68.7 to 119.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	71,148	(57,760 to 85,225)	90.7	(70.7 to 109.9)
						(115,276 to		
				2013	137,075	165,434)	89.7	(74.5 to 110.9)
						(366,977 to		
	DALYs	Ischemic Stroke	Male	1990	539,362	738,598)	1495.4	(1039.8 to 1983.7)
						(941,738 to		
				2013	1,251,249	1,582,713)	1829.1	(1408.6 to 2234.4)
				4000	505 700	(399,365 to	4000.0	(076.4) 4550.0
			Female	1990	505,700	652,093)	1233.2	(976.1 to 1550.2)
			† †			/01 / 02 F +c		
				2012	1 270 217	(914,035 to	1670 0	(1202 1 to 2011 2)
				2013	1,279,217	1,549,819)	1678.9	(1203.1 to 2011.2)
			Both			1,549,819) (849,322 to		
			Both	2013 1990 2013	1,279,217 1,045,062 2,530,467	1,549,819)	1678.9 1353.6 1756.8	(1203.1 to 2011.2) (1112.2 to 1643.5) (1419.3 to 2002.6)

						2,900,622)		
		Hemorrhagic				(715,370 to		
		Stroke	Male	1990	960,012	1,193,385)	1948.5	(1398.8 to 2454.9)
					,	(1,387,482 to		,
				2013	1,807,866	2,244,083)	1935.4	(1504.2 to 2429.1)
						(772,339 to		
			Female	1990	924,099	1,208,573)	1775.8	(1464.7 to 2287.9)
						(1,316,679 to		
				2013	1,694,594	2,137,775)	1779.6	(1386.6 to 2275.9)
						(1,594,844 to		
			Both	1990	1,884,112	2,234,051)	1858.5	(1541.4 to 2204.4)
						(2,947,099 to		
				2013	3,502,461	4,107,028)	1860.8	(1568.1 to 2221.4)
	YLDs	Ischemic Stroke	Male	1990	10,730	(7,120 to 15,061)	28.2	(18.7 to 39.5)
				2013	20,430	(12,300 to 30,428)	28.2	(16.8 to 42.6)
			Female	1990	9,788	(6,196 to 14,463)	22.5	(13.9 to 33.7)
				2013	20,013	(13,135 to 28,110)	24.6	(15.7 to 34.7)
			Both	1990	20,519	(13,670 to 28,171)	25.1	(16.7 to 34.8)
				2013	40,443	(26,972 to 55,704)	26.3	(17.2 to 36.3)
		Hemorrhagic						
		Stroke	Male	1990	13,335	(8,682 to 18,372)	30.9	(20.2 to 42.5)
				2013	24,736	(15,668 to 35,572)	30.5	(19.1 to 44.2)
			Female	1990	10,656	(6,791 to 15,155)	22.3	(14.1 to 32.5)
				2013	20,194	(13,254 to 28,600)	23.1	(15.0 to 32.9)
			Both	1990	23,991	(16,257 to 32,426)	26.3	(17.7 to 35.8)
			Both	2013	44,931	(29,796 to 61,998)	26.6	(17.6 to 36.7)
Iran	Deaths	Ischemic Stroke	Male	1990	6,621	(5,006 to 8,817)	80.9	(63.1 to 105.0)
				2013	7,479	(3,869 to 10,185)	37.4	(19.6 to 50.8)
			Female	1990	4,808	(3,855 to 6,038)	67.8	(54.1 to 84.4)
				2013	8,914	(5,299 to 11,122)	47	(28.0 to 58.4)

			Both	1990	11,429	(9,510 to 13,843)	74.6	(63.7 to 88.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	16,394	(10,036 to 19,890)	42	(25.7 to 50.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	6,721	(4,941 to 8,725)	61.6	(43.7 to 80.2)
				2013	5,745	(3,647 to 9,275)	26	(15.9 to 43.3)
			Female	1990	5,402	(4,053 to 6,582)	54.6	(41.2 to 69.3)
				2013	6,311	(3,343 to 8,180)	30	(15.8 to 39.0)
			Both	1990	12,123	(10,036 to 14,408)	58.2	(47.7 to 70.2)
				2013	12,057	(8,700 to 15,439)	27.9	(20.0 to 36.5)
						(95,006 to		
	DALYs	Ischemic Stroke	Male	1990	131,239	185,023)	1177.1	(890.7 to 1575.3)
						(52,722 to		
				2013	105,692	149,102)	465.4	(239.6 to 650.7)
						(67,103 to		
			Female	1990	90,575	125,490)	884.4	(710.7 to 1124.7)
						(64,346 to		
				2013	111,328	142,182)	506.6	(303.1 to 639.7)
			D - +l-	1000	224 045	(175,013 to	4025.5	/050 7 to 4202 7\
			Both	1990	221,815	281,609)	1035.5	(858.7 to 1262.7)
				2013	217,021	(127,286 to 269,256)	484.8	(290.9 to 592.3)
		Hemorrhagic		2013	217,021	(157,002 to	404.0	(230.3 (0 332.3)
		Stroke	Male	1990	207,657	273,236)	1248.9	(918.6 to 1627.1)
		or one	111010		207,007	(74,834 to		(310.0 to 1027.1)
				2013	115,921	169,769)	430.9	(275.2 to 660.4)
					,	(124,423 to		
			Female	1990	177,222	215,446)	1032	(779.6 to 1269.8)
				2013	113,264	(65,498 to	432.3	(242.8 to 555.2)

						146,914)		
						(316,529 to		
			Both	1990	384,879	455,210)	1143.3	(940.3 to 1358.4)
						(172,356 to		
				2013	229,185	282,547)	431	(319.2 to 536.8)
	YLDs	Ischemic Stroke	Male	1990	1,239	(745 to 1,807)	8.6	(5.1 to 12.7)
				2013	3,515	(2,317 to 4,907)	12.6	(8.5 to 17.5)
			Female	1990	930	(596 to 1,355)	6.9	(4.4 to 10.1)
				2013	2,793	(1,842 to 3,873)	9.9	(6.7 to 13.7)
			Both	1990	2,169	(1,435 to 3,065)	7.8	(5.2 to 11.0)
				2013	6,309	(4,313 to 8,643)	11.3	(7.9 to 15.4)
		Hemorrhagic						
		Stroke	Male	1990	268	(167 to 402)	1.6	(1.0 to 2.4)
				2013	649	(415 to 945)	2.1	(1.4 to 3.1)
			Female	1990	319	(201 to 456)	2	(1.3 to 2.9)
				2013	823	(537 to 1,182)	2.6	(1.7 to 3.7)
			Both	1990	587	(384 to 834)	1.8	(1.2 to 2.6)
			Both	2013	1,473	(980 to 2,059)	2.4	(1.6 to 3.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Iraq	Deaths	Ischemic Stroke	Male	1990	2,870	(2,108 to 3,763)	101.8	(75.8 to 130.5)
				2013	4,098	(2,841 to 5,685)	90.6	(64.0 to 123.1)
			Female	1990	3,479	(2,608 to 4,504)	104.6	(78.8 to 132.9)
				2013	5,888	(4,207 to 7,493)	100.1	(71.9 to 126.5)
			Both	1990	6,350	(5,252 to 7,640)	103.5	(86.1 to 123.0)
				2013	9,987	(7,732 to 12,064)	96.5	(75.9 to 115.7)
		Hemorrhagic				,		,
	Deaths	Stroke	Male	1990	1,994	(1,436 to 2,647)	53.3	(36.4 to 73.8)
				2013	2,993	(2,042 to 4,380)	49.2	(32.7 to 72.4)

					T.		
		Female	1990	2,495	(1,785 to 3,461)	65.7	(46.6 to 90.4)
			2013	4,417	(3,121 to 6,190)	63.7	(45.5 to 88.1)
		Both	1990	4,489	(3,549 to 5,665)	60.1	(46.3 to 76.6)
			2013	7,411	(5,739 to 9,413)	57.6	(45.0 to 72.9)
DALYs	Ischemic Stroke	Male	1990	50,293	(35,675 to 68,658)	1489.1	(1080.4 to 1973.1)
				,	(51,518 to		,
			2013	76,335	109,237)	1324.7	(915.3 to 1854.6)
		Female	1990	56,122	(41,239 to 75,808)	1474.3	(1092.2 to 1955.2)
					(65,991 to		
			2013	94,128	123,406)	1386.3	(980.6 to 1805.5)
					(85,144 to		
		Both	1990	106,415	130,888)	1483.1	(1198.4 to 1811.6)
					(128,978 to		
			2013	170,463	210,749)	1365.1	(1045.8 to 1667.6)
	Hemorrhagic						
	Stroke	Male	1990	67,907	(48,864 to 89,282)	1164.8	(817.3 to 1584.7)
					(64,074 to		
			2013	92,226	131,411)	1017.3	(685.7 to 1510.0)
		Female	1990	64,050	(46,069 to 92,515)	1339.8	(954.2 to 1876.4)
					(74,806 to		
			2013	108,314	158,285)	1237.8	(850.4 to 1772.7)
					(104,669 to		
		Both	1990	131,957	166,397)	1259	(986.3 to 1598.8)
					(154,883 to		(
			2013	200,541	254,940)	1140.6	(870.9 to 1461.6)
YLDs	Ischemic Stroke	Male	1990	370	(232 to 543)	9.2	(5.8 to 13.4)
			2013	719	(479 to 1,030)	9.2	(6.2 to 13.0)
		Female	1990	301	(191 to 421)	7.1	(4.5 to 10.0)
			2013	610	(396 to 861)	7.1	(4.7 to 9.9)
		Both	1990	672	(439 to 942)	8.1	(5.4 to 11.3)

measure
Deaths I
Deaths
DALYs I
Deaths

				2013	17,357	(14,538 to 21,520)	305.7	(256.5 to 377.4)
		Hemorrhagic						
		Stroke	Male	1990	9,536	(7,664 to 12,610)	576.3	(464.6 to 759.1)
				2013	7,899	(5,666 to 10,007)	310.1	(223.6 to 392.7)
			Female	1990	11,193	(9,009 to 14,367)	564	(455.9 to 719.4)
				2013	7,940	(5,873 to 10,206)	267.3	(199.8 to 344.1)
			Both	1990	20,729	(18,309 to 25,115)	569.5	(503.3 to 688.5)
				2013	15,839	(12,618 to 18,568)	287.6	(229.7 to 337.6)
	YLDs	Ischemic Stroke	Male	1990	1,514	(975 to 2,140)	91.1	(58.6 to 128.6)
				2013	2,266	(1,436 to 3,288)	88.2	(55.9 to 128.4)
			Female	1990	1,310	(858 to 1,868)	63.2	(41.6 to 89.9)
				2013	1,796	(1,179 to 2,563)	60.4	(39.5 to 86.1)
			Both	1990	2,824	(1,866 to 3,840)	75.7	(50.0 to 103.1)
				2013	4,062	(2,678 to 5,606)	73.5	(48.5 to 101.4)
		Hemorrhagic						
		Stroke	Male	1990	290	(170 to 457)	17.5	(10.3 to 27.2)
				2013	452	(214 to 727)	17.6	(8.5 to 28.4)
			Female	1990	269	(136 to 500)	13.6	(7.1 to 23.9)
				2013	381	(213 to 612)	13.3	(7.5 to 21.1)
			Both	1990	560	(345 to 841)	15.3	(9.6 to 22.7)
			Both	2013	833	(514 to 1,219)	15.3	(9.5 to 22.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Israel	Deaths	Ischemic Stroke	Male	1990	723	(574 to 924)	41.8	(33.1 to 53.4)
				2013	526	(321 to 709)	13.7	(8.4 to 18.4)
			Female	1990	855	(644 to 1,044)	40.5	(30.4 to 49.6)
				2013	1,000	(557 to 1,360)	17.3	(9.8 to 23.4)
			Both	1990	1,579	(1,343 to 1,848)	41.1	(35.0 to 48.2)

			2013	1,526	(968 to 1,892)	16	(10.1 to 19.8)
	Hemorrhagic						
Deaths	Stroke	Male	1990	514	(406 to 774)	29.5	(23.3 to 44.6)
			2013	481	(260 to 660)	12.7	(6.9 to 17.4)
		Female	1990	608	(491 to 843)	29.1	(23.4 to 40.4)
			2013	671	(335 to 957)	12.4	(6.4 to 17.5)
		Both	1990	1,123	(976 to 1,497)	29.3	(25.5 to 39.1)
			2013	1,153	(703 to 1,457)	12.6	(7.8 to 15.9)
DALYs	Ischemic Stroke	Male	1990	10,655	(8,604 to 13,305)	602.6	(485.7 to 751.3)
			2013	7,825	(5,479 to 9,801)	210.1	(147.3 to 264.0)
		Female	1990	11,139	(8,646 to 13,336)	523.1	(406.8 to 622.1)
			2013	10,124	(6,572 to 13,146)	199.4	(132.2 to 260.1)
		Both	1990	21,794	(18,798 to 25,246)	560.3	(482.7 to 646.7)
			2013	17,950	(12,534 to 21,673)	206	(145.6 to 249.1)
	Hemorrhagic						
	Stroke	Male	1990	9,282	(7,417 to 13,372)	522.6	(417.2 to 756.9)
			2013	7,344	(4,238 to 9,471)	198.2	(114.6 to 254.6)
		Female	1990	9,623	(7,856 to 12,930)	464.7	(377.8 to 622.6)
			2013	7,235	(4,187 to 10,186)	153.1	(89.6 to 218.1)
		Both	1990	18,906	(16,491 to 24,770)	492.4	(429.3 to 646.1)
			2013	14,579	(9,769 to 17,707)	174.7	(119.5 to 212.2)
YLDs	Ischemic Stroke	Male	1990	1,044	(704 to 1,419)	58.7	(39.4 to 80.0)
			2013	2,274	(1,485 to 3,187)	61.5	(40.1 to 86.1)
		Female	1990	1,151	(747 to 1,601)	55.3	(35.8 to 76.9)
			2013	2,236	(1,524 to 3,082)	48.7	(33.3 to 66.6)
		Both	1990	2,196	(1,471 to 2,991)	56.8	(38.0 to 77.5)
			2013	4,511	(3,102 to 6,075)	54.5	(37.9 to 73.1)
	Hemorrhagic Stroke	Male	1990	204	(121 to 206)	11.2	(7.2 to 15.0)
	Struke	iviale	1990	204	(131 to 286)	11.2	(7.3 to 15.9)

Country Reasure Stroke Type Sex Year Male 1990 11,229 12,224 to 33,9 to 37,197 24,6 (19,1 to 35.2)									
Both 1990 319 (212 to 437) 8.2 (3.3 to 7.4)					2013	445	(282 to 636)	12	(7.6 to 17.3)
Both 1990 319 (212 to 437) 8.2 (5.4 to 11.2)				Female	1990	115	(73 to 165)	5.6	(3.5 to 8.0)
Both 2013 670 (448 to 918) 8.3 (5.6 to 11.3)					2013	225	(146 to 324)	5.2	(3.3 to 7.4)
Italy Deaths Ischemic Stroke Male 1990 20,181 (16,326 to 23,775) 64.7 (52.2 to 76.0)				Both	1990	319	(212 to 437)	8.2	(5.4 to 11.2)
Country Measure Stroke Type Sex Year Mean 95% U Mean 95% U (11,861 to 20,108) 26.2 (19.6 to 32.9)				Both	2013	670	(448 to 918)	8.3	(5.6 to 11.3)
Female 1990 29,819 (22,224 to 33,658) 54.1 (40.5 to 60.7)	Italy	Deaths	Ischemic Stroke	Male	1990	20,181	(16,326 to 23,775)	64.7	(52.2 to 76.0)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI standardized Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Location 2013 26,278 (20,339 to 37,197) 24.6 (19.1 to 35.2) (19.1 to 35.2) Location Both 1990 50,000 (41,530 to 55,259) 58.5 (48.6 to 64.7) Location 2013 42,248 (35,128 to 54,098) 25.5 (21.3 to 32.7) Location Hemorrhagic 2013 11,299 (9,026 to 14,258) 34.7 (27.9 to 44.3) Location Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) Location Stroke Male 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Location Both 1990 26,096 (22,181 to 31,375) 31.6 (26,9 to 38.0) Location Both 1990 275,865					2013	15,970	(11,861 to 20,108)	26.2	(19.6 to 32.9)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Lountry Measure Stroke Type sex year Mean 95% UI Mean 95% UI Lountry Measure 2013 26,278 (20,339 to 37,197) 24.6 (19.1 to 35.2) Lountry Lountry Both 1990 50,000 (41,530 to 55,259) 58.5 (48.6 to 64.7) Lountry Hemorrhagic 2013 42,248 (35,128 to 54,098) 25.5 (21.3 to 32.7) Lountry Hemorrhagic Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) Lountry Email 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Lountry Female 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) Lountry Lountry 2013 28,308 (20,401 to 34,724) 19.1 (14.2 to 23.1) Lountry Lountry 183,571 <td< td=""><td></td><td></td><td></td><td>Female</td><td>1990</td><td>29,819</td><td>(22,224 to 33,658)</td><td>54.1</td><td>(40.5 to 60.7)</td></td<>				Female	1990	29,819	(22,224 to 33,658)	54.1	(40.5 to 60.7)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI 2013 26,278 (20,339 to 37,197) 24.6 (19.1 to 35.2) Both 1990 50,000 (41,530 to 55,259) 58.5 (48.6 to 64.7) Lemorrhagic 2013 42,248 (35,128 to 54,098) 25.5 (21.3 to 32.7) Deaths Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) Deaths Stroke Male 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Female 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Both 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) DALYs Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) DALYs Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) Female						Counts, All	Ages	Rate	Per 100,000
2013 26,278 (20,339 to 37,197) 24.6 (19.1 to 35.2)								Age-	standardized
Both 1990 50,000 (41,530 to 55,259) 58.5 (48.6 to 64.7)	Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Deaths Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) Deaths Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) Deaths Stroke Male 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Deaths Eemale 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) Deaths Both 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) DALys Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) DALys Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) DALys Ischemic Stroke Male 1990 301,820 335,568) 584.4 (443.5 to 650.3) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 20,356) 230.8 (188.6 to 326.2)					2013	26,278	(20,339 to 37,197)	24.6	(19.1 to 35.2)
Hemorrhagic Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3)				Both	1990	50,000	(41,530 to 55,259)	58.5	(48.6 to 64.7)
Deaths Stroke Male 1990 11,229 (9,026 to 14,258) 34.7 (27.9 to 44.3) 2013 11,899 (7,687 to 15,528) 20.9 (13.8 to 27.0) Female 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) 2013 16,409 (10,882 to 22,178) 17.5 (12.4 to 23.4) Both 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) 2013 28,308 (20,401 to 34,724) 19.1 (14.2 to 23.1) DALYs Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) 1 2013 183,571 223,323) 324.6 (258.7 to 394.5) 2 (228,800 to 301,820 335,568) 584.4 (443.5 to 650.3) 3 (167,359 to 301,820) 290,356) 230.8 (188.6 to 326.2)					2013	42,248	(35,128 to 54,098)	25.5	(21.3 to 32.7)
2013 11,899 (7,687 to 15,528) 20.9 (13.8 to 27.0)									
Female 1990 14,867 (11,219 to 18,709) 29.1 (22.2 to 36.1) 2013 16,409 (10,882 to 22,178) 17.5 (12.4 to 23.4) Both 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) 2013 28,308 (20,401 to 34,724) 19.1 (14.2 to 23.1) DALYS Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) 2013 183,571 223,323) 324.6 (258.7 to 394.5) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 200,356) 230.8 (188.6 to 326.2)		Deaths	Stroke	Male	1990	11,229	(9,026 to 14,258)	34.7	(27.9 to 44.3)
2013 16,409 (10,882 to 22,178) 17.5 (12.4 to 23.4)					2013	11,899	(7,687 to 15,528)	20.9	(13.8 to 27.0)
Both 1990 26,096 (22,181 to 31,375) 31.6 (26.9 to 38.0) 2013 28,308 (20,401 to 34,724) 19.1 (14.2 to 23.1) (226,930 to (226,930 to (320,626)) 842.7 (693.0 to 978.0) DALYS Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) 2013 183,571 223,323) 324.6 (258.7 to 394.5) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to (188.6 to 326.2)				Female	1990	14,867	(11,219 to 18,709)	29.1	(22.2 to 36.1)
DALYS Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0)					2013	16,409	(10,882 to 22,178)	17.5	(12.4 to 23.4)
DALYS Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) 2013 183,571 223,323) 324.6 (258.7 to 394.5) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 2013 206,307 290,356) 230.8 (188.6 to 326.2)				Both	1990	26,096	(22,181 to 31,375)	31.6	(26.9 to 38.0)
DALYs Ischemic Stroke Male 1990 275,865 320,626) 842.7 (693.0 to 978.0) (146,510 to (146,510 to (223,323) 324.6 (258.7 to 394.5) (228,800 to (228,800 to (335,568) 584.4 (443.5 to 650.3) (167,359 to (167,359 to (230.8 (188.6 to 326.2)					2013	28,308	(20,401 to 34,724)	19.1	(14.2 to 23.1)
2013 183,571 223,323) 324.6 (258.7 to 394.5) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 2013 206,307 290,356) 230.8 (188.6 to 326.2)									
2013 183,571 223,323) 324.6 (258.7 to 394.5) Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 2013 206,307 290,356) 230.8 (188.6 to 326.2)		DALYs	Ischemic Stroke	Male	1990	275,865		842.7	(693.0 to 978.0)
Female 1990 301,820 (228,800 to 335,568) 584.4 (443.5 to 650.3) (167,359 to 2013 206,307 290,356) 230.8 (188.6 to 326.2)					2042	402 574		2246	(250.71, 204.5)
Female 1990 301,820 335,568) 584.4 (443.5 to 650.3) (167,359 to 2013 206,307 290,356) 230.8 (188.6 to 326.2)					2013	183,5/1	· · · · · · · · · · · · · · · · · · ·	324.6	(258.7 to 394.5)
2013 206,307 (167,359 to 230.8 (188.6 to 326.2)				Female	1990	301 820		58 <i>1 1</i>	(443 5 to 650 3)
2013 206,307 290,356) 230.8 (188.6 to 326.2)				Terriale	1330	301,020	· · · · · · · · · · · · · · · · · · ·	504.4	(++5.5 (0 050.5)
					2013	206,307		230.8	(188.6 to 326.2)
				Both	1990	577,686	(490,458 to	693.1	(587.5 to 765.6)

						638,137)		
						(332,428 to		
				2013	389,879	484,070)	272.8	(232.7 to 337.1)
		Hemorrhagic				(168,517 to		
		Stroke	Male	1990	204,791	247,481)	619.1	(509.7 to 750.1)
						(112,137 to		
				2013	161,074	201,369)	319.8	(224.3 to 398.4)
						(161,917 to		
			Female	1990	205,317	248,340)	463	(368.5 to 561.0)
						(118,520 to		
				2013	159,200	210,745)	232.1	(182.3 to 302.8)
						(360,529 to		
			Both	1990	410,108	475,266)	533.4	(471.7 to 618.0)
						(249,788 to		
				2013	320,274	381,143)	272.2	(218.8 to 321.0)
	YLDs	Ischemic Stroke	Male	1990	25,154	(17,083 to 35,952)	74.5	(50.8 to 106.2)
				2013	31,133	(20,541 to 43,386)	61.1	(40.5 to 84.4)
			Female	1990	18,845	(12,874 to 26,145)	40.3	(27.4 to 55.7)
				2013	22,277	(13,749 to 32,539)	33.9	(21.5 to 48.4)
			Both	1990	43,999	(30,513 to 59,717)	54.8	(38.0 to 75.0)
				2013	53,411	(36,508 to 72,784)	46.2	(31.5 to 62.0)
		Hemorrhagic						
		Stroke	Male	1990	4,641	(3,006 to 6,634)	13.9	(9.0 to 19.8)
				2013	6,703	(4,268 to 9,262)	13.7	(8.8 to 18.8)
			Female	1990	4,734	(3,171 to 6,567)	10.9	(7.3 to 15.2)
				2013	6,052	(3,825 to 8,720)	10.4	(6.7 to 14.7)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	9,375	(6,274 to 12,744)	12.2	(8.1 to 16.5)

			Both	2013	12,755	(8,669 to 17,399)	11.8	(8.1 to 16.1)
Jamaica	Deaths	Ischemic Stroke	Male	1990	617	(518 to 715)	82.6	(69.1 to 95.9)
				2013	724	(542 to 933)	70.9	(53.1 to 91.3)
			Female	1990	998	(825 to 1,207)	100.4	(82.7 to 121.1)
				2013	1,116	(813 to 1,449)	79	(57.9 to 103.1)
			Both	1990	1,616	(1,418 to 1,832)	93.1	(81.7 to 105.6)
				2013	1,840	(1,459 to 2,209)	76.2	(60.4 to 91.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	264	(213 to 332)	34.6	(28.0 to 43.6)
				2013	369	(238 to 492)	34.5	(22.3 to 46.0)
			Female	1990	434	(354 to 571)	45.8	(37.3 to 59.8)
				2013	565	(400 to 768)	41.3	(29.5 to 56.6)
			Both	1990	698	(606 to 888)	40.9	(35.4 to 51.9)
				2013	934	(720 to 1,161)	38.6	(29.7 to 47.9)
	DALYs	Ischemic Stroke	Male	1990	9,413	(7,963 to 10,828)	1221.7	(1032.6 to 1405.8)
				2013	10,870	(7,989 to 14,091)	1013.8	(750.2 to 1314.0)
			Female	1990	12,601	(10,128 to 15,005)	1344.6	(1078.4 to 1602.5)
				2013	12,948	(9,533 to 17,230)	989.6	(722.5 to 1322.2)
			Both	1990	22,015	(19,165 to 24,851)	1294.8	(1127.9 to 1460.2)
				2013	23,819	(19,108 to 28,856)	1008.6	(807.1 to 1222.8)
		Hemorrhagic						
		Stroke	Male	1990	5,954	(4,900 to 7,324)	717	(587.6 to 888.9)
				2013	7,621	(5,292 to 10,286)	656.5	(451.6 to 885.3)
			Female	1990	8,094	(6,531 to 10,057)	882.9	(715.8 to 1098.4)
				2013	9,120	(6,531 to 12,765)	699.6	(498.6 to 983.5)
			Both	1990	14,048	(12,306 to 16,565)	806.7	(706.4 to 957.8)
				2013	16,742	(13,328 to 20,786)	682.5	(540.6 to 844.0)
	YLDs	Ischemic Stroke	Male	1990	359	(239 to 504)	47.6	(31.5 to 66.8)
				2013	494	(314 to 705)	45.7	(29.0 to 65.2)

			Female	1990	281	(177 to 405)	30.8	(19.4 to 43.9)
				2013	394	(253 to 574)	30.5	(19.5 to 44.6)
			Both	1990	640	(433 to 870)	38.4	(26.1 to 52.0)
				2013	888	(601 to 1,209)	37.4	(25.2 to 51.2)
		Hemorrhagic						
		Stroke	Male	1990	119	(74 to 176)	15.4	(9.6 to 22.8)
				2013	166	(100 to 272)	14.5	(8.7 to 24.5)
			Female	1990	92	(55 to 149)	10.2	(6.0 to 16.6)
				2013	132	(81 to 212)	10.1	(6.3 to 16.6)
			Both	1990	211	(136 to 304)	12.6	(8.1 to 18.1)
			Both	2013	298	(192 to 429)	12.2	(7.8 to 17.7)
Japan	Deaths	Ischemic Stroke	Male	1990	36,801	(31,584 to 42,286)	68.9	(58.8 to 78.8)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
·		•		2013	52,867	(42,940 to 63,346)	36.5	(29.7 to 43.7)
			Female	1990	45,489	(34,836 to 51,691)	52.8	(40.3 to 59.9)
				2013	64,703	(49,177 to 99,027)	25.1	(19.2 to 39.0)
			Both	1990	82,291	(70,344 to 90,340)	59.3	(50.6 to 65.1)
						(98,196 to		
				2013	117,570	156,199)	30.1	(25.1 to 39.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	23,262	(19,657 to 28,140)	37.1	(30.9 to 45.4)
				2013	32,670	(22,636 to 40,485)	26.2	(18.7 to 32.2)
			Female	1990	25,550	(20,483 to 32,867)	30	(24.1 to 38.5)
				2013	35,336	(27,376 to 47,837)	17.6	(14.3 to 23.8)
			Both	1990	48,813	(43,561 to 57,923)	33.4	(29.7 to 39.9)
				2013	68,006	(54,476 to 80,497)	21.7	(18.1 to 25.0)
						(434,189 to		,
	1		1		i			
	DALYs	Ischemic Stroke	Male	1990	509,293	581,929)	866.7	(742.2 to 987.3)

Ischemic Stroke	Male Female Both	1990 2013 1990 2013 1990 2013	980,182 967,981 57,454 105,410 45,727 85,368 103,181	1,106,043) (829,670 to 1,103,960) (38,479 to 80,973) (69,263 to 148,249) (30,683 to 64,000) (55,073 to 120,118) (71,665 to 141,523)	654.8 412.3 90.4 85.9 53.8 51.9	(589.9 to 736.8) (356.9 to 468.0) (61.1 to 125.8) (57.2 to 120.0) (36.2 to 75.0) (34.1 to 73.8) (48.1 to 95.4)
Ischemic Stroke	Male	2013 1990 2013 1990	967,981 57,454 105,410 45,727	(829,670 to 1,103,960) (38,479 to 80,973) (69,263 to 148,249) (30,683 to 64,000) (55,073 to 120,118)	412.3 90.4 85.9 53.8	(356.9 to 468.0) (61.1 to 125.8) (57.2 to 120.0) (36.2 to 75.0)
Ischemic Stroke	Male	2013 1990 2013 1990	967,981 57,454 105,410 45,727	(829,670 to 1,103,960) (38,479 to 80,973) (69,263 to 148,249) (30,683 to 64,000) (55,073 to	412.3 90.4 85.9 53.8	(356.9 to 468.0) (61.1 to 125.8) (57.2 to 120.0) (36.2 to 75.0)
Ischemic Stroke	Male	2013 1990 2013	967,981 57,454 105,410	(829,670 to 1,103,960) (38,479 to 80,973) (69,263 to 148,249) (30,683 to 64,000)	412.3 90.4 85.9	(356.9 to 468.0) (61.1 to 125.8) (57.2 to 120.0)
Ischemic Stroke	Male	2013 1990 2013	967,981 57,454 105,410	(829,670 to 1,103,960) (38,479 to 80,973) (69,263 to 148,249)	412.3 90.4 85.9	(356.9 to 468.0) (61.1 to 125.8) (57.2 to 120.0)
Ischemic Stroke		2013 1990	967,981 57,454	(829,670 to 1,103,960) (38,479 to 80,973) (69,263 to	412.3 90.4	(356.9 to 468.0) (61.1 to 125.8)
Ischemic Stroke		2013	967,981	(829,670 to 1,103,960) (38,479 to 80,973)	412.3	(356.9 to 468.0)
Josh ownin Chunkin		2013	967,981	(829,670 to 1,103,960)	412.3	(356.9 to 468.0)
	Both		,	(829,670 to		,
	Both	1990	980,182	· · · ·	654.8	(589.9 to 736.8)
	Doth	1000	000 103	1 106 042)	CE 4 O	(FOO 0 to 726 O)
				(882,687 to		
		2013	402,072		294.4	(249.8 to 391.3)
		2010		• •	•••	(0.40.0)
	Female	1990	438,354		531.4	(435.6 to 637.0)
				• •		
		2013	565,908	678,941)	538.9	(418.8 to 640.6)
				• •		
Stroke	Male	1990	541,828	629,226)	790.2	(684.1 to 919.3)
Hemorrhagic				(471,588 to		
		2013	1,116,524	1,414,590)	339.4	(289.1 to 425.1)
				(956,082 to		
	Both	1990	973,476	1,072,229)	674.5	(582.6 to 742.0)
			,	(841,171 to		,
		2013	495,096	•	238.9	(189.1 to 348.4)
			,			,
	Female	1990	464.182	• •	537.8	(413.3 to 601.8)
			, ,	· ·		(
		2013	621.428	• •	465.6	(390.6 to 543.0)
		Hemorrhagic	2013 Both 1990 2013 Hemorrhagic Stroke Male 1990 2013	Female 1990 464,182 2013 495,096 Both 1990 973,476 2013 1,116,524 Hemorrhagic Stroke Male 1990 541,828 2013 565,908 Female 1990 438,354	Female 1990 464,182 518,703) 2013 495,096 729,961) Both 1990 973,476 1,072,229) Both 1990 973,476 1,072,229) Hemorrhagic 2013 1,116,524 1,414,590) Hemorrhagic Stroke Male 1990 541,828 629,226) 2013 565,908 678,941) Female 1990 438,354 525,949) (357,110 to (357,1	Female 2013 621,428 725,371) 465.6 Female 1990 464,182 518,703) 537.8 392,474 to 2013 495,096 729,961) 238.9 Both 1990 973,476 1,072,229) 674.5 Both 1990 973,476 1,072,229) 674.5 Hemorrhagic (956,082 to 1,414,590) 339.4 Stroke Male 1990 541,828 629,226) 790.2 (430,152 to (430,152 to (430,152 to 678,941) 538.9 Female 1990 438,354 525,949) 531.4 (335,170 to 2013 402,072 536,662) 294.4

Counts, All Ages

Rate Per 100,000 Age- standardized

Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(129,962 to		
				2013	190,778	260,458)	67.5	(46.1 to 92.0)
		Hemorrhagic						
		Stroke	Male	1990	19,181	(12,214 to 28,300)	29.9	(19.3 to 43.8)
				2013	34,374	(21,221 to 50,086)	30.4	(19.1 to 43.8)
			Female	1990	15,919	(9,757 to 23,208)	19.3	(12.1 to 28.1)
				2013	31,797	(20,403 to 46,642)	22.8	(14.7 to 33.1)
			Both	1990	35,100	(22,985 to 49,256)	23.8	(15.7 to 33.3)
			Both	2013	66,172	(43,119 to 91,409)	26.2	(17.3 to 36.0)
Jordan	Deaths	Ischemic Stroke	Male	1990	670	(526 to 876)	121.8	(95.9 to 157.9)
				2013	696	(543 to 878)	70.2	(55.6 to 86.5)
			Female	1990	603	(483 to 758)	117.9	(93.8 to 148.5)
				2013	739	(589 to 898)	74.1	(59.7 to 89.3)
			Both	1990	1,273	(1,074 to 1,522)	119.8	(101.5 to 142.3)
				2013	1,435	(1,212 to 1,650)	72.4	(61.1 to 82.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	260	(176 to 357)	40.9	(27.4 to 57.2)
				2013	277	(198 to 381)	21.5	(15.4 to 29.9)
			Female	1990	226	(156 to 295)	37.3	(25.4 to 50.0)
				2013	259	(189 to 349)	20.5	(14.9 to 27.4)
			Both	1990	486	(374 to 611)	39.2	(30.1 to 49.8)
				2013	536	(435 to 657)	21	(17.2 to 26.1)
	DALYs	Ischemic Stroke	Male	1990	10,598	(8,128 to 14,455)	1700	(1311.7 to 2284.5)
				2013	11,445	(8,686 to 14,873)	861.4	(675.4 to 1090.5)
			Female	1990	9,182	(7,242 to 11,601)	1565.5	(1244.7 to 1980.3)
				2013	10,447	(7,977 to 12,975)	816.9	(643.2 to 996.6)
			Both	1990	19,780	(16,464 to 24,397)	1633	(1363.7 to 1988.2)
				2013	21,893	(18,140 to 25,903)	841	(707.0 to 972.7)

		Hemorrhagic						
		Stroke	Male	1990	6,484	(4,602 to 8,716)	787.7	(537.7 to 1098.8)
				2013	6,500	(4,589 to 8,949)	364	(259.4 to 500.5)
			Female	1990	5,584	(3,795 to 7,098)	718.5	(491.5 to 927.0)
				2013	5,664	(4,108 to 7,694)	324.7	(238.2 to 440.9)
			Both	1990	12,069	(9,400 to 14,797)	754.6	(580.9 to 941.1)
				2013	12,164	(9,744 to 15,042)	344.7	(278.2 to 424.9)
	YLDs	Ischemic Stroke	Male	1990	71	(47 to 103)	9.6	(6.2 to 13.9)
				2013	201	(127 to 288)	10.3	(6.5 to 14.7)
			Female	1990	50	(32 to 75)	7.2	(4.6 to 10.6)
				2013	140	(90 to 203)	7.2	(4.7 to 10.5)
			Both	1990	122	(84 to 167)	8.4	(5.8 to 11.5)
				2013	342	(227 to 484)	8.7	(5.9 to 12.4)
		Hemorrhagic						
		Stroke	Male	1990	10	(5 to 17)	1.3	(0.6 to 2.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	29	(17 to 45)	1.3	(0.8 to 1.9)
			Female	1990	9	(5 to 14)	1.2	(0.6 to 1.9)
				2013	27	(16 to 42)	1.3	(0.8 to 2.0)
			Both	1990	20	(12 to 29)	1.2	(0.7 to 1.8)
			Both	2013	57	(36 to 82)	1.3	(0.8 to 1.8)
Kazakhstan	Deaths	Ischemic Stroke	Male	1990	4,647	(3,732 to 5,633)	152.9	(125.8 to 183.3)
				2013	6,575	(5,234 to 8,082)	168.7	(135.0 to 205.0)
			Female	1990	8,491	(7,303 to 10,251)	128.1	(110.3 to 154.1)
				2013	9,760	(6,686 to 11,793)	138.1	(95.4 to 166.2)
					1			1
			Both	1990	13,138	(11,589 to 15,230)	137.2	(121.4 to 158.7)
			Both	1990 2013	13,138 16,335	(11,589 to 15,230) (12,757 to 18,792)	137.2 150.9	(121.4 to 158.7) (116.9 to 172.9)

		l I		1	I	ı	
D Ib	Hemorrhagic	24.1	4000	2.004	(2.007 4.720)	404.7	/7C F L. 424 A)
Deaths	Stroke	Male	1990	3,891	(3,007 to 4,729)	101.7	(76.5 to 124.4)
			2013	4,897	(3,582 to 6,271)	100.2	(72.4 to 129.3)
		Female	1990	5,692	(4,350 to 6,868)	83.2	(63.5 to 100.5)
			2013	5,852	(4,160 to 7,655)	77.9	(55.4 to 102.6)
		Both	1990	9,583	(7,783 to 10,806)	90.6	(72.9 to 103.0)
			2013	10,749	(8,455 to 12,953)	87.5	(68.1 to 107.4)
					(69,220 to		
DALYs	Ischemic Stroke	Male	1990	89,604	111,785)	2341.6	(1870.4 to 2839.9)
					(104,445 to		
			2013	131,713	165,230)	2676.6	(2141.5 to 3292.8)
					(99,589 to		
		Female	1990	116,569	142,856)	1702	(1458.1 to 2083.4)
					(93,240 to		
			2013	137,432	170,468)	1817.3	(1237.3 to 2243.8)
					(178,871 to		
		Both	1990	206,173	242,098)	1943.5	(1703.7 to 2278.9)
					(213,173 to		
			2013	269,146	315,928)	2171.7	(1717.7 to 2526.0)
	Hemorrhagic				(79,418 to		
	Stroke	Male	1990	100,867	122,680)	2088	(1634.8 to 2542.7)
					(96,746 to		
			2013	130,028	167,109)	2143.6	(1598.2 to 2741.4)
					(87,686 to		
		Female	1990	109,259	130,959)	1539.5	(1233.4 to 1846.0)
					(86,036 to		
			2013	116,724	148,624)	1441.8	(1047.9 to 1835.7)
					(174,244 to		
		Both	1990	210,127	238,003)	1767.9	(1458.2 to 1998.2)
					(201,263 to		
			2013	246,752	296,079)	1748.5	(1403.6 to 2095.7)

	YLDs	Ischemic Stroke	Male	1990	438	(267 to 653)	9.2	(5.7 to 13.5)
				2013	535	(339 to 797)	9.3	(6.1 to 13.6)
			Female	1990	597	(390 to 856)	8.4	(5.5 to 12.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	637	(407 to 933)	7.9	(5.0 to 11.4)
			Both	1990	1,036	(700 to 1,455)	8.8	(6.0 to 12.3)
				2013	1,173	(805 to 1,635)	8.4	(5.8 to 11.6)
		Hemorrhagic						
		Stroke	Male	1990	453	(280 to 668)	8.2	(5.2 to 12.2)
				2013	517	(328 to 767)	8	(5.1 to 11.7)
			Female	1990	338	(208 to 496)	4.6	(2.8 to 6.9)
				2013	377	(230 to 535)	4.5	(2.8 to 6.5)
			Both	1990	791	(525 to 1,122)	6.2	(4.1 to 8.7)
			Both	2013	895	(598 to 1,247)	6	(4.0 to 8.3)
Kenya	Deaths	Ischemic Stroke	Male	1990	1,261	(543 to 1,812)	46.3	(20.4 to 66.6)
				2013	2,347	(1,125 to 3,316)	46.6	(22.8 to 65.2)
			Female	1990	1,727	(877 to 2,468)	59.8	(30.3 to 85.9)
				2013	2,935	(1,443 to 4,208)	51.4	(25.0 to 74.1)
			Both	1990	2,989	(1,630 to 3,984)	53.6	(29.5 to 72.0)
				2013	5,283	(2,993 to 7,250)	49.4	(27.9 to 67.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,634	(1,132 to 2,352)	50	(33.7 to 75.3)
				2013	3,080	(2,188 to 4,613)	49.2	(33.6 to 77.3)
			Female	1990	2,046	(1,467 to 2,781)	60.4	(40.8 to 85.5)
				2013	3,180	(2,057 to 4,512)	46.8	(28.9 to 68.6)
			Both	1990	3,680	(2,788 to 4,717)	55.5	(39.0 to 74.5)
				2013	6,260	(4,759 to 8,474)	48	(34.2 to 67.1)

	DALYs	Ischemic Stroke	Male	1990	21,463	(9,399 to 31,063)	650.1	(283.5 to 933.7)
				2013	39,816	(18,632 to 58,336)	643.8	(307.8 to 918.8)
			Female	1990	25,706	(13,065 to 36,472)	767.4	(393.0 to 1100.7)
				2013	40,745	(20,082 to 60,279)	608.8	(301.0 to 884.5)
			Both	1990	47,170	(25,653 to 63,059)	712.8	(388.9 to 948.1)
						(45,913 to		
				2013	80,562	110,969)	627	(352.2 to 862.1)
		Hemorrhagic						
		Stroke	Male	1990	38,505	(28,262 to 52,245)	910.4	(637.1 to 1285.9)
						(53,318 to		
				2013	73,852	103,005)	879.1	(624.2 to 1301.5)
			Female	1990	43,972	(32,926 to 56,181)	1036.2	(759.5 to 1365.0)
				2013	65,631	(45,099 to 90,318)	737	(488.6 to 1036.6)
						(67,850 to		
			Both	1990	82,478	100,984)	976	(760.0 to 1232.2)
						(110,866 to		(
				2013	139,483	181,354)	805.2	(618.1 to 1081.7)
	YLDs	Ischemic Stroke	Male	1990	466	(291 to 675)	10.1	(6.2 to 14.9)
				2013	1,039	(613 to 1,543)	10.8	(6.3 to 16.1)
			Female	1990	346	(219 to 520)	7.2	(4.6 to 11.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	771	(490 to 1,143)	7.4	(4.7 to 10.8)
			Both	1990	812	(529 to 1,155)	8.6	(5.6 to 12.4)
				2013	1,810	(1,162 to 2,521)	9	(5.8 to 12.5)
		Hemorrhagic						
		Stroke	Male	1990	165	(100 to 248)	2.9	(1.7 to 4.3)
				2013	387	(234 to 609)	3.1	(1.8 to 4.9)
			Female	1990	108	(65 to 166)	1.9	(1.2 to 2.9)

				2013	230	(144 to 347)	1.8	(1.1 to 2.7)
			Both	1990	274	(174 to 401)	2.4	(1.5 to 3.4)
			Both	2013	617	(388 to 888)	2.4	(1.5 to 3.5)
Kiribati	Deaths	Ischemic Stroke	Male	1990	12	(8 to 15)	104.4	(78.8 to 131.5)
				2013	22	(15 to 30)	112.2	(79.3 to 145.7)
			Female	1990	6	(4 to 9)	44.5	(30.5 to 66.2)
				2013	12	(7 to 20)	51.7	(28.9 to 77.8)
			Both	1990	18	(14 to 23)	69.5	(56.2 to 85.2)
				2013	35	(25 to 45)	76.6	(56.8 to 96.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	25	(19 to 33)	162.2	(122.5 to 205.7)
				2013	54	(35 to 76)	196.3	(135.6 to 265.2)
			Female	1990	11	(8 to 18)	65.8	(45.2 to 102.9)
				2013	24	(13 to 36)	86	(49.5 to 124.8)
			Both	1990	37	(29 to 46)	108.7	(87.6 to 136.2)
				2013	78	(55 to 105)	135.6	(96.5 to 175.9)
	DALYs	Ischemic Stroke	Male	1990	314	(228 to 410)	2004.5	(1499.4 to 2558.6)
				2013	571	(373 to 817)	2095.6	(1434.8 to 2835.2)
			Female	1990	150	(98 to 227)	822.5	(550.3 to 1232.7)
				2013	264	(139 to 426)	868.3	(471.6 to 1360.6)
			Both	1990	464	(364 to 584)	1343.4	(1065.2 to 1683.6)
				2013	836	(577 to 1,115)	1412.3	(1013.1 to 1821.1)
		Hemorrhagic						
		Stroke	Male	1990	926	(695 to 1,194)	4304.5	(3215.5 to 5560.9)
				2013	1,830	(1,198 to 2,616)	5092.2	(3368.7 to 7149.8)
			Female	1990	345	(249 to 554)	1552.4	(1082.5 to 2469.4)
				2013	650	(364 to 994)	1835.7	(1033.9 to 2756.6)
			Both	1990	1,272	(1,022 to 1,599)	2836.8	(2251.9 to 3577.8)
				2013	2,481	(1,730 to 3,371)	3375.2	(2393.8 to 4519.3)

2013

160

14

	DALYs	Ischemic Stroke	Male	1990	2,197	(1,644 to 2,852)	708.8	(561.5 to 902.8)
				2013	4,108	(3,031 to 5,069)	615.3	(462.5 to 745.4)
			Female	1990	1,556	(1,211 to 2,228)	684.2	(547.0 to 956.3)
				2013	2,508	(1,622 to 3,002)	626.1	(423.3 to 731.7)
			Both	1990	3,754	(3,083 to 4,768)	695.7	(588.2 to 866.1)
				2013	6,616	(4,946 to 7,829)	613.5	(463.1 to 712.3)
		Hemorrhagic						
		Stroke	Male	1990	2,645	(1,910 to 3,312)	461.9	(304.5 to 597.2)
				2013	3,028	(2,277 to 3,967)	250.2	(190.0 to 350.0)
			Female	1990	2,244	(1,538 to 2,654)	524.5	(345.9 to 640.2)
				2013	1,832	(1,472 to 2,491)	246.1	(196.3 to 359.9)
			Both	1990	4,889	(3,781 to 5,717)	489.2	(351.0 to 588.3)
				2013	4,860	(4,106 to 6,012)	248.1	(208.7 to 328.8)
	YLDs	Ischemic Stroke	Male	1990	50	(31 to 76)	9.2	(5.9 to 13.4)
				2013	112	(72 to 162)	9.9	(6.5 to 13.7)
			Female	1990	25	(15 to 38)	7.4	(4.5 to 10.7)
				2013	51	(32 to 73)	7.6	(4.8 to 10.9)
			Both	1990	76	(49 to 108)	8.4	(5.6 to 11.9)
				2013	164	(106 to 228)	9	(6.0 to 12.2)
		Hemorrhagic						
		Stroke	Male	1990	10	(5 to 16)	1.5	(0.8 to 2.3)
				2013	21	(11 to 32)	1.5	(0.9 to 2.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	6	(3 to 10)	1.6	(0.8 to 2.6)
				2013	10	(5 to 16)	1.2	(0.7 to 1.9)
			Both	1990	17	(11 to 25)	1.5	(1.0 to 2.2)
			Both	2013	31	(20 to 46)	1.4	(0.9 to 2.0)

Kyrgyzstan	Deaths	Ischemic Stroke	Male	1990	1,346	(1,078 to 1,674)	174.2	(142.9 to 212.3)
				2013	1,682	(1,370 to 2,043)	156.7	(127.9 to 189.6)
			Female	1990	2,202	(1,803 to 2,610)	150.5	(123.0 to 177.7)
				2013	1,824	(1,511 to 2,251)	120.8	(100.6 to 149.5)
			Both	1990	3,549	(3,153 to 4,153)	160.3	(142.8 to 186.0)
				2013	3,507	(3,061 to 4,059)	136.6	(118.8 to 159.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	897	(617 to 1,130)	89	(60.0 to 113.4)
				2013	1,293	(1,026 to 1,622)	94.4	(74.3 to 119.0)
			Female	1990	1,195	(825 to 1,433)	76.6	(52.1 to 92.1)
				2013	1,252	(1,011 to 1,621)	72.9	(59.0 to 95.3)
			Both	1990	2,092	(1,568 to 2,433)	81.6	(60.1 to 95.8)
				2013	2,545	(2,167 to 3,023)	82.5	(69.9 to 99.3)
	DALYs	Ischemic Stroke	Male	1990	25,347	(19,449 to 32,911)	2683.9	(2129.1 to 3360.7)
				2013	33,159	(26,746 to 40,512)	2581.4	(2099.4 to 3141.2)
			Female	1990	29,929	(24,728 to 37,496)	1971.3	(1631.8 to 2448.3)
				2013	26,390	(21,509 to 32,195)	1625.6	(1329.6 to 1974.4)
			Both	1990	55,276	(48,086 to 67,303)	2260.6	(1981.1 to 2702.0)
				2013	59,549	(51,113 to 68,788)	2046.5	(1765.1 to 2358.2)
		Hemorrhagic						
		Stroke	Male	1990	23,978	(17,198 to 29,866)	1987.1	(1395.6 to 2492.7)
				2013	36,324	(28,972 to 44,873)	2186.7	(1731.9 to 2739.5)
			Female	1990	26,520	(19,899 to 31,125)	1640.2	(1227.0 to 1923.4)
				2013	30,008	(24,189 to 37,267)	1554.2	(1263.2 to 1968.7)
			Both	1990	50,499	(39,894 to 57,886)	1787.6	(1401.1 to 2056.2)
				2013	66,332	(56,146 to 76,868)	1842.1	(1567.4 to 2162.0)
	YLDs	Ischemic Stroke	Male	1990	99	(62 to 145)	8.6	(5.5 to 12.4)
				2013	136	(86 to 196)	8.9	(5.8 to 12.7)
			Female	1990	133	(80 to 198)	8.4	(5.0 to 12.3)

				2013	152	(98 to 219)	8.1	(5.1 to 11.5)
			Both	1990	233	(150 to 330)	8.5	(5.5 to 12.0)
				2013	289	(193 to 401)	8.4	(5.7 to 11.6)
		Hemorrhagic						
		Stroke	Male	1990	104	(66 to 150)	7.8	(5.1 to 11.5)
				2013	142	(89 to 202)	7.7	(4.9 to 11.0)
			Female	1990	75	(47 to 108)	4.5	(2.8 to 6.5)
				2013	97	(59 to 142)	4.7	(2.8 to 6.9)
			Both	1990	179	(121 to 251)	5.9	(4.0 to 8.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	2013	239	(156 to 335)	6.1	(4.0 to 8.5)
Laos	Deaths	Ischemic Stroke	Male	1990	557	(316 to 1,023)	90.3	(53.1 to 156.2)
				2013	1,130	(766 to 1,699)	104.7	(73.4 to 152.3)
			Female	1990	658	(414 to 1,108)	87.8	(56.9 to 141.6)
				2013	1,443	(793 to 2,101)	109.8	(64.0 to 156.0)
			Both	1990	1,215	(758 to 1,941)	89.3	(58.6 to 138.8)
				2013	2,574	(1,738 to 3,631)	108	(74.7 to 146.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	775	(418 to 1,223)	88.9	(40.6 to 143.4)
				2013	1,153	(654 to 1,865)	76.7	(44.5 to 122.3)
			Female	1990	821	(481 to 1,290)	85.4	(50.5 to 135.3)
				2013	1,290	(744 to 2,019)	77.9	(44.4 to 121.1)
			Both	1990	1,597	(992 to 2,420)	87.4	(54.2 to 130.1)
				2013	2,443	(1,446 to 3,789)	77.7	(48.8 to 117.5)
	DALYs	Ischemic Stroke	Male	1990	12,836	(7,512 to 23,556)	1492.4	(859.3 to 2746.4)
				2013	22,154	(14,331 to 35,139)	1603.6	(1076.9 to 2449.5)
			Female	1990	12,870	(7,381 to 22,065)	1332.7	(821.4 to 2279.9)

				2013	23,860	(11,917 to 37,093)	1525.1	(811.4 to 2261.8)
			Both	1990	25,707	(15,454 to 41,328)	1410.5	(876.6 to 2259.9)
				2013	46,015	(29,662 to 69,108)	1566.8	(1051.4 to 2269.8)
		Hemorrhagic						
		Stroke	Male	1990	26,948	(15,355 to 41,025)	2164.3	(1158.0 to 3413.5)
				2013	36,875	(21,137 to 59,752)	1829.8	(1042.3 to 2978.6)
			Female	1990	24,521	(13,988 to 38,043)	1914.4	(1115.7 to 3013.2)
				2013	35,142	(18,730 to 56,379)	1686	(933.5 to 2671.1)
			Both	1990	51,469	(31,624 to 77,259)	2038.5	(1236.1 to 3101.1)
						(42,385 to		
				2013	72,018	111,807)	1759.5	(1039.4 to 2753.1)
	YLDs	Ischemic Stroke	Male	1990	214	(138 to 301)	27.9	(18.1 to 39.0)
				2013	375	(238 to 543)	27	(16.4 to 39.4)
			Female	1990	200	(128 to 280)	21.2	(14.0 to 29.5)
				2013	385	(253 to 548)	23.8	(15.6 to 34.2)
			Both	1990	414	(276 to 573)	24.1	(16.2 to 32.7)
				2013	760	(523 to 1,063)	25.3	(17.2 to 35.5)
		Hemorrhagic						
		Stroke	Male	1990	265	(175 to 376)	30.6	(20.4 to 43.4)
				2013	456	(293 to 647)	28.8	(18.6 to 41.1)
			Female	1990	215	(138 to 301)	21	(13.8 to 29.2)
				2013	401	(257 to 564)	22.7	(14.3 to 32.0)
			Both	1990	480	(324 to 657)	25.3	(17.3 to 34.1)
			Both	2013	857	(572 to 1,171)	25.5	(17.0 to 35.0)
Latvia	Deaths	Ischemic Stroke	Male	1990	1,538	(1,268 to 1,859)	158.2	(131.1 to 190.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,419	(1,174 to 1,650)	118.8	(98.5 to 138.3)

		Female	1990	3,088	(2,621 to 3,716)	134.1	(113.9 to 162.0)
			2013	2,532	(2,113 to 2,995)	83.7	(69.9 to 98.3)
		Both	1990	4,627	(4,101 to 5,356)	142.9	(126.8 to 165.7)
			2013	3,952	(3,429 to 4,406)	96.4	(83.5 to 107.3)
	Hemorrhagic						
Deaths	Stroke	Male	1990	803	(540 to 1,097)	74.4	(49.1 to 103.6)
			2013	486	(370 to 667)	40.4	(30.8 to 55.6)
		Female	1990	1,405	(880 to 1,738)	64.1	(40.8 to 79.0)
			2013	635	(465 to 1,121)	24.5	(18.7 to 40.3)
		Both	1990	2,208	(1,531 to 2,733)	68.5	(47.6 to 85.0)
			2013	1,122	(884 to 1,646)	30.6	(24.6 to 43.4)
DALYs	Ischemic Stroke	Male	1990	23,850	(19,195 to 29,410)	2253	(1844.3 to 2747.8)
			2013	20,913	(17,055 to 24,194)	1696.5	(1380.4 to 1963.0)
		Female	1990	35,993	(30,437 to 43,478)	1632.2	(1383.3 to 1981.0)
			2013	25,057	(20,852 to 29,501)	938.2	(781.8 to 1121.6)
		Both	1990	59,844	(52,314 to 69,297)	1858.9	(1626.3 to 2151.0)
			2013	45,970	(40,012 to 51,205)	1223.1	(1061.0 to 1362.4)
	Hemorrhagic						
	Stroke	Male	1990	17,261	(12,017 to 22,619)	1444.4	(997.7 to 1927.0)
			2013	9,771	(7,621 to 12,954)	804.1	(630.1 to 1061.0)
		Female	1990	22,025	(14,845 to 26,224)	1084	(743.4 to 1293.6)
			2013	8,652	(6,783 to 13,532)	414.5	(332.6 to 623.6)
		Both	1990	39,287	(29,092 to 48,189)	1233.4	(916.5 to 1506.7)
			2013	18,424	(15,325 to 24,781)	575.3	(483.1 to 749.0)
YLDs	Ischemic Stroke	Male	1990	437	(273 to 628)	35.6	(22.5 to 50.4)
			2013	525	(328 to 760)	43.9	(27.9 to 63.4)
		Female	1990	481	(310 to 668)	25	(16.2 to 34.6)
			2013	461	(293 to 669)	24.1	(15.2 to 34.7)
		Both	1990	918	(604 to 1,246)	29.4	(19.4 to 40.0)

				2013	987	(650 to 1,356)	32.3	(21.0 to 44.4)
		Hemorrhagic						
		Stroke	Male	1990	28	(14 to 48)	2.3	(1.2 to 3.8)
				2013	29	(13 to 53)	2.5	(1.2 to 4.6)
			Female	1990	28	(16 to 46)	1.6	(0.9 to 2.6)
				2013	28	(15 to 50)	1.7	(0.9 to 3.0)
			Both	1990	56	(33 to 87)	1.9	(1.1 to 2.9)
			Both	2013	57	(33 to 90)	2.1	(1.2 to 3.2)
Lebanon	Deaths	Ischemic Stroke	Male	1990	234	(155 to 336)	37.4	(25.0 to 52.5)
				2013	199	(49 to 334)	11.7	(2.9 to 19.6)
			Female	1990	559	(323 to 782)	73.8	(43.4 to 100.9)
				2013	975	(260 to 1,429)	49.1	(13.0 to 72.3)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Country	IIICasarc	Stroke Type	Both	1990	793	(532 to 1,035)	56.8	(38.6 to 73.6)
			Dotti	2013	1,174	(351 to 1,639)	31.6	(9.4 to 44.2)
		Hemorrhagic		2013	1,174	(331 to 1,033)	31.0	(3.4 to 44.2)
	Deaths	Stroke	Male	1990	216	(136 to 317)	29	(17.2 to 43.4)
				2013	160	(70 to 353)	8.8	(3.6 to 19.8)
			Female	1990	532	(249 to 778)	62.5	(28.9 to 93.1)
				2013	736	(130 to 1,306)	36.5	(6.4 to 65.1)
			Both	1990	748	(434 to 1,007)	46.6	(26.4 to 63.5)
				2013	897	(257 to 1,500)	23.2	(6.5 to 39.0)
	DALYs	Ischemic Stroke	Male	1990	3,766	(2,439 to 5,524)	482.1	(318.5 to 693.1)
								, ,
				2013	2,641	(702 to 4,714)	134.4	(36.5 to 236.1)
			Female	2013 1990	2,641 8,690	(702 to 4,714) (4,630 to 12,509)	134.4 1000.8	(36.5 to 236.1) (553.1 to 1425.0)
			Female		-			

	_	1	, , , , , , , , , , , , , , , , , , , 		П	T		
				2013	14,235	(4,211 to 20,096)	352.3	(104.8 to 493.3)
		Hemorrhagic						
		Stroke	Male	1990	5,553	(3,885 to 8,191)	552.8	(368.8 to 818.3)
				2013	2,964	(1,522 to 5,726)	136.3	(69.0 to 268.3)
			Female	1990	12,389	(5,769 to 17,436)	1200.6	(555.4 to 1718.1)
				2013	12,088	(2,442 to 20,200)	571.9	(115.1 to 964.1)
			Both	1990	17,943	(11,094 to 23,550)	884.1	(536.0 to 1171.9)
				2013	15,053	(4,796 to 23,730)	351.8	(110.4 to 559.4)
	YLDs	Ischemic Stroke	Male	1990	81	(53 to 115)	8.7	(5.7 to 12.3)
				2013	213	(136 to 309)	9.3	(5.9 to 13.4)
			Female	1990	65	(42 to 93)	6.7	(4.5 to 9.6)
				2013	157	(99 to 227)	7.4	(4.6 to 10.7)
			Both	1990	147	(99 to 198)	7.7	(5.3 to 10.4)
				2013	371	(248 to 512)	8.4	(5.6 to 11.5)
		Hemorrhagic						
		Stroke	Male	1990	12	(7 to 20)	1.3	(0.7 to 2.0)
				2013	32	(19 to 50)	1.3	(0.8 to 2.1)
			Female	1990	13	(7 to 20)	1.3	(0.7 to 2.0)
				2013	29	(16 to 47)	1.3	(0.8 to 2.2)
			Both	1990	25	(16 to 38)	1.3	(0.8 to 1.9)
			Both	2013	61	(38 to 93)	1.3	(0.8 to 2.0)
Lesotho	Deaths	Ischemic Stroke	Male	1990	154	(78 to 219)	59	(30.3 to 83.7)
				2013	283	(173 to 389)	83.8	(52.3 to 114.4)
			Female	1990	169	(89 to 243)	49.7	(26.6 to 70.6)
				2013	305	(168 to 489)	58.9	(32.1 to 93.8)
			Both	1990	323	(215 to 437)	53.7	(35.6 to 72.0)
				2013	589	(414 to 778)	68.7	(48.4 to 90.8)
								•
					Counts, All	Δσες	Rate	Per 100.000

Counts, All Ages

Rate Per 100,000

							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
·		Hemorrhagic						
	Deaths	Stroke	Male	1990	237	(164 to 326)	80	(54.4 to 112.3)
				2013	405	(295 to 556)	101.7	(73.2 to 142.4)
			Female	1990	239	(158 to 315)	63.2	(41.4 to 84.7)
				2013	345	(231 to 620)	58.9	(38.6 to 105.2)
			Both	1990	476	(350 to 586)	70.4	(50.6 to 88.1)
				2013	750	(570 to 1,063)	75.8	(56.8 to 108.9)
	DALYs	Ischemic Stroke	Male	1990	3,045	(1,501 to 4,409)	971.1	(490.0 to 1392.7)
				2013	5,583	(3,299 to 7,914)	1447	(873.4 to 2012.7)
			Female	1990	2,874	(1,531 to 4,322)	711.7	(381.6 to 1044.7)
				2013	4,832	(2,680 to 8,092)	837.3	(463.0 to 1376.4)
			Both	1990	5,919	(3,849 to 8,140)	823.4	(540.9 to 1115.2)
				2013	10,415	(7,323 to 13,711)	1078.7	(758.6 to 1425.9)
		Hemorrhagic						
		Stroke	Male	1990	5,827	(4,109 to 7,830)	1583.7	(1103.1 to 2152.1)
				2013	11,475	(8,477 to 15,274)	2293.2	(1657.2 to 3130.6)
			Female	1990	5,170	(3,444 to 6,781)	1121.3	(737.5 to 1484.7)
				2013	8,910	(5,872 to 15,301)	1250.9	(837.5 to 2192.0)
			Both	1990	10,997	(8,345 to 13,475)	1322.7	(981.9 to 1623.7)
				2013	20,385	(15,627 to 28,552)	1671.6	(1281.9 to 2348.7)
	YLDs	Ischemic Stroke	Male	1990	69	(46 to 98)	19.5	(13.1 to 27.7)
				2013	84	(53 to 117)	19.2	(12.2 to 26.6)
			Female	1990	88	(56 to 127)	19.1	(12.1 to 27.1)
				2013	114	(73 to 164)	18.7	(11.9 to 27.1)
			Both	1990	158	(107 to 218)	19.3	(13.0 to 26.6)
				2013	198	(130 to 272)	18.9	(12.5 to 26.1)
		Hemorrhagic	Male	1990	20	(12 to 30)	5.2	(3.2 to 7.8)

		Stroke						
				2013	27	(17 to 39)	5.3	(3.3 to 7.6)
			Female	1990	23	(14 to 35)	4.7	(3.0 to 7.0)
				2013	29	(17 to 42)	4.4	(2.6 to 6.5)
			Both	1990	44	(28 to 62)	4.9	(3.2 to 6.9)
			Both	2013	56	(36 to 78)	4.7	(3.0 to 6.7)
Liberia	Deaths	Ischemic Stroke	Male	1990	193	(136 to 258)	71.4	(49.5 to 94.2)
				2013	673	(370 to 914)	122.3	(71.0 to 159.7)
			Female	1990	245	(187 to 370)	85.4	(65.5 to 127.9)
				2013	791	(524 to 1,115)	128.3	(87.9 to 179.8)
			Both	1990	439	(361 to 564)	78.8	(65.1 to 101.1)
				2013	1,465	(1,045 to 1,796)	125.7	(92.2 to 153.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	229	(147 to 302)	55.9	(32.7 to 76.0)
				2013	475	(290 to 758)	63.4	(36.4 to 109.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	251	(184 to 351)	62.9	(44.5 to 90.0)
				2013	481	(299 to 929)	60.1	(36.0 to 124.4)
			Both	1990	480	(362 to 581)	59.7	(42.6 to 72.8)
				2013	957	(679 to 1,580)	61.7	(41.3 to 109.3)
	DALYs	Ischemic Stroke	Male	1990	4,704	(3,283 to 6,433)	1129.9	(795.2 to 1513.0)
				2013	13,961	(7,421 to 19,685)	1894.8	(1044.7 to 2569.0)
			Female	1990	5,523	(3,958 to 8,933)	1299.9	(994.4 to 1949.6)
				2013	14,955	(9,484 to 21,912)	1897.2	(1258.1 to 2682.5)
			Both	1990	10,228	(8,162 to 13,919)	1219.8	(1002.5 to 1565.0)
				2013	28,916	(20,615 to 36,178)	1899.4	(1351.7 to 2333.2)
		Hemorrhagic	Male	1990	8,967	(6,005 to 12,249)	1247	(779.0 to 1645.0)

		Stroke						
		Stroke		2013	14,545	(9,442 to 21,183)	1337	(820.7 to 2128.8)
			Female	1990	8,405	(5,817 to 12,261)	1301.1	(957.5 to 1814.8)
			Terriale	2013	13,113	(8,474 to 24,450)	1218	(765.8 to 2338.9)
			Both	1990	-	(12,559 to 22,074)	1277.3	(959.8 to 1540.2)
			BULII	2013	17,372	· · · · · · · · · · · · · · · · · · ·		•
	VI D -	lashani'a Chualia	N 4 = 1 =		27,659 72	(20,629 to 42,420)	1276.7	(909.7 to 2090.4)
	YLDs	Ischemic Stroke	Male	1990		(46 to 107)	15.8	(10.4 to 23.1)
				2013	144	(93 to 205)	15.2	(9.8 to 21.7)
			Female	1990	57	(36 to 84)	11.5	(7.5 to 16.7)
				2013	122	(78 to 174)	11.7	(7.6 to 16.7)
			Both	1990	130	(87 to 186)	13.6	(9.2 to 19.2)
				2013	266	(181 to 366)	13.3	(9.1 to 18.4)
		Hemorrhagic						
		Stroke	Male	1990	22	(14 to 33)	4.2	(2.6 to 6.1)
				2013	45	(28 to 65)	3.9	(2.5 to 5.7)
			Female	1990	16	(10 to 24)	2.9	(1.8 to 4.2)
				2013	34	(21 to 50)	2.8	(1.8 to 4.2)
			Both	1990	39	(25 to 57)	3.5	(2.4 to 5.0)
			Both	2013	79	(51 to 112)	3.3	(2.2 to 4.7)
Libya	Deaths	Ischemic Stroke	Male	1990	362	(256 to 491)	66.6	(48.8 to 88.3)
				2013	862	(616 to 1,185)	69.7	(49.7 to 94.1)
			Female	1990	455	(337 to 588)	79.6	(59.2 to 101.5)
				2013	1,149	(773 to 1,518)	76.9	(52.0 to 101.3)
			Both	1990	817	(649 to 1,000)	73.4	(59.4 to 88.9)
				2013	2,012	(1,558 to 2,464)	73.8	(57.2 to 90.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	464	(342 to 634)	49.7	(34.9 to 68.8)
				2013	705	(474 to 976)	46.3	(29.9 to 64.8)
			Female	1990	571	(411 to 731)	66.3	(47.2 to 86.6)

				2013	960	(662 to 1,316)	56.4	(38.2 to 77.7)
			Both	1990	1,035	(819 to 1,271)	58.1	(45.5 to 73.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,665	(1,272 to 2,166)	51.9	(38.9 to 68.3)
	DALYs	Ischemic Stroke	Male	1990	7,577	(5,093 to 11,063)	876.6	(622.2 to 1193.4)
				2013	13,519	(9,558 to 19,051)	887.4	(632.8 to 1231.3)
			Female	1990	8,775	(5,935 to 12,565)	1046.3	(754.9 to 1364.9)
				2013	15,792	(10,601 to 21,029)	928	(619.0 to 1237.2)
			Both	1990	16,352	(12,314 to 21,221)	961.2	(760.5 to 1180.8)
				2013	29,311	(22,395 to 36,395)	910.3	(702.2 to 1125.1)
		Hemorrhagic						
		Stroke	Male	1990	20,875	(14,799 to 29,401)	1141.9	(841.0 to 1561.5)
				2013	18,698	(13,648 to 25,446)	874	(612.9 to 1198.0)
			Female	1990	22,937	(15,133 to 31,302)	1461.5	(1056.9 to 1885.6)
				2013	21,972	(15,322 to 29,915)	1008.6	(705.2 to 1388.0)
			Both	1990	43,813	(33,945 to 55,185)	1294.4	(1027.0 to 1588.0)
				2013	40,671	(32,273 to 51,090)	945.2	(737.4 to 1204.1)
	YLDs	Ischemic Stroke	Male	1990	26	(16 to 41)	2.5	(1.5 to 3.9)
				2013	43	(27 to 64)	2.2	(1.4 to 3.2)
			Female	1990	18	(10 to 28)	1.9	(1.2 to 3.0)
				2013	38	(23 to 59)	1.9	(1.1 to 2.8)
			Both	1990	44	(28 to 66)	2.2	(1.4 to 3.3)
				2013	82	(52 to 118)	2	(1.3 to 3.0)
		Hemorrhagic						
		Stroke	Male	1990	4	(2 to 6)	0.3	(0.2 to 0.5)
				2013	6	(3 to 10)	0.3	(0.2 to 0.5)
			Female	1990	4	(2 to 7)	0.4	(0.2 to 0.7)

				2013	6	(3 to 9)	0.3	(0.2 to 0.4)
			Both	1990	8	(4 to 13)	0.4	(0.2 to 0.6)
			Both	2013	12	(7 to 18)	0.3	(0.2 to 0.4)
Lithuania	Deaths	Ischemic Stroke	Male	1990	1,307	(1,065 to 1,626)	90.3	(73.8 to 112.2)
				2013	1,362	(1,092 to 1,623)	73.8	(59.2 to 87.9)
			Female	1990	2,240	(1,961 to 2,932)	81.6	(71.6 to 106.3)
				2013	2,774	(1,988 to 3,208)	64.4	(47.1 to 73.8)
			Both	1990	3,547	(3,145 to 4,339)	85.2	(75.8 to 103.9)
				2013	4,137	(3,274 to 4,667)	69	(55.0 to 77.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	507	(344 to 685)	31.9	(21.5 to 43.3)
				2013	415	(298 to 546)	23.1	(16.6 to 30.2)
			Female	1990	736	(488 to 903)	28.4	(19.3 to 34.6)
				2013	642	(483 to 911)	18.2	(14.4 to 25.2)
			Both	1990	1,244	(909 to 1,486)	30.1	(22.1 to 36.0)
				2013	1,057	(846 to 1,379)	20.5	(16.8 to 26.1)
	DALYs	Ischemic Stroke	Male	1990	20,653	(16,759 to 25,583)	1362	(1119.3 to 1676.2)
				2013	20,456	(16,498 to 24,307)	1112.5	(895.0 to 1320.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	27,570	(24,371 to 35,450)	1049.4	(924.0 to 1349.9)
				2013	27,707	(20,947 to 31,442)	737.7	(564.1 to 839.6)
			Both	1990	48,224	(42,800 to 58,354)	1176.5	(1045.2 to 1421.0)
				2013	48,163	(39,200 to 53,850)	894.7	(735.7 to 1003.6)
		Hemorrhagic						
		Stroke	Male	1990	12,406	(8,807 to 16,367)	732	(519.5 to 970.5)
				2013	9,321	(6,794 to 11,961)	530.1	(385.7 to 677.2)
			Female	1990	13,966	(9,558 to 17,053)	574.5	(397.6 to 698.1)

				2013	9,625	(7,764 to 13,114)	344.6	(281.7 to 452.2)
			Doth		-			,
			Both	1990	26,372	(20,061 to 31,331)	646.5	(494.8 to 769.0)
			_	2013	18,947	(15,669 to 23,266)	428.7	(355.9 to 520.4)
	YLDs	Ischemic Stroke	Male	1990	502	(319 to 724)	29.9	(19.1 to 43.1)
				2013	603	(380 to 870)	34.3	(21.7 to 49.4)
			Female	1990	491	(322 to 699)	20.3	(13.4 to 29.0)
				2013	558	(366 to 810)	20.2	(13.2 to 29.4)
			Both	1990	993	(675 to 1,375)	24.3	(16.6 to 33.8)
				2013	1,161	(801 to 1,610)	26.2	(18.0 to 36.0)
		Hemorrhagic						
		Stroke	Male	1990	36	(19 to 62)	2.1	(1.1 to 3.6)
				2013	32	(17 to 60)	2	(1.1 to 3.5)
			Female	1990	30	(18 to 49)	1.3	(0.8 to 2.1)
				2013	37	(20 to 66)	1.5	(0.8 to 2.8)
			Both	1990	66	(40 to 102)	1.7	(1.0 to 2.5)
			Both	2013	69	(42 to 111)	1.7	(1.1 to 2.8)
Luxembourg	Deaths	Ischemic Stroke	Male	1990	133	(102 to 161)	74.7	(57.0 to 89.9)
				2013	97	(73 to 127)	28	(21.0 to 36.6)
			Female	1990	232	(153 to 268)	68.3	(45.1 to 78.7)
				2013	149	(105 to 243)	26	(18.5 to 42.8)
			Both	1990	366	(275 to 415)	71.1	(53.4 to 80.6)
				2013	246	(193 to 358)	27	(21.3 to 39.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	85	(68 to 115)	45.6	(35.7 to 62.5)
				2013	90	(61 to 114)	26.2	(17.9 to 33.2)
			Female	1990	135	(99 to 170)	42.1	(31.4 to 52.7)
				2013	111	(79 to 161)	21	(15.2 to 31.1)
			Both	1990	221	(187 to 273)	43.8	(37.2 to 54.0)
				2013	201	(152 to 249)	23.4	(17.9 to 28.9)

	DALYs	Ischemic Stroke	Male	1990	1,932	(1,503 to 2,306)	1007.9	(784.9 to 1202.2)
				2013	1,353	(1,064 to 1,716)	397	(312.4 to 503.2)
			Female	1990	2,572	(1,767 to 2,937)	805.7	(553.6 to 919.1)
				2013	1,461	(1,097 to 2,247)	302.5	(229.2 to 471.6)
			Both	1990	4,504	(3,519 to 5,055)	888.1	(693.5 to 999.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	2,815	(2,272 to 3,848)	346.3	(280.8 to 473.2)
		Hemorrhagic						
		Stroke	Male	1990	1,587	(1,277 to 2,033)	787.1	(639.5 to 1013.9)
				2013	1,340	(959 to 1,672)	396.1	(283.3 to 494.6)
			Female	1990	1,883	(1,420 to 2,284)	659.5	(494.0 to 793.7)
				2013	1,208	(877 to 1,798)	280.8	(200.6 to 425.1)
			Both	1990	3,470	(3,013 to 4,108)	716.4	(620.3 to 845.7)
				2013	2,548	(2,030 to 3,059)	335.3	(265.6 to 402.1)
	YLDs	Ischemic Stroke	Male	1990	174	(117 to 244)	86.7	(58.1 to 121.3)
				2013	309	(207 to 429)	91.5	(61.3 to 126.4)
			Female	1990	175	(111 to 248)	59.9	(38.9 to 85.6)
				2013	252	(170 to 348)	61.1	(41.0 to 85.0)
			Both	1990	349	(238 to 478)	70.8	(48.5 to 96.9)
				2013	562	(385 to 761)	75.1	(51.1 to 101.3)
		Hemorrhagic						
		Stroke	Male	1990	33	(18 to 56)	16.5	(8.8 to 27.9)
				2013	57	(31 to 96)	17.1	(9.4 to 29.0)
			Female	1990	33	(18 to 55)	12.4	(6.8 to 19.8)
				2013	50	(27 to 83)	12.9	(7.3 to 21.8)
			Both	1990	67	(41 to 105)	14.1	(8.6 to 21.7)
			Both	2013	107	(67 to 165)	14.8	(9.3 to 22.6)

Macedonia	Deaths	Ischemic Stroke	Male	1990	1,364	(1,209 to 1,508)	228.9	(203.6 to 252.2)
				2013	1,952	(1,657 to 2,226)	193.3	(164.9 to 221.0)
			Female	1990	1,231	(1,089 to 1,395)	183.8	(163.0 to 208.8)
				2013	2,165	(1,737 to 2,453)	158.1	(126.3 to 178.6)
			Both	1990	2,596	(2,394 to 2,786)	204.8	(189.3 to 219.7)
				2013	4,117	(3,593 to 4,487)	173.6	(151.6 to 189.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	464	(386 to 561)	60.1	(49.2 to 73.8)
				2013	548	(425 to 693)	47.5	(36.7 to 61.1)
			Female	1990	527	(450 to 648)	62.2	(52.3 to 77.4)
				2013	674	(520 to 839)	48.5	(37.5 to 60.7)
			Both	1990	991	(869 to 1,146)	61.3	(53.2 to 71.8)
				2013	1,223	(1,027 to 1,454)	48.2	(40.5 to 58.0)
	DALYs	Ischemic Stroke	Male	1990	19,320	(16,803 to 21,661)	2836.6	(2500.1 to 3152.7)
				2013	25,812	(21,623 to 29,496)	2321.3	(1958.2 to 2644.3)
			Female	1990	15,581	(13,690 to 18,038)	2102.3	(1870.1 to 2404.7)
				2013	23,636	(19,009 to 26,912)	1676.8	(1340.6 to 1907.1)
			Both	1990	34,902	(31,875 to 38,007)	2447.5	(2256.2 to 2641.3)
				2013	49,449	(43,109 to 54,225)	1965.4	(1719.9 to 2153.6)
		Hemorrhagic						
		Stroke	Male	1990	11,784	(9,929 to 14,007)	1321.4	(1114.5 to 1576.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	11,866	(9,383 to 14,416)	959.2	(756.6 to 1164.4)
			Female	1990	12,420	(10,860 to 15,208)	1310.4	(1139.1 to 1600.5)
				2013	12,627	(10,024 to 15,291)	915.2	(724.4 to 1105.5)
			Both	1990	24,204	(21,561 to 27,580)	1317.7	(1172.3 to 1512.0)
				2013	24,494	(20,896 to 28,217)	937.6	(796.5 to 1078.2)

	YLDs	Ischemic Stroke	Male	1990	200	(125 to 288)	23.2	(14.6 to 33.3)
				2013	327	(211 to 484)	26.2	(16.9 to 38.7)
			Female	1990	140	(89 to 199)	15.5	(9.8 to 22.1)
	Hemorrhagic Stroke			2013	200	(125 to 293)	14.6	(9.2 to 21.3)
			Both	1990	341	(225 to 469)	19.2	(12.7 to 26.1)
			2013	528	(347 to 737)	20.1	(13.3 to 28.0)	
		Male	1990	44	(27 to 66)	4.8	(3.0 to 7.3)	
				2013	64	(40 to 96)	5.2	(3.3 to 7.7)
			Female	1990	32	(19 to 50)	3.4	(2.0 to 5.1)
				2013	46	(26 to 70)	3.5	(2.1 to 5.3)
			Both	1990	77	(48 to 112)	4.1	(2.6 to 5.9)
			Both	2013	110	(72 to 156)	4.3	(2.9 to 6.1)
Madagascar	Deaths	Ischemic Stroke	Male	1990	3,089	(1,662 to 4,179)	174.3	(94.5 to 233.6)
				2013	5,162	(2,451 to 7,749)	166.1	(80.5 to 243.1)
			Female	1990	2,637	(1,843 to 3,448)	168.4	(116.0 to 217.1)
				2013	5,486	(3,341 to 7,575)	168.8	(104.7 to 228.0)
			Both	1990	5,726	(3,913 to 7,080)	171.8	(117.9 to 213.4)
				2013	10,649	(6,421 to 14,005)	167.7	(103.3 to 214.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,604	(3,474 to 6,030)	199.7	(142.1 to 278.0)
				2013	8,187	(5,653 to 11,862)	192.7	(133.3 to 294.1)
			Female	1990	3,858	(3,040 to 4,800)	188.3	(143.9 to 243.9)
				2013	7,832	(5,265 to 11,071)	187.1	(125.5 to 266.4)
			Both	1990	8,463	(6,898 to 10,224)	194	(152.3 to 243.6)
				2013	16,019	(12,157 to 21,196)	190.2	(144.3 to 258.8)
	DALYs	Ischemic Stroke	Male	1990	64,166	(34,815 to 86,318)	2854.8	(1538.2 to 3860.2)
				2013	105,517	(47,536 to 162,224)	2631.8	(1252.5 to 3987.1)

			Female	1990	47,191	(33,221 to 63,667)	2435.6	(1717.8 to 3205.9)
						(54,915 to		
				2013	94,555	135,395)	2372.8	(1421.9 to 3304.0)
						(74,522 to		
			Both	1990	111,358	138,943)	2649.7	(1787.7 to 3278.6)
						(114,705 to		
				2013	200,073	268,535)	2500.4	(1484.8 to 3291.1)
		Hemorrhagic				(110,052 to		
		Stroke	Male	1990	139,107	170,878)	4699.6	(3627.3 to 6065.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(168,983 to		
				2013	247,677	347,353)	4376.2	(3012.7 to 6352.7)
						(81,807 to		
			Female	1990	104,723	126,537)	3950.1	(3102.6 to 4887.2)
						(131,002 to		
				2013	198,704	279,348)	3740.6	(2492.8 to 5242.1)
						(202,658 to		
			Both	1990	243,831	286,051)	4320	(3539.1 to 5177.9)
						(334,436 to		
				2013	446,381	584,877)	4059	(3068.2 to 5385.8)
	YLDs	Ischemic Stroke	Male	1990	294	(184 to 431)	10.8	(6.8 to 16.1)
				2013	599	(382 to 865)	11.6	(7.3 to 16.7)
			Female	1990	189	(114 to 284)	7.2	(4.4 to 10.7)
				2013	436	(280 to 636)	7.7	(4.8 to 11.4)
			Both	1990	484	(314 to 682)	9.1	(5.9 to 12.9)
				2013	1,035	(690 to 1,418)	9.6	(6.4 to 13.4)
		Hemorrhagic		2013	1,000	(050 to 1,710)	5.0	(0.7 (0 13.7)
		Stroke	Male	1990	89	(55 to 132)	2.8	(1.7 to 4.2)
		23.2		2013	188	(117 to 274)	3	(1.8 to 4.3)
				2013	100	(11/102/7)	,	(1.0 (0 7.5)

			Female	1990	53	(33 to 82)	1.7	(1.1 to 2.7)
				2013	116	(70 to 174)	1.7	(1.0 to 2.6)
			Both	1990	143	(92 to 206)	2.3	(1.4 to 3.3)
			Both	2013	304	(199 to 429)	2.3	(1.5 to 3.3)
Malawi	Deaths	Ischemic Stroke	Male	1990	320	(69 to 498)	33.5	(6.8 to 52.2)
				2013	974	(242 to 1,513)	49.2	(12.4 to 74.5)
			Female	1990	606	(238 to 927)	53.3	(20.0 to 80.8)
				2013	1,271	(492 to 1,969)	55.9	(21.9 to 85.1)
			Both	1990	927	(390 to 1,339)	44.5	(18.4 to 64.9)
				2013	2,245	(914 to 3,273)	53	(22.0 to 76.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	592	(416 to 834)	42.3	(27.2 to 65.5)
				2013	1,433	(948 to 2,267)	58.3	(36.5 to 95.8)
			Female	1990	1,064	(774 to 1,484)	69.9	(47.1 to 104.3)
				2013	1,821	(1,275 to 2,812)	66.3	(44.7 to 107.5)
			Both	1990	1,657	(1,265 to 2,191)	57.5	(41.1 to 81.8)
				2013	3,255	(2,402 to 4,703)	62.9	(44.4 to 94.9)
	DALYs	Ischemic Stroke	Male	1990	6,768	(2,069 to 10,279)	473.5	(106.0 to 732.1)
				2013	17,339	(4,540 to 26,958)	696.8	(176.5 to 1074.6)
			Female	1990	10,997	(4,490 to 16,716)	728	(288.6 to 1101.4)
				2013	19,598	(7,361 to 30,674)	710.1	(275.6 to 1099.2)
			Both	1990	17,766	(7,661 to 25,038)	614.5	(256.2 to 885.6)
				2013	36,938	(15,001 to 54,440)	705.6	(292.5 to 1024.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	18,674	(13,926 to 24,976)	857.2	(602.6 to 1205.3)

				2013	35,517	(24,992 to 53,306)	1084.6	(726.2 to 1702.1)
			Female	1990	27,871	(20,948 to 38,616)	1314.9	(960.9 to 1826.2)
				2013	38,876	(27,857 to 56,055)	1109.7	(784.8 to 1684.6)
			Both	1990	46,545	(37,235 to 58,915)	1107.2	(850.4 to 1458.6)
						(57,214 to		
				2013	74,393	101,348)	1102.2	(820.9 to 1563.6)
	YLDs	Ischemic Stroke	Male	1990	188	(117 to 281)	9.8	(6.1 to 14.6)
				2013	343	(218 to 504)	9.9	(6.2 to 14.8)
			Female	1990	147	(90 to 214)	6.9	(4.2 to 10.1)
				2013	274	(166 to 402)	7.1	(4.3 to 10.4)
			Both	1990	336	(222 to 477)	8.2	(5.5 to 11.5)
				2013	617	(411 to 861)	8.4	(5.5 to 11.8)
		Hemorrhagic						
		Stroke	Male	1990	71	(43 to 106)	3	(1.8 to 4.4)
				2013	124	(74 to 194)	2.9	(1.7 to 4.5)
			Female	1990	47	(28 to 71)	1.8	(1.1 to 2.7)
				2013	83	(50 to 122)	1.8	(1.1 to 2.7)
			Both	1990	118	(76 to 169)	2.4	(1.5 to 3.3)
			Both	2013	208	(133 to 303)	2.3	(1.5 to 3.4)
Malaysia	Deaths	Ischemic Stroke	Male	1990	2,305	(1,737 to 2,800)	76.6	(57.6 to 94.2)
				2013	4,323	(3,409 to 5,933)	57.3	(45.0 to 79.1)
			Female	1990	2,500	(1,811 to 2,916)	82.2	(59.9 to 96.6)
				2013	3,335	(2,696 to 4,426)	51	(41.3 to 67.4)
			Both	1990	4,806	(3,865 to 5,437)	79.7	(63.7 to 91.1)
				2013	7,658	(6,486 to 9,931)	53.9	(45.4 to 70.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,445	(1,908 to 2,965)	61.4	(46.1 to 75.0)
				2013	4,227	(3,288 to 5,376)	45.5	(34.8 to 58.1)
			Female	1990	2,442	(1,796 to 2,803)	63	(44.6 to 73.9)

				2013	3,046	(2,443 to 4,192)	35.8	(28.2 to 50.1)
			Both	1990	4,887	(3,982 to 5,576)	62.3	(50.1 to 71.6)
				2013	7,274	(6,194 to 8,956)	40.5	(34.0 to 49.9)
	DALYs	Ischemic Stroke	Male	1990	42,812	(31,919 to 51,986)	1158.2	(870.3 to 1403.2)
						(61,917 to		
				2013	79,368	108,129)	879	(694.3 to 1199.3)
			Female	1990	40,283	(29,777 to 47,078)	1108.9	(813.2 to 1290.2)
				2013	53,402	(43,014 to 71,146)	650.3	(532.3 to 857.2)
			Both	1990	83,096	(67,089 to 94,748)	1135.3	(915.9 to 1285.2)
						(112,624 to		
				2013	132,770	171,026)	761.6	(647.1 to 965.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	71,465	(57,505 to 85,613)	1362.6	(1068.8 to 1645.2)
						(88,561 to		
				2013	112,558	142,011)	1003.9	(790.7 to 1272.3)
			Female	1990	63,585	(47,817 to 72,197)	1286	(956.8 to 1470.1)
						(59,729 to		
				2013	74,047	100,271)	675.9	(547.9 to 915.3)
			5	4000	125.050	(112,095 to	4005.6	(4000 0 : 4504 0)
			Both	1990	135,050	153,031)	1325.6	(1090.9 to 1504.9)
				2012	100 005	(159,669 to	025.4	/715 0 to 1022 0\
	V/I D :	Laster and a Charle	D.A. L.	2013	186,605	228,120)	835.4	(715.8 to 1022.9)
	YLDs	Ischemic Stroke	Male	1990	1,052	(689 to 1,526)	27.3	(17.5 to 39.8)
				2013	2,965	(1,938 to 4,226)	32.6	(21.5 to 46.1)
			Female	1990	1,006	(672 to 1,442)	25.7	(16.9 to 37.6)
				2013	2,457	(1,598 to 3,524)	26.4	(17.1 to 37.8)

	1	T					ı	
			Both	1990	2,059	(1,373 to 2,891)	26.5	(17.8 to 37.1)
				2013	5,423	(3,649 to 7,455)	29.5	(20.0 to 40.5)
		Hemorrhagic						
		Stroke	Male	1990	1,299	(832 to 1,858)	29.7	(19.1 to 42.7)
				2013	3,401	(2,232 to 4,813)	34.5	(22.6 to 49.3)
			Female	1990	1,079	(710 to 1,496)	24.9	(16.3 to 35.3)
				2013	2,535	(1,654 to 3,560)	24.9	(16.4 to 34.6)
			Both	1990	2,379	(1,595 to 3,217)	27.3	(18.7 to 37.2)
			Both	2013	5,936	(4,006 to 8,066)	29.7	(20.4 to 40.2)
Maldives	Deaths	Ischemic Stroke	Male	1990	44	(28 to 58)	134.3	(84.5 to 169.9)
				2013	51	(39 to 74)	59.3	(45.2 to 84.5)
			Female	1990	16	(7 to 31)	85.6	(44.1 to 144.9)
				2013	40	(23 to 51)	61	(34.8 to 76.8)
			Both	1990	61	(40 to 81)	116.2	(81.6 to 147.1)
				2013	92	(69 to 111)	59.5	(43.8 to 71.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	52	(33 to 68)	88.4	(52.7 to 123.7)
				2013	22	(16 to 36)	21.6	(14.9 to 37.5)
			Female	1990	22	(14 to 41)	59.7	(41.1 to 98.5)
				2013	17	(11 to 22)	21.4	(13.5 to 27.3)
			Both	1990	74	(57 to 90)	76.5	(58.5 to 98.4)
				2013	40	(31 to 54)	21.2	(16.2 to 30.3)
	DALYs	Ischemic Stroke	Male	1990	937	(581 to 1,279)	1978.9	(1262.3 to 2545.1)
				2013	751	(579 to 1,034)	763.2	(588.6 to 1066.6)
			Female	1990	365	(127 to 751)	1221.7	(563.6 to 2143.4)
				2013	536	(312 to 705)	664.3	(386.4 to 853.3)
			Both	1990	1,302	(770 to 1,806)	1684.1	(1116.4 to 2193.4)
				2013	1,288	(986 to 1,562)	711.6	(543.5 to 858.3)
	<u>'</u>				Counts, All	Ages	Rate	Per 100,000

							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	2,141	(1,500 to 2,663)	2349.4	(1505.1 to 3016.5)
				2013	750	(584 to 1,004)	561.8	(426.9 to 812.8)
			Female	1990	802	(448 to 1,735)	1406.8	(939.0 to 2473.7)
				2013	479	(310 to 597)	423.2	(277.0 to 525.6)
			Both	1990	2,943	(2,330 to 3,708)	1929.4	(1494.2 to 2318.8)
				2013	1,229	(1,001 to 1,477)	492.1	(396.8 to 612.5)
	YLDs	Ischemic Stroke	Male	1990	12	(8 to 18)	26.4	(16.7 to 37.7)
				2013	28	(18 to 42)	28.9	(18.0 to 42.8)
			Female	1990	7	(5 to 10)	24.2	(15.8 to 34.0)
				2013	21	(14 to 30)	24.2	(15.8 to 34.9)
			Both	1990	20	(13 to 28)	25.5	(17.1 to 35.2)
				2013	49	(34 to 70)	26.6	(18.2 to 36.9)
		Hemorrhagic						
		Stroke	Male	1990	15	(10 to 21)	29.1	(19.0 to 40.7)
				2013	32	(21 to 48)	31	(19.8 to 45.8)
			Female	1990	8	(5 to 11)	23.2	(15.3 to 31.9)
				2013	22	(14 to 30)	22.7	(15.2 to 31.8)
			Both	1990	24	(16 to 33)	26.6	(17.9 to 36.0)
			Both	2013	54	(37 to 75)	27	(18.2 to 37.4)
Mali	Deaths	Ischemic Stroke	Male	1990	669	(408 to 903)	52	(32.0 to 70.0)
				2013	1,394	(808 to 1,893)	79.1	(46.0 to 106.3)
			Female	1990	1,316	(787 to 1,783)	88.7	(52.2 to 116.1)
				2013	2,651	(1,359 to 3,578)	125	(65.4 to 166.0)
			Both	1990	1,986	(1,420 to 2,442)	71.9	(50.8 to 88.1)
				2013	4,046	(2,443 to 5,114)	104.2	(62.4 to 132.0)
	Deaths	Hemorrhagic	Male	1990	667	(373 to 869)	39.7	(20.0 to 54.3)

		Stroke						
				2013	1,006	(654 to 1,507)	38.4	(22.0 to 64.6)
			Female	1990	1,397	(877 to 1,835)	75.3	(44.4 to 101.6)
				2013	1,965	(1,331 to 3,048)	66.6	(42.8 to 110.8)
			Both	1990	2,065	(1,471 to 2,481)	58.9	(39.3 to 73.7)
				2013	2,972	(2,102 to 4,168)	53.6	(35.3 to 82.6)
	DALYs	Ischemic Stroke	Male	1990	14,890	(9,484 to 20,607)	854.5	(515.2 to 1155.4)
				2013	29,175	(16,320 to 40,231)	1239.6	(708.7 to 1691.6)
			Female	1990	27,869	(15,904 to 39,410)	1438	(852.0 to 1962.9)
				2013	52,521	(25,758 to 72,730)	1943.7	(982.3 to 2634.6)
			Both	1990	42,760	(30,614 to 53,903)	1168.9	(831.0 to 1440.5)
						(49,270 to		
				2013	81,696	104,739)	1625.3	(972.4 to 2059.2)
		Hemorrhagic						
		Stroke	Male	1990	22,655	(14,204 to 28,866)	883.9	(496.0 to 1152.5)
				2013	37,161	(25,697 to 51,535)	872.1	(557.4 to 1328.8)
					Counts, All	Ages	Rate	Per 100,000
		a: 1 =				0=0(1)	Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	42,362	(27,571 to 54,558)	1677.1	(1052.3 to 2191.1)
				2013	63,862	(44,518 to 93,242)	1572.3	(1072.1 to 2392.3)
			Both	1990	65,017	(47,766 to 77,467)	1309.5	(933.5 to 1579.4)
						(76,440 to		
				2013	101,023	132,204)	1244.1	(884.1 to 1750.7)
	YLDs	Ischemic Stroke	Male	1990	293	(191 to 414)	15.9	(10.6 to 22.5)
				2013	422	(273 to 624)	14.8	(9.5 to 21.7)
			Female	1990	250	(166 to 357)	11.8	(7.9 to 16.7)
				2013	401	(252 to 567)	12.3	(7.8 to 17.6)
			Both	1990	543	(368 to 740)	13.7	(9.4 to 18.5)

				2013	824	(555 to 1,141)	13.4	(9.0 to 18.6)
		Hemorrhagic			0_ :	(000 to 1)1 :-)	2011	(5.5 to 25.5)
		Stroke	Male	1990	88	(56 to 127)	4.3	(2.7 to 6.3)
				2013	137	(85 to 196)	3.8	(2.4 to 5.6)
			Female	1990	69	(43 to 100)	3	(1.8 to 4.4)
				2013	104	(63 to 154)	2.7	(1.7 to 4.1)
			Both	1990	157	(104 to 217)	3.6	(2.4 to 4.9)
			Both	2013	241	(160 to 339)	3.3	(2.1 to 4.6)
Malta	Deaths	Ischemic Stroke	Male	1990	65	(52 to 84)	50.3	(40.3 to 64.7)
				2013	72	(54 to 91)	29.6	(21.9 to 37.3)
			Female	1990	91	(72 to 113)	46.6	(36.9 to 57.5)
				2013	108	(78 to 133)	30.3	(21.7 to 37.5)
			Both	1990	157	(137 to 189)	48.3	(42.2 to 58.1)
				2013	181	(143 to 213)	30.3	(23.5 to 35.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	50	(37 to 63)	36.1	(26.9 to 45.7)
				2013	56	(41 to 71)	21.4	(15.3 to 27.4)
			Female	1990	67	(47 to 80)	34.3	(24.1 to 40.8)
				2013	67	(49 to 90)	18.9	(13.8 to 25.8)
			Both	1990	117	(94 to 135)	35.2	(28.2 to 40.8)
				2013	123	(100 to 150)	20.2	(16.3 to 24.9)
	DALYs	Ischemic Stroke	Male	1990	992	(809 to 1,258)	698.9	(569.7 to 888.3)
				2013	1,153	(893 to 1,421)	401.1	(313.0 to 492.6)
			Female	1990	1,114	(906 to 1,350)	567.5	(460.9 to 686.2)
				2013	1,168	(906 to 1,420)	318	(247.1 to 387.7)
			Both	1990	2,107	(1,856 to 2,495)	624.5	(551.3 to 738.1)
				2013	2,321	(1,940 to 2,718)	357.8	(297.1 to 418.8)
		Hemorrhagic Stroke	Male	1990	952	(742 to 1,177)	619.9	(476.8 to 771.5)

	1		ı ı			1	1	
				2013	960	(710 to 1,203)	330.8	(246.9 to 414.7)
			Female	1990	1,032	(763 to 1,194)	527.8	(390.3 to 610.6)
				2013	814	(624 to 1,074)	239.3	(185.5 to 313.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	1,984	(1,652 to 2,262)	569.4	(472.7 to 650.3)
				2013	1,775	(1,481 to 2,120)	283.3	(236.1 to 336.6)
	YLDs	Ischemic Stroke	Male	1990	122	(78 to 167)	82	(52.2 to 112.1)
				2013	273	(168 to 405)	88.8	(54.3 to 130.3)
			Female	1990	117	(79 to 164)	59.8	(40.6 to 83.4)
				2013	208	(134 to 299)	58.3	(37.9 to 83.1)
			Both	1990	239	(164 to 321)	69.3	(47.4 to 93.0)
				2013	482	(319 to 676)	72.3	(47.9 to 101.2)
		Hemorrhagic						
		Stroke	Male	1990	27	(14 to 48)	17.5	(9.2 to 31.5)
				2013	50	(25 to 90)	16.9	(8.6 to 29.8)
			Female	1990	25	(14 to 41)	12.8	(7.1 to 20.9)
				2013	41	(23 to 71)	12.6	(7.2 to 20.7)
			Both	1990	52	(32 to 79)	14.8	(9.3 to 22.4)
			Both	2013	92	(55 to 142)	14.5	(8.9 to 21.9)
Marshall Islands	Deaths	Ischemic Stroke	Male	1990	5	(2 to 7)	90.8	(48.4 to 121.6)
				2013	9	(5 to 13)	85.4	(45.3 to 121.4)
			Female	1990	2	(1 to 4)	46.6	(32.7 to 68.0)
				2013	7	(3 to 10)	54.6	(28.5 to 79.1)
			Both	1990	8	(5 to 10)	66.2	(48.8 to 83.4)
				2013	16	(10 to 22)	68.1	(43.1 to 88.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	10	(6 to 13)	128.3	(80.8 to 168.1)

				2013	19	(11 to 28)	124.1	(75.9 to 172.9)
			Female	1990	4	(3 to 6)	62.7	(44.9 to 88.7)
				2013	12	(6 to 17)	76.6	(41.7 to 111.9)
			Both	1990	15	(11 to 19)	93.5	(71.6 to 118.6)
				2013	31	(20 to 42)	98.7	(64.7 to 131.6)
	DALYs	Ischemic Stroke	Male	1990	125	(66 to 181)	1532.7	(827.1 to 2085.3)
				2013	220	(116 to 332)	1426.3	(776.8 to 2052.9)
			Female	1990	60	(39 to 87)	731.9	(506.4 to 1013.1)
				2013	141	(62 to 224)	851.6	(407.4 to 1289.9)
			Both	1990	186	(117 to 251)	1106.5	(752.0 to 1428.7)
				2013	361	(207 to 493)	1116.1	(672.3 to 1474.3)
		Hemorrhagic						
		Stroke	Male	1990	347	(247 to 455)	2925.1	(1940.5 to 3840.5)
				2013	601	(371 to 894)	2852.8	(1756.1 to 4172.1)
			Female	1990	142	(105 to 186)	1273.8	(939.3 to 1737.6)
				2013	318	(161 to 476)	1557.2	(829.3 to 2295.7)
			Both	1990	490	(384 to 605)	2084.5	(1599.1 to 2598.2)
				2013	919	(592 to 1,266)	2185.9	(1425.3 to 2971.5)
	YLDs	Ischemic Stroke	Male	1990	2	(1 to 2)	28.2	(18.2 to 39.7)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	3	(2 to 4)	24.5	(15.8 to 34.4)
			Female	1990	1	(1 to 2)	20.1	(12.7 to 28.5)
				2013	3	(2 to 4)	19.2	(12.5 to 27.7)
			Both	1990	3	(2 to 5)	23.7	(15.7 to 32.7)
				2013	6	(4 to 9)	21.6	(14.7 to 29.6)
		Hemorrhagic						
		Stroke	Male	1990	2	(1 to 4)	33.2	(22.2 to 47.9)

				2013	4	(3 to 6)	29.3	(19.1 to 41.1)
			Female	1990	1	(1 to 2)	21.4	(14.0 to 29.7)
				2013	3	(2 to 5)	21	(13.5 to 30.1)
			Both	1990	4	(3 to 6)	26.9	(18.1 to 37.2)
			Both	2013	8	(5 to 11)	24.8	(16.7 to 33.3)
Mauritania	Deaths	Ischemic Stroke	Male	1990	180	(111 to 252)	70	(44.9 to 95.4)
				2013	545	(288 to 738)	105.9	(59.5 to 139.9)
			Female	1990	265	(187 to 358)	85	(61.4 to 117.5)
				2013	710	(404 to 973)	113.3	(65.6 to 151.8)
			Both	1990	445	(355 to 555)	78.6	(62.9 to 98.4)
				2013	1,256	(810 to 1,599)	110.5	(72.0 to 138.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	165	(109 to 220)	49.5	(30.6 to 66.9)
				2013	337	(195 to 547)	47.7	(25.7 to 85.9)
			Female	1990	250	(183 to 359)	66.4	(48.0 to 96.3)
				2013	446	(264 to 858)	57.1	(32.9 to 118.5)
			Both	1990	416	(322 to 529)	59.1	(44.8 to 76.0)
				2013	784	(499 to 1,332)	53.1	(31.8 to 97.6)
	DALYs	Ischemic Stroke	Male	1990	3,894	(2,351 to 5,536)	1123.6	(690.1 to 1556.5)
				2013	11,347	(5,797 to 16,019)	1652.3	(879.4 to 2232.6)
			Female	1990	5,144	(3,470 to 6,915)	1299.7	(909.3 to 1753.2)
				2013	12,638	(6,830 to 17,708)	1638.3	(921.2 to 2251.3)
			Both	1990	9,039	(7,042 to 11,435)	1224.4	(974.8 to 1524.7)
				2013	23,986	(14,540 to 31,252)	1653.5	(1061.4 to 2110.7)
		Hemorrhagic						
		Stroke	Male	1990	4,996	(3,509 to 6,452)	1025.6	(675.1 to 1355.4)
				2013	10,347	(6,317 to 15,479)	1011.8	(586.7 to 1645.1)
			Female	1990	6,478	(4,725 to 9,079)	1284.3	(937.9 to 1845.7)
				2013	11,452	(7,163 to 20,478)	1124.3	(675.5 to 2118.2)

			Both	1990	11,474	(9,040 to 14,349)	1170.9	(907.6 to 1477.6)
				2013	21,800	(14,611 to 33,603)	1077.5	(693.4 to 1803.8)
	YLDs	Ischemic Stroke	Male	1990	64	(42 to 90)	15.4	(10.2 to 21.3)
				2013	146	(91 to 209)	16.4	(10.5 to 23.4)
			Female	1990	56	(35 to 83)	11.7	(7.4 to 17.0)
				2013	121	(79 to 174)	12.1	(8.0 to 17.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	121	(81 to 167)	13.4	(9.0 to 18.3)
				2013	268	(177 to 377)	14	(9.3 to 19.6)
		Hemorrhagic						
		Stroke	Male	1990	21	(13 to 30)	4.2	(2.6 to 6.1)
				2013	44	(26 to 66)	4.1	(2.5 to 6.1)
			Female	1990	16	(10 to 24)	3	(1.8 to 4.3)
				2013	32	(20 to 48)	2.8	(1.8 to 4.2)
			Both	1990	37	(24 to 51)	3.5	(2.4 to 4.9)
			Both	2013	76	(49 to 109)	3.4	(2.2 to 4.9)
Mauritius	Deaths	Ischemic Stroke	Male	1990	302	(257 to 373)	172.7	(145.0 to 210.6)
				2013	484	(365 to 572)	114.5	(87.1 to 134.9)
			Female	1990	277	(243 to 349)	108	(94.8 to 136.7)
				2013	415	(335 to 552)	69.5	(56.5 to 92.0)
			Both	1990	580	(520 to 695)	133.7	(119.9 to 161.5)
				2013	899	(770 to 1,038)	88.5	(76.0 to 102.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	212	(162 to 257)	79	(57.6 to 96.8)
				2013	249	(183 to 331)	41.2	(30.3 to 55.9)
			Female	1990	161	(115 to 183)	50.1	(34.3 to 57.8)
				2013	137	(100 to 227)	20.1	(14.5 to 33.4)

			Both	1990	373	(296 to 427)	63.1	(48.2 to 73.0)
				2013	386	(306 to 497)	29.9	(23.6 to 39.3)
	DALYs	Ischemic Stroke	Male	1990	6,052	(5,130 to 7,508)	2685.9	(2288.0 to 3294.0)
				2013	8,241	(6,216 to 9,995)	1638	(1234.2 to 1944.9)
			Female	1990	4,153	(3,535 to 5,156)	1407	(1218.5 to 1757.3)
				2013	5,040	(4,049 to 6,930)	800.5	(646.5 to 1097.1)
			Both	1990	10,206	(9,067 to 12,240)	1940.8	(1735.9 to 2311.8)
				2013	13,282	(11,112 to 15,403)	1167.5	(992.5 to 1346.8)
		Hemorrhagic						
		Stroke	Male	1990	6,169	(4,946 to 7,473)	1872.1	(1453.4 to 2263.9)
				2013	7,128	(5,295 to 9,238)	1062.2	(797.8 to 1386.7)
			Female	1990	4,082	(3,114 to 4,569)	1084.8	(808.7 to 1221.4)
				2013	3,422	(2,568 to 5,361)	473.3	(357.0 to 736.7)
			Both	1990	10,251	(8,515 to 11,664)	1446.7	(1169.7 to 1651.5)
				2013	10,550	(8,498 to 13,255)	753.6	(607.9 to 949.1)
	YLDs	Ischemic Stroke	Male	1990	65	(41 to 94)	27.3	(17.4 to 38.8)
				2013	158	(105 to 223)	29.6	(19.6 to 42.1)
			Female	1990	74	(48 to 105)	23.3	(15.1 to 33.3)
				2013	170	(114 to 240)	25.5	(17.0 to 36.1)
			Both	1990	140	(94 to 193)	25	(17.0 to 34.3)
				2013	329	(224 to 444)	27.3	(18.7 to 37.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic				(=		(
		Stroke	Male	1990	82	(54 to 117)	30.1	(19.5 to 42.2)
				2013	181	(120 to 261)	31.9	(21.1 to 45.6)
			Female	1990	77	(50 to 110)	22.7	(14.8 to 32.4)

				2013	163	(109 to 225)	23.9	(15.9 to 33.1)
			Both	1990	160	(107 to 217)	25.9	(17.5 to 34.8)
			Both	2013	344	(237 to 470)	27.5	(19.1 to 37.6)
Mexico	Deaths	Ischemic Stroke	Male	1990	3,545	(2,688 to 4,254)	22	(16.5 to 26.3)
				2013	5,690	(4,488 to 8,068)	16.7	(13.2 to 23.8)
			Female	1990	4,229	(3,034 to 5,146)	20.7	(14.9 to 25.2)
				2013	6,527	(5,017 to 8,959)	14.2	(10.9 to 19.5)
			Both	1990	7,775	(6,381 to 8,948)	21.3	(17.6 to 24.4)
				2013	12,218	(10,715 to 15,926)	15.4	(13.4 to 19.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	6,853	(5,751 to 8,166)	38.5	(31.9 to 45.8)
				2013	13,695	(11,294 to 16,775)	38	(31.2 to 46.7)
			Female	1990	7,980	(6,331 to 9,739)	36.5	(29.1 to 45.0)
				2013	14,805	(11,564 to 18,475)	31.5	(24.5 to 39.0)
			Both	1990	14,833	(13,112 to 17,254)	37.4	(33.1 to 43.5)
				2013	28,500	(24,803 to 32,864)	34.4	(29.8 to 39.5)
	DALYs	Ischemic Stroke	Male	1990	64,366	(49,820 to 77,631)	332.4	(256.8 to 400.0)
						(74,290 to		
				2013	92,942	127,285)	242.6	(194.7 to 334.8)
			Female	1990	66,451	(47,483 to 80,671)	287.5	(206.5 to 349.2)
						(69,561 to		(
				2013	88,906	123,319)	184.8	(144.8 to 255.5)
			Both	1990	120 017	(108,175 to 149,141)	307.8	(252.1 +0.252.0)
			BOILL	1990	130,817	(157,868 to	307.8	(253.1 to 353.0)
				2013	181,849	239,592)	210.9	(184.2 to 279.1)
		Hemorrhagic				(141,188 to		(== ::= := = ::=)
		Stroke	Male	1990	167,065	196,415)	720.5	(605.4 to 847.1)
						(237,974 to		
				2013	281,834	340,374)	672	(564.8 to 809.8)

						(139,878 to		
			Female	1990	174,348	212,233)	662.4	(534.8 to 799.9)
						(215,903 to		
				2013	260,143	331,984)	510.8	(421.5 to 650.4)
				1000		(302,153 to		(640 = 4 = 700 = 1)
			Both	1990	341,413	386,699)	688	(610.5 to 782.5)
				2013	F41 070	(486,778 to 621,913)	583.9	/F21 2 to 672 2\
	VI Da	la ab anaia Chualta	Mala		541,978			(521.3 to 672.2)
	YLDs	Ischemic Stroke	Male	1990	1,273	(883 to 1,708)	6.6	(4.6 to 8.8)
				2013	2,806	(1,954 to 3,722)	7.1	(5.0 to 9.5)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	moacuro	Stroka Typa	cov	voor	Mean	95% UI	Mean	95% UI
Country	measure	Stroke Type	Sex	year				
			Female	1990	1,194	(819 to 1,603)	5.3	(3.6 to 7.1)
			_	2013	2,782	(1,926 to 3,742)	5.7	(3.9 to 7.6)
			Both	1990	2,468	(1,710 to 3,300)	5.9	(4.1 to 7.8)
				2013	5,589	(3,921 to 7,481)	6.3	(4.4 to 8.5)
		Hemorrhagic				()		(10.7)
		Stroke	Male	1990	4,310	(2,976 to 5,772)	19.7	(13.7 to 26.2)
				2013	8,931	(6,201 to 11,841)	20.9	(14.6 to 27.5)
			Female	1990	3,204	(2,213 to 4,265)	12.8	(8.9 to 17.1)
				2013	6,539	(4,505 to 8,695)	12.5	(8.6 to 16.5)
			Both	1990	7,515	(5,199 to 10,015)	16	(11.1 to 21.3)
			Both	2013	15,471	(10,773 to 20,490)	16.2	(11.3 to 21.4)
Moldova	Deaths	Ischemic Stroke	Male	1990	1,301	(1,067 to 1,619)	106.3	(86.8 to 130.3)
				2013	1,511	(1,156 to 1,787)	105.6	(81.0 to 125.8)
			Female	1990	1,958	(1,524 to 2,442)	95.4	(74.2 to 116.6)
				2013	1,841	(1,249 to 2,106)	71	(48.1 to 81.3)
			Both	1990	3,259	(2,834 to 3,881)	99.9	(85.8 to 118.0)
				2013	3,352	(2,552 to 3,767)	84.2	(64.4 to 94.7)

		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,243	(857 to 1,496)	83.4	(56.9 to 101.4)
				2013	1,326	(1,081 to 1,711)	78.7	(64.2 to 103.8)
			Female	1990	1,691	(1,139 to 1,967)	75	(49.4 to 87.4)
				2013	1,375	(1,162 to 2,012)	53.8	(45.5 to 78.4)
			Both	1990	2,934	(2,187 to 3,275)	78.6	(58.0 to 88.3)
				2013	2,701	(2,333 to 3,497)	64.5	(55.6 to 83.9)
	DALYs	Ischemic Stroke	Male	1990	23,366	(19,230 to 29,831)	1623.2	(1338.4 to 2024.7)
				2013	26,167	(19,850 to 31,037)	1598.8	(1215.5 to 1884.2)
			Female	1990	29,223	(23,293 to 37,959)	1314.9	(1053.1 to 1686.2)
				2013	23,880	(15,980 to 27,126)	926.2	(619.2 to 1054.3)
			Both	1990	52,589	(46,194 to 65,309)	1439.7	(1264.0 to 1752.7)
				2013	50,048	(38,110 to 56,792)	1196.6	(910.8 to 1356.4)
		Hemorrhagic						
		Stroke	Male	1990	29,823	(21,403 to 35,779)	1752.3	(1235.0 to 2099.0)
				2013	30,926	(25,372 to 38,974)	1653	(1358.7 to 2091.5)
			Female	1990	35,513	(25,467 to 41,219)	1498.7	(1067.8 to 1747.9)
				2013	24,758	(21,257 to 35,224)	983.8	(845.1 to 1397.8)
			Both	1990	65,337	(50,643 to 72,798)	1607.4	(1236.7 to 1793.8)
				2013	55,684	(48,502 to 69,895)	1279.9	(1114.3 to 1609.6)
	YLDs	Ischemic Stroke	Male	1990	671	(441 to 954)	38.5	(25.5 to 54.2)
				2013	761	(485 to 1,099)	41.2	(26.5 to 58.8)
			Female	1990	556	(334 to 836)	23.8	(14.4 to 35.6)
				2013	594	(381 to 848)	24.4	(15.6 to 34.6)
			Both	1990	1,227	(816 to 1,724)	30.1	(20.1 to 42.1)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,355	(905 to 1,862)	31.7	(21.0 to 43.6)

			Female	1990	23,653	(20,274 to 28,484)	3772.3	(3236.7 to 4578.8)
				2013	37,845	(28,684 to 46,881)	3858.3	(2928.0 to 4689.5)
			Both	1990	47,851	(42,350 to 53,940)	4018.4	(3564.0 to 4530.3)
				2013	81,732	(69,927 to 94,756)	4464.9	(3783.6 to 5103.7)
	YLDs	Ischemic Stroke	Male	1990	43	(28 to 63)	8.3	(5.5 to 12.1)
				2013	61	(36 to 92)	7.7	(4.5 to 11.7)
			Female	1990	42	(26 to 61)	7.3	(4.5 to 10.5)
				2013	68	(42 to 101)	7.4	(4.6 to 11.1)
			Both	1990	86	(56 to 118)	7.8	(5.1 to 10.8)
				2013	130	(84 to 182)	7.6	(4.9 to 10.6)
		Hemorrhagic						
		Stroke	Male	1990	55	(35 to 78)	9.2	(5.9 to 13.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	79	(48 to 116)	8.1	(4.9 to 11.7)
			Female	1990	33	(21 to 47)	5.1	(3.2 to 7.3)
				2013	50	(31 to 74)	4.8	(3.0 to 7.2)
			Both	1990	88	(58 to 121)	7	(4.7 to 9.7)
			Both	2013	130	(82 to 183)	6.3	(4.1 to 8.8)
Montenegro	Deaths	Ischemic Stroke	Male	1990	86	(51 to 155)	41.6	(24.9 to 74.4)
				2013	188	(123 to 243)	59	(38.6 to 76.9)
				2013	100	(123 (0 243)	39	(30.0 to 70.3)
			Female	1990	109	(71 to 191)	32.3	(21.1 to 56.3)
			Female			,		, ,
			Female Both	1990	109	(71 to 191)	32.3	(21.1 to 56.3)
				1990 2013	109 227	(71 to 191) (131 to 288)	32.3 49.9	(21.1 to 56.3) (28.9 to 63.4)
		Hemorrhagic		1990 2013 1990	109 227 195	(71 to 191) (131 to 288) (133 to 333)	32.3 49.9 36	(21.1 to 56.3) (28.9 to 63.4) (24.7 to 61.1)
	Deaths	Hemorrhagic Stroke		1990 2013 1990	109 227 195	(71 to 191) (131 to 288) (133 to 333)	32.3 49.9 36	(21.1 to 56.3) (28.9 to 63.4) (24.7 to 61.1)

Female 1990 658 (504 to 821) 197 (150.0 to 246.3)							
Both 1990		Female	1990	658	(504 to 821)	197	(150.0 to 246.3)
DALYS Ischemic Stroke Male 1990 1,468 (857 to 2,727) 627.1 (156.3 to 203.7) DALYS Ischemic Stroke Male 1990 1,468 (857 to 2,727) 627.1 (373.1 to 1128.6) DALYS Ischemic Stroke Male 1990 1,468 (857 to 2,727) 627.1 (373.1 to 1128.6) Emale 1990 1,392 (920 to 2,449) 435.5 (287.4 to 766.7) DALYS Ischemic Stroke Male 1990 2,860 (1,910 to 4,832) 517.4 (346.5 to 873.1) Hemorrhagic Stroke Male 1990 8,935 (6,698 to 11,273) 3639.1 (2756.9 to 4552.5) DALYS Ischemic Stroke Male 1990 9,442 (6,763 to 11,954) 2988 (2141.6 to 3796.7) Both 1990 18,378 (14,564 to 22,823) 2535.5 (2213.7 to 2898.1) YLDS Ischemic Stroke Male 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Female 1990 15 (9 to 23) 5.8 (3.7 to 8.4) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) Hemorrhagic Stroke Male 1990 15 (9 to 24) 4 (2.5 to 6.0)			2013	800	(681 to 958)	177.8	(150.7 to 213.8)
DALYS Ischemic Stroke Male 1990		Both	1990	1,102	(904 to 1,296)	202.4	(167.0 to 237.9)
2013 2,890 (1,965 to 3,743) 828.8 (566.1 to 1057.9)			2013	1,365	(1,206 to 1,565)	177.1	(156.3 to 203.7)
Female 1990 1,392 (920 to 2,449) 435.5 (287.4 to 766.7) 2013 2,858 (1,711 to 3,583) 626.2 (379.5 to 783.5) Both 1990 2,860 (1,910 to 4,832) 517.4 (346.5 to 873.1) 2013 5,748 (4,017 to 6,970) 717.6 (501.2 to 867.9) Hemorrhagic Stroke Male 1990 8,935 (6,698 to 11,273) 3639.1 (2756.9 to 4552.5) 2013 9,432 (7,535 to 11,334) 2688.8 (2160.1 to 3233.7) Female 1990 9,442 (6,763 to 11,954) 2988 (2141.6 to 3796.7) Both 1990 18,378 (14,564 to 21,759) 3295.4 (2628.5 to 3900.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (213.7 to 2898.1) YLDs Ischemic Stroke Male 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Female 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1)	DALYs Ischemic Stroke	Male	1990	1,468	(857 to 2,727)	627.1	(373.1 to 1128.6)
Both 1990 2,860 (1,711 to 3,583) 626.2 (379.5 to 783.5)			2013	2,890	(1,965 to 3,743)	828.8	(566.1 to 1057.9)
Both 1990 2,860 (1,910 to 4,832) 517.4 (346.5 to 873.1)		Female	1990	1,392	(920 to 2,449)	435.5	(287.4 to 766.7)
Hemorrhagic Stroke Male 1990 8,935 (6,698 to 11,273) 3639.1 (2756.9 to 4552.5)			2013	2,858	(1,711 to 3,583)	626.2	(379.5 to 783.5)
Hemorrhagic Stroke Male 1990 8,935 (6,698 to 11,273) 3639.1 (2756.9 to 4552.5)		Both	1990	2,860	(1,910 to 4,832)	517.4	(346.5 to 873.1)
Stroke Male 1990 8,935 (6,698 to 11,273) 3639.1 (2756.9 to 4552.5) 2013 9,432 (7,535 to 11,334) 2688.8 (2160.1 to 3233.7) Female 1990 9,442 (6,763 to 11,954) 2988 (2141.6 to 3796.7) Both 1990 18,378 (14,564 to 21,759) 3295.4 (2628.5 to 3900.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (2213.7 to 2898.1) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.5 to 33.4) YLDs Ischemic Stroke Male 1990 47 (30 to 67) 14.9 (9.6 to 21.3) YLDs Both 1990 47 (30 to 67) 14.9 (9.6 to 21.3) YLDs Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) YLDs Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) YLDs <td< td=""><td></td><td></td><td>2013</td><td>5,748</td><td>(4,017 to 6,970)</td><td>717.6</td><td>(501.2 to 867.9)</td></td<>			2013	5,748	(4,017 to 6,970)	717.6	(501.2 to 867.9)
2013 9,432 (7,535 to 11,334) 2688.8 (2160.1 to 3233.7)		Male	1990	8.935	(6.698 to 11.273)	3639.1	(2756.9 to 4552.5)
Female 1990 9,442 (6,763 to 11,954) 2988 (2141.6 to 3796.7) Both 1990 18,378 (14,564 to 21,759) 3295.4 (2628.5 to 3900.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.5 to 33.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.7 to 35.0) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Stroke 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 6 1990				†			` '
Both 1990 18,378 (14,564 to 21,759) 3295.4 (2628.5 to 3900.4)		Female		· -			` '
Both 1990 18,378 (14,564 to 21,759) 3295.4 (2628.5 to 3900.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.5 to 33.4) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.7 to 35.0) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) Stroke Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.0)				·	, , , , , , , , , , , , , , , , , , , ,		,
YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (2213.7 to 2898.1) YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.5 to 33.4) 2013 92 (57 to 129) 25 (15.7 to 35.0) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) 2013 63 (40 to 91) 14.8 (9.5 to 21.1) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic 2013 156 (103 to 218) 19.5 (13.0 to 27.1) Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)		Both					, ,
YLDs Ischemic Stroke Male 1990 60 (39 to 85) 23.5 (15.5 to 33.4) 2013 92 (57 to 129) 25 (15.7 to 35.0) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) 2013 63 (40 to 91) 14.8 (9.5 to 21.1) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Lead of the control of the cont			2013				, ,
2013 92 (57 to 129) 25 (15.7 to 35.0) Female 1990 47 (30 to 67) 14.9 (9.6 to 21.3) 2013 63 (40 to 91) 14.8 (9.5 to 21.1) Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)	YLDs Ischemic Stroke	Male	1990				
2013 63 (40 to 91) 14.8 (9.5 to 21.1)			2013	92	(57 to 129)	25	·
Both 1990 107 (73 to 145) 18.7 (12.9 to 25.3) Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)		Female	1990	47	(30 to 67)	14.9	(9.6 to 21.3)
2013 156 (103 to 218) 19.5 (13.0 to 27.1)			2013	63	(40 to 91)	14.8	(9.5 to 21.1)
Hemorrhagic Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)		Both	1990	107	(73 to 145)	18.7	(12.9 to 25.3)
Stroke Male 1990 15 (9 to 23) 5.8 (3.7 to 8.4) 2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)			2013	156	(103 to 218)	19.5	(13.0 to 27.1)
2013 21 (12 to 32) 5.9 (3.5 to 9.0) Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)	Hemorrhagic						
Female 1990 13 (7 to 19) 4.1 (2.5 to 6.1) 2013 16 (9 to 24) 4 (2.5 to 6.0)	Stroke	Male	1990	†	· · · · · · · · · · · · · · · · · · ·		,
2013 16 (9 to 24) 4 (2.5 to 6.0)			2013	21	(12 to 32)	5.9	(3.5 to 9.0)
		Female	1990	+	(7 to 19)	4.1	(2.5 to 6.1)
Counts, All Ages Rate Per 100,000			2013	16	(9 to 24)	4	(2.5 to 6.0)
				Counts, All	Ages	Rate	Per 100,000

							Age- standardized		
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI	
			Both	1990	28	(18 to 40)	4.9	(3.2 to 6.8)	
			Both	2013	37	(23 to 53)	4.9	(3.1 to 7.0)	
Morocco	Deaths	Ischemic Stroke	Male	1990	2,233	(1,737 to 2,857)	68.6	(52.5 to 87.0)	
				2013	3,206	(2,438 to 4,271)	45.5	(34.7 to 59.4)	
			Female	1990	2,779	(2,045 to 3,271)	68.6	(49.8 to 82.6)	
				2013	4,744	(3,541 to 6,340)	50.2	(37.6 to 66.2)	
			Both	1990	5,012	(4,062 to 5,777)	68.8	(55.0 to 79.4)	
				2013	7,950	(6,526 to 9,828)	48.4	(39.8 to 59.5)	
	Deaths	Hemorrhagic Stroke	Male	1990	1,962	(1,471 to 2,524)	37.8	(26.7 to 50.7)	
				2013	2,367	(1,768 to 3,248)	25.9	(19.2 to 35.8)	
			Female	1990	2,530	(1,990 to 3,151)	43.9	(33.4 to 55.5)	
				2013	3,244	(2,442 to 4,551)	29.5	(21.9 to 41.3)	
			Both	1990	4,493	(3,711 to 5,329)	41.4	(32.7 to 50.3)	
				2013	5,612	(4,571 to 7,206)	28	(22.7 to 36.3)	
	DALYs	Ischemic Stroke	Male	1990	46,640	(35,787 to 61,305)	1010.3	(785.2 to 1287.6)	
				2013	57,113	(42,672 to 77,032)	647.4	(493.4 to 864.3)	
			Female	1990	48,803	(36,396 to 59,361)	917.9	(682.4 to 1081.7)	
				2013	71,005	(52,348 to 96,138)	663.4	(492.9 to 893.9)	
			Both	1990	95,444	(78,809 to 113,768)	961.2	(783.8 to 1110.0)	
				2013	128,118	(104,463 to 160,843)	658.6	(536.8 to 822.4)	
		Hemorrhagic Stroke	Male	1990	76,909	(58,307 to 101,217)	895.4	(675.7 to 1148.2)	
				2013	68,989	(51,782 to 95,731)	557.3	(417.8 to 769.2)	
			Female	1990	82,937	(63,269 to 109,622)	964.5	(762.5 to 1190.8)	

Company								
2013 2,548 (1,653 to 3,679) 50.7 (31.9 to 76.3)				2013	2,864	(1,960 to 4,084)	66.5	(43.1 to 97.8)
Both 1990 2,510 (1,699 to 3,295) 48.9 (30.5 to 68.5)			Female	1990	1,345	(840 to 1,925)	48.2	(28.2 to 74.4)
DALYS Ischemic Stroke Male 1990 16,798 (6,200 to 27,371) 708.3 (242.8 to 1149.1)				2013	2,548	(1,653 to 3,679)	50.7	(31.9 to 76.3)
DALYS Ischemic Stroke Male 1990 16,798 (6,200 to 27,371) 708.3 (242.8 to 1149.1)			Both	1990	2,510	(1,699 to 3,295)	48.9	(30.5 to 68.5)
2013				2013	5,413	(3,887 to 7,149)	58	(39.4 to 79.1)
Female 1990 17,504 (8,671 to 26,428) 632.1 (304.7 to 941.4)	DALYs	Ischemic Stroke	Male	1990	16,798	(6,200 to 27,371)	708.3	(242.8 to 1149.1)
2013 36,595 (20,343 to 54,768) 729.2 (394.4 to 1091.8) Both 1990 34,303 (19,835 to 47,025) 669 (382.1 to 919.9) 2013 81,406 107,215 876.2 (565.0 to 1163.2) Hemorrhagic Stroke Male 1990 33,262 (20,380 to 45,244) 1095.1 (607.5 to 1563.8) 2013 75,542 102,974 1440.3 (996.7 to 2029.8) Female 1990 32,766 (21,641 to 44,631) 922 (582.3 to 1293.8) 2013 55,359 (46,656 to 83,186) 1001.8 (684.5 to 1301.4) 2013 130,901 164,073 1147 (852.3 to 1484.1) YLDS Ischemic Stroke Male 1990 286 (187 to 412) 9.9 (6.4 to 14.0) Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)				2013	44,810	(24,689 to 63,151)	1055.5	(576.9 to 1496.5)
Both 1990 34,303 (19,835 to 47,025) 669 (382.1 to 919.9)			Female	1990	17,504	(8,671 to 26,428)	632.1	(304.7 to 941.4)
Hemorrhagic Stroke Male 1990 33,262 (20,380 to 45,244) 1095.1 (607.5 to 1563.8)				2013	36,595	(20,343 to 54,768)	729.2	(394.4 to 1091.8)
Hemorrhagic Stroke Male 1990 33,262 (20,380 to 45,244) 1095.1 (607.5 to 1563.8)			Both	1990	34,303	(19,835 to 47,025)	669	(382.1 to 919.9)
Hemorrhagic Stroke Male 1990 33,262 (20,380 to 45,244) 1095.1 (607.5 to 1563.8)						(53,082 to		
Stroke Male 1990 33,262 (20,380 to 45,244) 1095.1 (607.5 to 1563.8) 2013 75,542 102,974) 1440.3 (996.7 to 2029.8) Female 1990 32,766 (21,641 to 44,631) 922 (582.3 to 1293.8) 2013 55,359 (38,639 to 75,887) 905.6 (608.6 to 1276.6) 8oth 1990 66,029 (46,656 to 83,186) 1001.8 (684.5 to 1301.4) 101,085 to (101,085 to (101,085 to (101,085 to (101,085 to (101,085 to (6.4 to 14.0) 101,085 to 2013 130,901 164,073) 1147 (852.3 to 1484.1) 101,085 to 2013 581 (363 to 866) 10.7 (6.4 to 14.0) 101,085 to 2013 581 (363 to 866) 10.7 (6.7 to 15.9) 101,085 to 2013 581 (363 to 866) 10.7 (6.7 to 15.9) 101,085 to 2013 581 (363 to 866) 10.7 (6.7 to 15.9) 101,085 to 2013 510				2013	81,406	107,215)	876.2	(565.0 to 1163.2)
2013 75,542 102,974 1440.3 (996.7 to 2029.8)								
2013 75,542 102,974 1440.3 (996.7 to 2029.8)		Stroke	Male	1990	33,262	, , , , ,	1095.1	(607.5 to 1563.8)
Female 1990 32,766 (21,641 to 44,631) 922 (582.3 to 1293.8) 2013 55,359 (38,639 to 75,887) 905.6 (608.6 to 1276.6) Both 1990 66,029 (46,656 to 83,186) 1001.8 (684.5 to 1301.4) 2013 130,901 164,073) 1147 (852.3 to 1484.1) YLDS Ischemic Stroke Male 1990 286 (187 to 412) 9.9 (6.4 to 14.0) 2013 581 (363 to 866) 10.7 (6.7 to 15.9) Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)				2042	75 5 40	1 ' '	4.440.0	(0067, 00000)
2013 55,359 (38,639 to 75,887) 905.6 (608.6 to 1276.6)			<u> </u>		-	. ,		
Both 1990 66,029 (46,656 to 83,186) 1001.8 (684.5 to 1301.4) 2013 130,901 164,073) 1147 (852.3 to 1484.1) YLDs Ischemic Stroke Male 1990 286 (187 to 412) 9.9 (6.4 to 14.0) 2013 581 (363 to 866) 10.7 (6.7 to 15.9) Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)			Female			, , ,		, ,
2013 130,901 164,073) 1147 (852.3 to 1484.1)					-			
YLDs Ischemic Stroke Male 1990 286 (187 to 412) 9.9 (6.4 to 14.0) 1 2013 581 (363 to 866) 10.7 (6.7 to 15.9) 1 Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2 2013 510 (309 to 760) 7.6 (4.6 to 11.3) 3 Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) 4 2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) 4 Hemorrhagic Male 1990 102 (64 to 149) 3 (1.8 to 4.3)			Both	1990	66,029	, , , , ,	1001.8	(684.5 to 1301.4)
YLDs Ischemic Stroke Male 1990 286 (187 to 412) 9.9 (6.4 to 14.0) 2013 581 (363 to 866) 10.7 (6.7 to 15.9) Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) 2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)						• •		
2013 581 (363 to 866) 10.7 (6.7 to 15.9)					· ·	. ,		
Female 1990 256 (156 to 377) 7.1 (4.2 to 10.4) 2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) 2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)	YLDs	Ischemic Stroke	Male					` '
2013 510 (309 to 760) 7.6 (4.6 to 11.3) Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) 2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)				2013	581	(363 to 866)	10.7	(6.7 to 15.9)
Both 1990 542 (361 to 763) 8.3 (5.6 to 11.6) 2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)			Female	1990	256	(156 to 377)	7.1	(4.2 to 10.4)
2013 1,092 (717 to 1,545) 9 (5.9 to 12.7) Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)				2013	510	(309 to 760)	7.6	(4.6 to 11.3)
Hemorrhagic Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)			Both	1990	542	(361 to 763)	8.3	(5.6 to 11.6)
Stroke Male 1990 102 (64 to 149) 3 (1.8 to 4.3)				2013	1,092	(717 to 1,545)	9	(5.9 to 12.7)
		Hemorrhagic						
2013 202 (126 to 211) 2 (1.9 to 4.6)		Stroke	Male	1990	102	(64 to 149)	3	(1.8 to 4.3)
2013 203 (120 t0 311) 3 (1.8 t0 4.0)				2013	203	(126 to 311)	3	(1.8 to 4.6)

			Female	1990	80	(51 to 124)	2	(1.2 to 3.0)
				2013	151	(91 to 225)	2	(1.2 to 3.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	182	(118 to 260)	2.4	(1.6 to 3.4)
			Both	2013	354	(225 to 511)	2.4	(1.6 to 3.5)
Myanmar	Deaths	Ischemic Stroke	Male	1990	8,657	(4,555 to 13,670)	125.1	(68.6 to 185.0)
				2013	13,332	(5,690 to 21,192)	117	(51.4 to 175.3)
			Female	1990	10,074	(5,094 to 17,164)	116.8	(60.2 to 192.9)
				2013	14,600	(4,687 to 22,481)	100.7	(32.1 to 152.7)
			Both	1990	18,731	(10,463 to 28,579)	121	(69.2 to 177.2)
				2013	27,933	(12,648 to 41,425)	108.1	(49.5 to 155.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	11,161	(6,397 to 17,958)	123.8	(71.4 to 195.6)
				2013	14,548	(7,656 to 25,744)	96.5	(51.0 to 176.3)
			Female	1990	11,585	(6,348 to 19,518)	115.2	(65.4 to 191.4)
				2013	13,985	(7,069 to 25,704)	81.6	(41.2 to 153.2)
			Both	1990	22,746	(13,928 to 35,408)	120	(75.7 to 181.2)
				2013	28,533	(15,667 to 48,075)	88.7	(50.6 to 155.3)
						(90,082 to		
	DALYs	Ischemic Stroke	Male	1990	174,875	291,835)	1970.7	(1046.0 to 3112.7)
						(98,413 to		/=
				2013	244,522	417,295)	1718	(744.6 to 2738.4)
			Female	1990	167,062	(77,633 to 302,914)	1640.5	(814.6 to 2856.8)
			remale	1330	107,002	(68,905 to	1040.3	(014.0 (0 2030.0)
				2013	214,316	347,862)	1287.3	(419.0 to 2003.9)
						(184,266 to		,
			Both	1990	341,938	543,429)	1798.1	(1000.2 to 2759.5)

2013 458,838 731,173) 1482.3 (682.5 to 2241.3) Hemorrhagic (185,612 to Stroke Male 1990 317,092 494,966) 2728.3 (1587.0 to 4335.8) (203,176 to 670,039) 2038.5 (1086.3 to 3512.7) (135,903 to Female 1990 258,287 439,664) 2158.9 (1158.6 to 3629.8)
Hemorrhagic Stroke Male 1990 317,092 494,966) 2728.3 (1587.0 to 4335.8) (203,176 to 2013 388,127 670,039) 2038.5 (1086.3 to 3512.7) (135,903 to
Stroke Male 1990 317,092 494,966) 2728.3 (1587.0 to 4335.8) (203,176 to (203,176 to <t< td=""></t<>
2013 388,127 (203,176 to 670,039) 2038.5 (1086.3 to 3512.7) (135,903 to
2013 388,127 670,039) 2038.5 (1086.3 to 3512.7) (135,903 to
(135,903 to
2013 298,066 (149,530 to (727.7 to 2626.1)
2013 298,066 526,591) 1446.9 (727.7 to 2626.1) (347,073 to
Both 1990 575,380 903,983) 2439.2 (1483.7 to 3772.3)
(370,227 to
2013 686,193 1,120,966) 1725 (948.3 to 2839.9)
YLDs Ischemic Stroke Male 1990 2,335 (1,446 to 3,373) 25.9 (16.4 to 37.2)
2013 4,392 (2,742 to 6,355) 29.1 (18.9 to 41.4)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Per 100,000
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age- Standardized
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Age-Age-Standardized Age-Age-Age-Age-Age-Age-Age-Age-Age-Age-
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Stroke Type Sex Year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Type Sex Year Mean 95% UI Mean 95% UI
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Age-Age-Age-Stroke Type Sex Year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Male 1990 2,887 (1,836 to 4,138) 28.5 (18.2 to 40.2)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Stroke Type Sex Year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Type Sex Year Mean 95% UI Mean 95% UI
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Age-Age-Age-Stroke Type Sex Year Mean 95% UI Mean 95% UI Hemorrhagic Stroke Male 1990 2,887 (1,836 to 4,138) 28.5 (18.2 to 40.2)
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Age-Age-Age-Age-Age-Age-Age-Age-Age-
Female 1990 2,384 (1,479 to 3,505) 21.7 (13.7 to 31.7) 2013 4,388 (2,887 to 6,296) 23.4 (15.3 to 33.3) Both 1990 4,720 (3,181 to 6,508) 23.5 (15.8 to 32.4) 2013 8,781 (5,927 to 12,030) 25.9 (17.6 to 35.2) Counts, All Ages Rate Age-Age-Age-Age-Age-Age-Age-Age-Age-Age-

Namibia	Deaths	Ischemic Stroke	Male	1990	129	(82 to 171)	68.4	(43.7 to 90.9)
				2013	275	(194 to 349)	83.5	(59.5 to 105.0)
			Female	1990	187	(127 to 265)	76.5	(52.7 to 108.9)
				2013	350	(221 to 460)	79.2	(51.0 to 104.5)
			Both	1990	316	(236 to 411)	73.6	(55.3 to 96.0)
			2013	625	(454 to 753)	81.5	(59.2 to 98.7)	
		Hemorrhagic						
	Deaths	Stroke	Male	1990	192	(140 to 246)	86.4	(61.1 to 110.7)
				2013	322	(240 to 439)	78.1	(58.2 to 107.9)
			Female	1990	248	(174 to 302)	90.1	(62.4 to 110.8)
				2013	384	(292 to 558)	74.8	(57.0 to 109.9)
			Both	1990	441	(333 to 522)	88.9	(65.7 to 106.0)
				2013	706	(573 to 927)	76.7	(61.8 to 103.3)
	DALYs	Ischemic Stroke	Male	1990	2,614	(1,625 to 3,513)	1106.5	(706.3 to 1469.0)
				2013	5,613	(3,882 to 7,253)	1365.6	(965.3 to 1729.1)
			Female	1990	3,253	(2,133 to 4,715)	1125.8	(763.3 to 1599.1)
				2013	5,685	(3,506 to 7,508)	1101.2	(686.7 to 1447.0)
			Both	1990	5,867	(4,319 to 7,729)	1123.7	(838.7 to 1456.4)
				2013	11,299	(8,105 to 13,824)	1218.5	(885.9 to 1474.3)
		Hemorrhagic			-			
		Stroke	Male	1990	4,918	(3,645 to 6,210)	1679.8	(1222.0 to 2140.3)
				2013	8,913	(6,560 to 12,020)	1629.9	(1209.7 to 2228.2)
			Female	1990	5,497	(3,887 to 6,716)	1631.4	(1154.9 to 1979.6)
				2013	8,687	(6,508 to 12,656)	1354.2	(1018.3 to 1981.3)
			Both	1990	10,415	(8,018 to 12,181)	1658	(1266.3 to 1951.1)
				2013	17,600	(14,152 to 22,771)	1481.5	(1199.2 to 1944.4)
	YLDs	Ischemic Stroke	Male	1990	56	(35 to 80)	19.4	(12.4 to 27.3)
				2013	98	(63 to 143)	19	(12.2 to 27.5)
			Female	1990	62	(39 to 88)	17.9	(11.4 to 25.7)

				2013	115	(73 to 169)	17.3	(10.9 to 25.7)
			Both	1990	119	(79 to 162)	18.6	(12.6 to 25.6)
				2013	214	(142 to 299)	18	(12.0 to 25.4)
		Hemorrhagic						
		Stroke	Male	1990	17	(10 to 25)	5.1	(3.2 to 7.5)
				2013	31	(18 to 45)	5.1	(3.1 to 7.5)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	17	(10 to 25)	4.3	(2.7 to 6.3)
				2013	31	(19 to 46)	4.1	(2.6 to 6.3)
			Both	1990	34	(23 to 48)	4.7	(3.1 to 6.7)
			Both	2013	62	(40 to 86)	4.5	(3.0 to 6.3)
Nepal	Deaths	Ischemic Stroke	Male	1990	1,421	(627 to 2,523)	54.7	(24.7 to 90.3)
				2013	3,873	(1,930 to 6,012)	69.5	(35.4 to 104.2)
			Female	1990	1,336	(662 to 2,687)	46.5	(24.0 to 87.5)
				2013	4,179	(2,129 to 6,045)	67.8	(35.2 to 96.3)
			Both	1990	2,757	(1,660 to 4,142)	50.3	(30.8 to 72.3)
				2013	8,052	(5,139 to 10,650)	68.7	(45.6 to 88.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,741	(843 to 2,784)	49.8	(20.4 to 84.9)
				2013	3,901	(2,261 to 6,025)	57.7	(32.5 to 93.2)
			Female	1990	1,602	(999 to 2,780)	43.2	(24.7 to 76.2)
				2013	4,143	(2,131 to 6,277)	57.1	(28.6 to 87.6)
			Both	1990	3,344	(2,156 to 4,759)	46.3	(27.2 to 68.5)
				2013	8,044	(5,067 to 11,047)	57.3	(35.8 to 79.9)
	DALYs	Ischemic Stroke	Male	1990	29,325	(13,620 to 54,521)	826.7	(383.4 to 1458.7)
						(32,292 to		
				2013	66,566	107,443)	953.2	(478.5 to 1489.3)

			Female	1990	25,098	(12,692 to 52,373)	670.6	(339.9 to 1338.4)
				2013	64,927	(32,516 to 99,587)	880.4	(456.3 to 1295.9)
			Both	1990	54,424	(31,816 to 83,252)	744.6	(452.3 to 1107.8)
						(83,523 to		
				2013	131,493	181,364)	915.9	(581.3 to 1217.4)
		Hemorrhagic						
		Stroke	Male	1990	49,980	(29,223 to 75,701)	1030.7	(512.9 to 1633.7)
						(56,181 to		
				2013	91,651	135,001)	1077.2	(647.0 to 1607.3)
			Female	1990	40,930	(27,048 to 69,323)	843.1	(537.5 to 1447.6)
						(46,885 to		
				2013	87,975	132,704)	1010.2	(534.9 to 1520.3)
			5	1000	00.044	(63,882 to	000.0	(64.6.2)
			Both	1990	90,911	123,466)	933.8	(616.3 to 1318.2)
				2013	170 627	(116,731 to	1040.1	(662.0 to 1405.0)
	\/I 5		24.1		179,627	241,964)		(662.9 to 1405.0)
	YLDs	Ischemic Stroke	Male	1990	1,220	(740 to 1,811)	27.5	(17.0 to 40.5)
				2013	2,513	(1,661 to 3,544)	28.8	(19.2 to 41.1)
			Female	1990	1,168	(729 to 1,657)	24.4	(15.0 to 34.5)
				2013	2,217	(1,424 to 3,141)	23.9	(15.3 to 33.9)
			Both	1990	2,389	(1,550 to 3,344)	25.8	(16.9 to 36.0)
				2013	4,731	(3,219 to 6,399)	26.2	(17.8 to 35.6)
		Hemorrhagic						
		Stroke	Male	1990	690	(445 to 1,001)	14.1	(9.1 to 20.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,327	(826 to 1,881)	14.5	(9.1 to 20.5)
			Female	1990	573	(351 to 832)	11	(6.8 to 15.7)
				2013	1,055	(668 to 1,502)	10.6	(6.7 to 15.3)

			Both	1990	1,264	(826 to 1,754)	12.5	(8.1 to 17.1)
			Both	2013	2,382	(1,596 to 3,285)	12.5	(8.3 to 17.2)
Netherlands	Deaths	Ischemic Stroke	Male	1990	3,028	(2,424 to 3,793)	44.3	(35.4 to 55.3)
				2013	2,462	(1,764 to 3,151)	20.8	(15.0 to 26.8)
			Female	1990	4,623	(3,481 to 5,375)	36.3	(27.3 to 42.0)
				2013	4,273	(3,204 to 6,078)	22.4	(17.0 to 32.3)
			Both	1990	7,652	(6,457 to 8,726)	39.6	(33.5 to 45.2)
				2013	6,735	(5,365 to 8,383)	22.1	(17.7 to 27.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,138	(1,684 to 2,714)	29.7	(23.3 to 37.9)
				2013	2,083	(1,407 to 2,707)	17.5	(11.8 to 22.9)
			Female	1990	2,918	(2,128 to 3,666)	24.9	(18.6 to 31.2)
				2013	3,064	(2,117 to 4,471)	17.7	(12.4 to 25.1)
			Both	1990	5,057	(4,135 to 6,017)	27	(22.2 to 32.0)
				2013	5,148	(3,831 to 6,678)	17.7	(13.4 to 22.7)
	DALYs	Ischemic Stroke	Male	1990	44,196	(36,548 to 54,821)	606.7	(501.4 to 749.6)
				2013	34,861	(26,870 to 42,879)	288.4	(222.0 to 354.6)
			Female	1990	48,929	(38,491 to 56,427)	417.3	(328.0 to 484.6)
				2013	41,695	(32,713 to 57,339)	250.8	(198.6 to 342.5)
						(81,107 to		
			Both	1990	93,125	105,610)	496.8	(434.2 to 566.3)
				2013	76,557	(63,906 to 92,325)	269.9	(225.6 to 324.8)
		Hemorrhagic				(2.1.2.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1		(
		Stroke	Male	1990	39,206	(31,594 to 48,534)	513.4	(412.1 to 638.3)
				2013	31,939	(23,261 to 39,965)	268.9	(194.9 to 336.6)
			Female	1990	41,272	(32,262 to 50,950)	407.6	(322.4 to 508.1)
				2013	37,385	(27,489 to 51,295)	263.5	(197.6 to 352.2)
			Both	1990	80,478	(69,056 to 93,054)	452.4	(388.2 to 521.9)
				2013	69,324	(55,479 to 84,566)	265.1	(213.3 to 320.9)

	YLDs	Ischemic Stroke	Male	1990	4,881	(3,256 to 6,774)	64.4	(43.2 to 89.3)
				2013	6,996	(4,416 to 10,056)	57.4	(36.5 to 82.7)
			Female	1990	5,165	(3,405 to 7,437)	48.8	(32.5 to 70.3)
				2013	6,579	(4,307 to 9,245)	44.9	(29.8 to 62.4)
			Both	1990	10,046	(6,715 to 13,727)	55.4	(37.0 to 75.5)
				2013	13,575	(8,981 to 18,652)	50.7	(33.5 to 69.8)
		Hemorrhagic						
		Stroke	Male	1990	1,123	(698 to 1,732)	14.6	(9.1 to 22.5)
				2013	1,794	(1,035 to 2,723)	15.3	(9.0 to 22.8)
			Female	1990	1,221	(662 to 1,873)	12.3	(6.9 to 18.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,525	(956 to 2,291)	11.5	(7.3 to 17.1)
			Both	1990	2,345	(1,460 to 3,367)	13.2	(8.3 to 19.0)
			Both	2013	3,320	(2,119 to 4,695)	13.1	(8.4 to 18.6)
New Zealand	Deaths	Ischemic Stroke	Male	1990	700	(568 to 874)	52	(42.4 to 65.0)
				2013	748	(544 to 956)	25.2	(18.3 to 32.1)
			Female	1990	1,100	(860 to 1,333)	49	(38.3 to 59.4)
				2013	1,218	(881 to 1,495)	29.4	(21.2 to 35.8)
			Both	1990	1,800	(1,541 to 2,064)	50.8	(43.4 to 58.4)
				2013	1,966	(1,544 to 2,321)	27.8	(21.8 to 32.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	366	(290 to 474)	24.3	(19.1 to 31.8)
				2013	383	(261 to 498)	13.3	(9.0 to 17.2)
			Female	1990	509	(412 to 676)	25.2	(20.5 to 33.0)
				2013	550	(372 to 731)	15.1	(10.6 to 19.6)
			Both	1990	876	(766 to 1,085)	24.9	(21.7 to 30.7)
				2013	933	(713 to 1,128)	14.3	(11.1 to 17.0)

	DALYs	Ischemic Stroke	Male	1990	9,658	(8,041 to 11,823)	659.1	(547.5 to 810.1)
				2013	8,123	(6,133 to 10,071)	283.4	(213.3 to 350.2)
			Female	1990	11,058	(8,911 to 13,280)	505.4	(406.6 to 605.5)
				2013	9,637	(7,137 to 11,510)	257.2	(192.2 to 300.7)
			Both	1990	20,717	(18,324 to 23,800)	575.6	(508.8 to 662.6)
				2013	17,760	(14,434 to 20,466)	271.2	(220.6 to 312.2)
		Hemorrhagic						
		Stroke	Male	1990	8,233	(6,705 to 10,393)	511.6	(417.6 to 651.0)
				2013	6,927	(5,176 to 8,640)	249.9	(186.7 to 310.7)
			Female	1990	9,577	(8,078 to 12,178)	511.1	(430.6 to 645.3)
				2013	8,045	(6,135 to 10,096)	256.3	(197.8 to 318.7)
			Both	1990	17,810	(15,858 to 21,229)	510.1	(453.2 to 608.5)
				2013	14,972	(12,470 to 17,253)	253.4	(211.0 to 292.4)
	YLDs	Ischemic Stroke	Male	1990	717	(482 to 1,002)	46.3	(31.4 to 64.5)
				2013	897	(577 to 1,276)	32	(20.4 to 45.5)
			Female	1990	655	(443 to 890)	32.5	(22.1 to 44.2)
				2013	747	(481 to 1,051)	23.4	(15.0 to 32.8)
			Both	1990	1,373	(946 to 1,847)	38.7	(26.8 to 52.0)
				2013	1,644	(1,117 to 2,261)	27.4	(18.6 to 37.6)
		Hemorrhagic						
		Stroke	Male	1990	535	(360 to 756)	34	(22.9 to 47.8)
				2013	748	(477 to 1,083)	27.3	(17.7 to 39.3)
			Female	1990	432	(290 to 595)	22.2	(14.9 to 30.6)
				2013	508	(334 to 716)	16.7	(10.9 to 23.5)
			Both	1990	967	(669 to 1,311)	27.4	(18.9 to 37.3)
			Both	2013	1,256	(836 to 1,711)	21.7	(14.4 to 29.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

Nicaragua	Deaths	Ischemic Stroke	Male	1990	295	(154 to 390)	56.7	(29.3 to 75.7)
				2013	248	(173 to 436)	18.4	(12.8 to 32.2)
			Female	1990	383	(185 to 470)	58.4	(28.3 to 72.2)
				2013	249	(180 to 431)	14.9	(10.8 to 25.8)
			Both	1990	678	(373 to 836)	57.9	(31.3 to 71.6)
				2013	498	(384 to 813)	16.5	(12.7 to 26.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	246	(182 to 386)	40.2	(27.8 to 66.4)
				2013	762	(550 to 941)	54.5	(39.0 to 67.8)
			Female	1990	304	(253 to 463)	40.2	(32.4 to 64.2)
				2013	788	(607 to 988)	46.1	(35.4 to 57.8)
			Both	1990	551	(452 to 805)	40.2	(31.7 to 61.7)
				2013	1,551	(1,259 to 1,771)	50	(40.3 to 57.2)
	DALYs	Ischemic Stroke	Male	1990	6,060	(3,192 to 7,897)	838.1	(442.8 to 1101.8)
				2013	3,888	(2,809 to 6,601)	255.5	(183.9 to 437.0)
			Female	1990	6,794	(3,235 to 8,329)	809.3	(389.3 to 991.8)
				2013	3,195	(2,357 to 5,432)	181.7	(133.9 to 310.2)
			Both	1990	12,854	(7,075 to 15,854)	823.9	(457.3 to 1015.2)
				2013	7,083	(5,559 to 11,344)	216.3	(168.6 to 348.9)
		Hemorrhagic						
		Stroke	Male	1990	6,715	(5,208 to 9,818)	733.1	(545.8 to 1132.3)
				2013	13,871	(10,461 to 16,794)	848.4	(636.0 to 1029.3)
			Female	1990	7,657	(6,369 to 10,557)	749.2	(629.3 to 1080.0)
				2013	12,305	(9,844 to 15,627)	659.3	(527.3 to 835.1)
			Both	1990	14,373	(12,315 to 19,275)	739.4	(616.3 to 1041.4)
				2013	26,177	(22,075 to 29,969)	748.2	(624.1 to 856.5)
	YLDs	Ischemic Stroke	Male	1990	39	(25 to 55)	5.1	(3.3 to 7.2)
				2013	100	(62 to 146)	6.3	(4.0 to 9.1)
			Female	1990	43	(26 to 62)	5.1	(3.1 to 7.4)

				2013	101	(63 to 148)	5.7	(3.5 to 8.4)
			Both	1990	82	(54 to 114)	5.1	(3.4 to 7.1)
				2013	202	(137 to 281)	6	(4.1 to 8.4)
		Hemorrhagic						
		Stroke	Male	1990	122	(84 to 169)	13.2	(9.2 to 18.9)
				2013	328	(217 to 464)	18.3	(12.4 to 26.0)
			Female	1990	110	(69 to 155)	11.2	(7.0 to 15.9)
				2013	251	(163 to 364)	12.6	(8.2 to 18.4)
			Both	1990	232	(160 to 311)	12.2	(8.5 to 16.5)
			Both	2013	580	(399 to 811)	15.3	(10.5 to 21.2)
Niger	Deaths	Ischemic Stroke	Male	1990	431	(202 to 710)	51.2	(26.8 to 81.3)
				2013	944	(114 to 1,603)	46.9	(6.5 to 78.1)
			Female	1990	590	(266 to 993)	61.7	(30.8 to 98.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,417	(99 to 2,319)	68.6	(5.2 to 108.7)
			Both	1990	1,022	(639 to 1,376)	57.2	(35.7 to 76.0)
				2013	2,361	(469 to 3,619)	58.6	(12.4 to 88.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	628	(400 to 893)	47.8	(24.5 to 72.9)
	Deaths		Male	1990 2013	628 901	(400 to 893) (461 to 1,760)	47.8 32.9	(24.5 to 72.9) (14.4 to 74.5)
	Deaths		Male Female			· , , , , , , , , , , , , , , , , , , ,		, ,
	Deaths			2013	901	(461 to 1,760)	32.9	(14.4 to 74.5)
	Deaths			2013 1990	901 728	(461 to 1,760) (358 to 1,027)	32.9 52.3	(14.4 to 74.5) (23.4 to 80.1)
	Deaths		Female	2013 1990 2013	901 728 1,167	(461 to 1,760) (358 to 1,027) (372 to 2,259)	32.9 52.3 42.4	(14.4 to 74.5) (23.4 to 80.1) (13.1 to 89.9)
	Deaths		Female	2013 1990 2013 1990	901 728 1,167 1,356	(461 to 1,760) (358 to 1,027) (372 to 2,259) (881 to 1,744)	32.9 52.3 42.4 50.3	(14.4 to 74.5) (23.4 to 80.1) (13.1 to 89.9) (30.0 to 66.6)
		Stroke	Female Both	2013 1990 2013 1990 2013	901 728 1,167 1,356 2,068	(461 to 1,760) (358 to 1,027) (372 to 2,259) (881 to 1,744) (1,176 to 3,874)	32.9 52.3 42.4 50.3 37.8	(14.4 to 74.5) (23.4 to 80.1) (13.1 to 89.9) (30.0 to 66.6) (19.2 to 79.8)

			2013	27,204	(1,995 to 47,022)	986.2	(78.7 to 1616.9)
		Both	1990	26,310	(16,693 to 36,949)	915.6	(583.3 to 1221.6)
			2013	47,651	(9,760 to 74,847)	860.8	(182.6 to 1313.4)
	Hemorrhagic						
	Stroke	Male	1990	25,534	(16,658 to 36,643)	1066.7	(644.8 to 1538.2)
		2013	27,618	(16,100 to 46,684)	649.8	(328.6 to 1281.5)	
	Female	1990	26,938	(14,931 to 37,687)	1107.9	(535.3 to 1591.7)	
		2013	33,160	(10,677 to 59,077)	829.2	(265.9 to 1580.9)	
	Both	1990	52,472	(35,680 to 68,606)	1090.4	(690.9 to 1398.4)	
					(36,723 to		
			2013	60,778	101,771)	738.7	(419.1 to 1380.8)
YLDs	Ischemic Stroke	Male	1990	231	(150 to 334)	15.4	(10.1 to 22.8)
			2013	566	(356 to 805)	15.2	(9.7 to 21.4)
		Female	1990	194	(122 to 279)	11.5	(7.4 to 16.6)
			2013	429	(277 to 609)	11.8	(7.8 to 16.7)
		Both	1990	425	(281 to 593)	13.3	(9.0 to 18.6)
			2013	996	(661 to 1,354)	13.5	(8.8 to 18.2)
	Hemorrhagic						
	Stroke	Male	1990	76	(46 to 114)	4.2	(2.6 to 6.3)
			2013	173	(104 to 252)	3.9	(2.3 to 5.9)
		Female	1990	57	(36 to 83)	2.9	(1.9 to 4.2)
			2013	117	(70 to 170)	2.8	(1.7 to 4.0)
		Both	1990	134	(86 to 187)	3.5	(2.3 to 4.9)
		Both	2013	290	(182 to 411)	3.3	(2.1 to 4.7)
Deaths	Ischemic Stroke	Male	1990	6,759	(4,530 to 9,934)	62.5	(43.4 to 88.7)
			2013	15,729	(9,608 to 21,473)	84.7	(55.5 to 110.2)
		Female	1990	10,135	(7,700 to 14,202)	81	(61.4 to 108.8)
			2013	19,204	(12,150 to 25,482)	99.9	(63.0 to 128.4)
		Both	1990	16,895	(13,643 to 21,455)	72.9	(59.2 to 90.4)
		YLDs Ischemic Stroke Hemorrhagic Stroke	Hemorrhagic Stroke Female Both YLDs Ischemic Stroke Male Female Both Hemorrhagic Stroke Male Female Both Hemorrhagic Stroke Male Female Female Female Female Female Female Female Female	Both 1990 2013 Hemorrhagic Stroke Male 1990 2013 Female 1990 2013 Both 1990 2013 Female 1990 2013 Female 1990 2013 Female 1990 2013 Both 1990 2013 Hemorrhagic Stroke Male 1990 2013 Female 1990 2013 Female 1990 2013 Both 1990 2013 Deaths Ischemic Stroke Male 1990 2013 Female 1990 2013 Emale 1990 2013 Female 1990 2013	Both 1990 26,310 2013 47,651 47,651 47,651	Both 1990 26,310 (16,693 to 36,949) 2013 47,651 (9,760 to 74,847)	Both 1990 26,310 (16,693 to 36,949) 915.6

					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	34,933	(23,101 to 43,286)	92.8	(64.5 to 112.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	6,559	(3,331 to 8,821)	43.3	(19.2 to 59.9)
				2013	9,537	(5,936 to 16,626)	35.2	(20.6 to 66.5)
			Female	1990	8,622	(5,625 to 11,446)	52.9	(30.4 to 71.1)
				2013	10,944	(7,132 to 19,738)	39.5	(24.8 to 75.2)
			Both	1990	15,181	(10,485 to 18,527)	48.6	(30.8 to 60.6)
				2013	20,481	(14,789 to 34,536)	37.5	(26.0 to 66.3)
					,	(101,533 to		,
	DALYs	Ischemic Stroke	Male	1990	151,415	232,063)	922.1	(625.4 to 1344.5)
						(190,074 to		
				2013	323,230	454,666)	1205.6	(749.5 to 1629.1)
						(154,803 to		
			Female	1990	207,793	306,585)	1166.6	(886.1 to 1627.8)
						(227,975 to		()
				2013	362,406	488,626)	1339.3	(848.3 to 1768.5)
			D. H	4000	250 200	(286,745 to	4056.7	(055.21, 4224.5)
			Both	1990	359,208	474,946)	1056.7	(855.3 to 1331.5)
				2013	685,636	(447,557 to 854,507)	1277.6	(851.0 to 1570.8)
		Hemorrhagic		2013	065,030	(127,825 to	12//.0	(001.0 to 1070.0)
		Stroke	Male	1990	214,271	279,870)	831.7	(413.8 to 1112.7)
		- Grone	maic	1330		(199,534 to	001.7	(110.0 to 1112.7)
				2013	306,570	477,853)	687.3	(428.4 to 1214.2)
						(173,213 to		,
			Female	1990	248,195	348,402)	985.5	(639.1 to 1303.3)
						(218,944 to		
				2013	324,134	566,523)	751.5	(491.9 to 1336.8)

						(331,654 to		
			Both	1990	462,466	580,327)	914.7	(625.2 to 1122.1)
				2012	620.704	(470,371 to	720.7	(F20 0 + - 4242 0)
				2013	630,704	991,236)	720.7	(520.0 to 1213.8)
	YLDs	Ischemic Stroke	Male	1990	3,218	(2,065 to 4,644)	15.3	(9.8 to 21.8)
				2013	5,609	(3,339 to 8,466)	14.6	(8.8 to 22.4)
			Female	1990	2,570	(1,656 to 3,643)	11.4	(7.4 to 15.9)
				2013	4,626	(2,979 to 6,755)	11.9	(7.8 to 17.5)
			Both	1990	5,789	(3,975 to 7,959)	13.2	(9.2 to 18.0)
				2013	10,236	(6,479 to 14,740)	13.2	(8.5 to 19.3)
		Hemorrhagic Stroke	Male	1990	1,039	(625 to 1,554)	4.1	(2.5 to 6.2)
		Stroke	Widie	2013	1,819	(1,121 to 2,680)	3.9	(2.3 to 5.9)
			Female	1990	769	(467 to 1,133)	2.9	(1.8 to 4.3)
			remare	2013	1,263	(767 to 1,883)	2.7	(1.7 to 4.1)
			Both	1990	1,809	(1,203 to 2,533)	3.5	(2.3 to 4.9)
			Both	2013	3,083	(1,991 to 4,448)	3.3	(2.1 to 4.8)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
North Korea	Deaths	Ischemic Stroke	Male	1990	1,544	(698 to 2,382)	59.8	(27.6 to 90.1)
				2013	4,872	(2,516 to 7,389)	76.5	(41.6 to 109.4)
			Female	1990	2,818	(1,328 to 4,424)	50.2	(24.6 to 77.7)
				2013	7,807	(4,058 to 10,223)	60.5	(31.2 to 79.8)
			Both	1990	4,363	(2,350 to 6,188)	53.1	(29.1 to 77.3)
				2013	12,679	(8,108 to 16,483)	66.2	(42.3 to 84.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	6,424	(4,240 to 9,287)	181.3	(127.9 to 251.3)
				2013	14,370	(10,487 to 19,512)	182	(134.8 to 243.0)
			Female	1990	10,116	(6,854 to 14,777)	160.5	(111.5 to 231.0)

2013 18,025 (13,738 to 24,128) 135.6 Both 1990 16,541 (12,344 to 21,771) 168.2 2013 32,395 (26,754 to 41,223) 153.7 DALYS Ischemic Stroke Male 1990 36,571 (16,064 to 58,528) 930.7 (52,772 to 159,657) 1171.5 Female 1990 49,461 (21,953 to 80,721) 734.2	(104.9 to 181.3) (127.5 to 220.5) (127.1 to 196.2) (445.3 to 1416.4) (633.5 to 1753.1) (344.1 to 1159.5)
2013 32,395 (26,754 to 41,223) 153.7 DALYS Ischemic Stroke Male 1990 36,571 (16,064 to 58,528) 930.7	(127.1 to 196.2) (445.3 to 1416.4) (633.5 to 1753.1)
DALYS Ischemic Stroke Male 1990 36,571 (16,064 to 58,528) 930.7	(445.3 to 1416.4) (633.5 to 1753.1)
2013 100,875 (52,772 to 1171.5	(633.5 to 1753.1)
2013 100,875 159,657) 1171.5	· · · · · · · · · · · · · · · · · · ·
	· · · · · · · · · · · · · · · · · · ·
Female 1990 49 461 (21 052 to 90 721) 724 2	(3/// 1 to 1150 5)
	(344.1 (0 1133.3)
(60,674 to	
2013 115,892 153,236) 827	(426.3 to 1083.0)
(45,646 to	
Both 1990 86,032 125,537) 802	(445.9 to 1129.1)
(139,817 to	
2013 216,768 293,219) 962.3	(621.4 to 1270.2)
Hemorrhagic (124,688 to	
Stroke Male 1990 191,359 283,938) 3482.9	(2366.8 to 4931.7)
(267,214 to	
2013 365,442 501,237) 3504.9	(2588.8 to 4741.2)
(142,926 to	
Female 1990 221,571 334,890) 2857.4	(1897.6 to 4203.1)
(244,908 to	
2013 330,636 440,265) 2327.2	(1737.2 to 3103.6)
(304,756 to	(
Both 1990 412,930 547,128) 3114.6	(2337.2 to 4088.4)
(563,087 to	(2222.012504.2)
2013 696,078 878,669) 2835.5	(2332.0 to 3594.2)
YLDs Ischemic Stroke Male 1990 1,363 (815 to 2,110) 35.7	(21.5 to 55.8)
2013 3,097 (1,850 to 4,553) 36.2	(21.8 to 55.1)
Female 1990 1,613 (1,012 to 2,380) 23.1	(14.4 to 33.3)
2013 3,317 (2,087 to 4,896) 23.5	(14.8 to 34.1)
Both 1990 2,977 (1,939 to 4,292) 26.9	(17.3 to 38.6)
2013 6,414 (4,253 to 9,022) 27.7	(18.5 to 38.9)

		Hemorrhagic						
		Stroke	Male	1990	1,656	(1,028 to 2,464)	37.6	(22.7 to 56.7)
				2013	3,663	(2,138 to 5,555)	40.5	(23.2 to 61.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	1,692	(1,100 to 2,455)	22.2	(14.5 to 31.7)
				2013	3,112	(1,946 to 4,482)	22.1	(13.9 to 31.9)
			Both	1990	3,348	(2,200 to 4,741)	26.9	(17.5 to 38.6)
			Both	2013	6,776	(4,456 to 9,660)	28.5	(18.6 to 40.6)
Norway	Deaths	Ischemic Stroke	Male	1990	1,366	(1,081 to 1,696)	52.4	(41.4 to 65.3)
				2013	980	(720 to 1,220)	27.9	(20.6 to 34.7)
			Female	1990	2,016	(1,440 to 2,332)	45.1	(32.1 to 52.2)
				2013	1,396	(1,059 to 2,039)	23.9	(18.3 to 34.9)
			Both	1990	3,382	(2,816 to 3,840)	48.4	(40.4 to 54.9)
				2013	2,377	(1,953 to 3,060)	25.8	(21.4 to 33.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	858	(669 to 1,198)	33.2	(25.8 to 46.2)
				2013	752	(465 to 979)	21.6	(13.5 to 28.1)
			Female	1990	1,211	(909 to 1,528)	30.1	(23.1 to 37.1)
				2013	889	(678 to 1,314)	16.8	(13.2 to 24.4)
			Both	1990	2,070	(1,746 to 2,547)	31.6	(26.9 to 39.0)
				2013	1,642	(1,254 to 2,107)	19	(14.8 to 24.0)
	DALYs	Ischemic Stroke	Male	1990	19,039	(15,813 to 23,865)	716.7	(593.6 to 899.8)
				2013	12,758	(9,927 to 15,452)	372	(289.0 to 451.2)
			Female	1990	21,114	(16,016 to 24,045)	516.6	(399.4 to 592.9)
				2013	12,459	(9,947 to 17,108)	255.7	(206.0 to 352.4)
			Both	1990	40,153	(34,372 to 45,563)	605	(520.7 to 686.5)
				2013	25,218	(21,243 to 30,694)	309.4	(258.7 to 376.2)

347

	Hemorrhagic						
Deaths	Stroke	Male	1990	170	(114 to 243)	73.5	(49.1 to 101.6)
			2013	312	(217 to 421)	53.4	(37.4 to 72.7)
		Female	1990	224	(161 to 314)	77	(56.3 to 105.1)
			2013	247	(175 to 332)	48.8	(34.6 to 65.5)
		Both	1990	395	(304 to 500)	74.7	(57.9 to 93.0)
			2013	560	(437 to 701)	51.2	(40.2 to 64.4)
DALYs	Ischemic Stroke	Male	1990	1,059	(628 to 1,873)	411.2	(251.0 to 688.0)
			2013	3,639	(2,328 to 4,905)	566.8	(361.8 to 748.5)
		Female	1990	1,054	(625 to 2,012)	386.6	(236.2 to 716.3)
			2013	2,240	(1,317 to 2,994)	446.1	(261.9 to 591.6)
		Both	1990	2,113	(1,405 to 3,589)	397.6	(271.9 to 651.4)
			2013	5,880	(4,057 to 7,535)	511.7	(350.9 to 652.1)
	Hemorrhagic						
	Stroke	Male	1990	5,510	(3,597 to 7,961)	1273.4	(846.0 to 1809.8)
			2013	7,070	(4,869 to 9,772)	871.5	(594.7 to 1172.4)
		Female	1990	7,713	(4,906 to 11,249)	1530.1	(1086.9 to 2169.4)
			2013	5,258	(3,668 to 7,262)	798.8	(549.0 to 1098.2)
		Both	1990	13,223	(9,758 to 17,777)	1381.5	(1054.3 to 1752.2)
			2013	12,328	(9,541 to 15,642)	834.9	(645.9 to 1045.2)
YLDs	Ischemic Stroke	Male	1990	36	(23 to 54)	9.2	(5.9 to 13.2)
			2013	94	(62 to 138)	9.2	(6.1 to 13.5)
		Female	1990	23	(14 to 33)	7.1	(4.5 to 10.2)
			2013	48	(32 to 67)	7.3	(5.0 to 10.2)
_		Both	1990	60	(39 to 83)	8.1	(5.5 to 11.1)
			2013	143	(97 to 200)	8.4	(5.8 to 11.7)
	Hemorrhagic						
	Stroke	Male	1990	7	(3 to 11)	1.4	(0.7 to 2.2)
			2013	18	(10 to 30)	1.3	(0.7 to 2.2)

			Female	1990	6	(3 to 9)	1.6	(0.9 to 2.4)
				2013	10	(6 to 17)	1.4	(0.8 to 2.1)
			Both	1990	13	(8 to 19)	1.4	(0.9 to 2.1)
			Both	2013	29	(17 to 44)	1.3	(0.8 to 2.0)
Pakistan	Deaths	Ischemic Stroke	Male	1990	11,151	(5,581 to 16,584)	54.3	(28.1 to 80.0)
				2013	28,709	(16,587 to 39,150)	77.5	(46.0 to 105.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	13,603	(7,721 to 18,854)	81.3	(45.1 to 110.2)
				2013	30,526	(17,267 to 44,951)	89.4	(51.3 to 129.6)
			Both	1990	24,754	(16,667 to 32,015)	66.4	(44.1 to 85.2)
				2013	59,235	(41,444 to 77,651)	83.2	(58.4 to 108.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	11,457	(6,194 to 17,303)	46	(23.0 to 71.5)
				2013	28,546	(18,292 to 41,241)	64.3	(39.7 to 94.0)
			Female	1990	15,090	(8,705 to 20,702)	73.1	(40.4 to 103.7)
				2013	30,762	(17,030 to 46,336)	74.9	(39.7 to 115.0)
			Both	1990	26,548	(17,496 to 35,559)	58.3	(37.1 to 79.5)
				2013	59,309	(39,064 to 79,834)	69.5	(45.6 to 92.9)
						(98,756 to		
	DALYs	Ischemic Stroke	Male	1990	198,261	310,379)	784	(397.3 to 1182.7)
						(285,867 to		
				2013	491,635	695,166)	1111.9	(647.4 to 1538.8)
						(130,979 to		
			Female	1990	230,938	326,433)	1129.6	(653.7 to 1565.0)
				2012	405.040	(273,127 to	4400	(676 A : 4776 A
				2013	485,310	738,714)	1192	(676.4 to 1778.1)
			Both	1990	429,199	(286,948 to 556,141)	939.6	(634.3 to 1215.2)
			DULII	1990	429,139	330,141)	959.0	(034.3 (0 1213.2)

						(668,936 to		
				2013	976,946	1,311,036)	1150.4	(797.4 to 1521.4)
		Hemorrhagic				(179,615 to		
		Stroke	Male	1990	293,479	419,222)	913.2	(519.4 to 1348.6)
						(461,016 to		
				2013	712,097	1,025,732)	1277.8	(832.6 to 1831.7)
						(222,814 to		
			Female	1990	366,233	490,920)	1417.3	(824.5 to 1940.1)
						(421,802 to		
				2013	710,786	1,084,402)	1400.8	(798.9 to 2135.2)
						(459,253 to		
			Both	1990	659,712	851,372)	1142.6	(769.4 to 1504.7)
						(967,336 to		
				2013	1,422,883	1,894,784)	1337.7	(889.8 to 1789.6)
	YLDs	Ischemic Stroke	Male	1990	8,413	(5,341 to 11,847)	27.9	(18.2 to 38.9)
				2013	19,148	(12,559 to 26,984)	37	(24.5 to 52.0)
			Female	1990	5,777	(3,724 to 8,198)	22.4	(14.5 to 31.7)
				2013	10,900	(6,887 to 15,695)	20.8	(13.2 to 30.4)
			Both	1990	14,191	(9,325 to 19,445)	25.4	(16.8 to 34.5)
				2013	30,049	(20,472 to 42,089)	29.1	(19.9 to 41.1)
		Hemorrhagic						
		Stroke	Male	1990	4,690	(2,947 to 6,674)	14.5	(9.3 to 20.3)
				2013	9,976	(6,471 to 14,246)	17.9	(11.8 to 25.5)
			Female	1990	2,782	(1,707 to 4,086)	9.8	(6.0 to 14.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	5,227	(3,202 to 7,669)	9.1	(5.5 to 13.5)
			Both	1990	7,472	(4,924 to 10,434)	12.3	(8.1 to 17.1)
			Both	2013	15,203	(10,216 to 20,804)	13.6	(9.1 to 18.7)

Palestine	Deaths	Ischemic Stroke	Male	1990	56	(33 to 86)	28.6	(17.3 to 42.9)
				2013	176	(105 to 257)	32.9	(19.4 to 47.9)
			Female	1990	71	(38 to 105)	31.2	(16.7 to 45.3)
				2013	176	(101 to 251)	31.7	(18.1 to 44.4)
			Both	1990	128	(85 to 178)	30.1	(19.7 to 41.4)
				2013	352	(232 to 453)	32.6	(21.4 to 42.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	293	(207 to 393)	116.9	(83.3 to 156.2)
				2013	625	(442 to 868)	102.4	(73.6 to 139.3)
			Female	1990	397	(291 to 538)	142.7	(104.5 to 190.5)
				2013	593	(456 to 805)	94.4	(74.4 to 126.3)
			Both	1990	691	(551 to 863)	131.2	(103.6 to 162.0)
				2013	1,219	(983 to 1,520)	98.9	(81.1 to 121.0)
	DALYs	Ischemic Stroke	Male	1990	1,258	(761 to 1,961)	444	(269.9 to 668.7)
				2013	3,573	(2,235 to 5,211)	491.3	(296.5 to 713.7)
			Female	1990	1,454	(800 to 2,243)	453.9	(245.5 to 674.3)
				2013	2,777	(1,690 to 4,122)	384.1	(228.2 to 555.6)
			Both	1990	2,712	(1,929 to 3,842)	448.7	(304.8 to 625.3)
				2013	6,351	(4,384 to 8,202)	438.1	(296.2 to 560.6)
		Hemorrhagic						
		Stroke	Male	1990	9,717	(7,014 to 12,969)	2038.6	(1408.1 to 2771.6)
				2013	15,796	(11,105 to 21,718)	1702.4	(1194.7 to 2370.2)
			Female	1990	12,031	(8,408 to 17,084)	2384.7	(1701.7 to 3306.4)
				2013	12,562	(9,438 to 16,783)	1344.9	(1013.2 to 1836.4)
			Both	1990	21,748	(17,034 to 27,922)	2223.4	(1759.9 to 2799.9)
				2013	28,358	(22,651 to 35,389)	1521.7	(1216.4 to 1913.3)
	YLDs	Ischemic Stroke	Male	1990	43	(28 to 61)	11.6	(7.7 to 16.2)
				2013	114	(74 to 163)	12	(7.7 to 16.9)
			Female	1990	34	(23 to 50)	8.8	(5.9 to 12.4)

				2013	91	(58 to 128)	9.4	(6.1 to 13.2)
			Both	1990	77	(52 to 108)	10.1	(7.0 to 13.7)
				2013	205	(137 to 281)	10.6	(7.1 to 14.5)
		Hemorrhagic						
		Stroke	Male	1990	8	(4 to 13)	1.8	(1.0 to 2.7)
				2013	19	(11 to 30)	1.7	(1.0 to 2.6)
			Female	1990	8	(5 to 13)	1.8	(1.1 to 2.9)
				2013	19	(11 to 30)	1.7	(1.0 to 2.7)
			Both	1990	16	(10 to 24)	1.8	(1.1 to 2.6)
			Both	2013	38	(23 to 57)	1.7	(1.0 to 2.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Panama	Deaths	Ischemic Stroke	Male	1990	342	(223 to 435)	65.1	(42.0 to 82.8)
				2013	356	(211 to 645)	26.5	(15.7 to 47.9)
			Female	1990	340	(199 to 408)	57.2	(33.5 to 68.7)
				2013	321	(206 to 620)	20.6	(13.2 to 39.8)
			Both	1990	683	(457 to 820)	61	(41.1 to 73.3)
				2013	678	(443 to 1,188)	23.4	(15.3 to 41.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	218	(151 to 362)	38.3	(25.8 to 66.4)
				2013	769	(454 to 1,018)	55.9	(32.9 to 74.1)
			Female	1990	223	(168 to 376)	35.5	(26.1 to 61.5)
				2013	718	(421 to 901)	45.9	(26.9 to 57.6)
			Both	1990	441	(332 to 699)	36.9	(27.2 to 60.2)
				2013	1,488	(954 to 1,845)	50.8	(32.3 to 63.0)
	DALYs	Ischemic Stroke	Male	1990	5,442	(3,445 to 6,887)	882.9	(566.5 to 1115.2)
				2013	5,031	(3,153 to 8,848)	352.1	(219.5 to 620.8)
			Female	1990	4,641	(2,739 to 5,550)	705.7	(419.0 to 841.4)

				2013	3,561	(2,332 to 6,912)	227.3	(148.8 to 440.1)
			Both	1990	10,083	(6,642 to 12,107)	791.6	(528.2 to 948.8)
				2013	8,592	(5,810 to 14,844)	287.3	(194.0 to 495.5)
		Hemorrhagic						
		Stroke	Male	1990	4,590	(3,457 to 6,989)	656.6	(480.9 to 1032.2)
				2013	12,792	(8,041 to 16,984)	853.4	(532.1 to 1130.0)
			Female	1990	4,232	(3,483 to 6,397)	577.4	(465.4 to 898.2)
				2013	9,466	(6,188 to 11,708)	588.6	(382.3 to 729.1)
			Both	1990	8,822	(7,134 to 12,714)	616	(486.8 to 921.1)
				2013	22,259	(15,605 to 27,511)	717.2	(498.0 to 885.7)
	YLDs	Ischemic Stroke	Male	1990	45	(28 to 65)	7	(4.4 to 10.2)
				2013	105	(63 to 159)	7.2	(4.3 to 11.0)
			Female	1990	40	(24 to 58)	6	(3.7 to 8.8)
				2013	90	(57 to 132)	5.7	(3.6 to 8.4)
			Both	1990	86	(56 to 119)	6.5	(4.3 to 9.1)
				2013	195	(122 to 278)	6.4	(4.1 to 9.1)
		Hemorrhagic						
		Stroke	Male	1990	133	(85 to 186)	18.2	(11.8 to 25.7)
				2013	315	(207 to 445)	20.3	(13.2 to 29.1)
			Female	1990	91	(62 to 128)	12.3	(8.2 to 17.3)
				2013	204	(135 to 290)	12.4	(8.1 to 17.7)
			Both	1990	225	(153 to 313)	15.2	(10.4 to 21.0)
			Both	2013	519	(347 to 715)	16.2	(10.8 to 22.3)
	Deaths	Ischemic Stroke	Male	1990	124	(69 to 189)	26.8	(15.3 to 38.5)
Papua New				2013	276	(152 to 429)	28.6	(16.8 to 41.2)
Guinea								
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

				•		1	
		Female	1990	95	(58 to 160)	16.4	(10.5 to 27.2)
			2013	211	(137 to 343)	16.9	(11.1 to 27.1)
		Both	1990	220	(137 to 320)	20.6	(13.7 to 28.9)
			2013	487	(308 to 718)	21.6	(14.4 to 30.7)
	Hemorrhagic						
Deaths	Stroke	Male	1990	398	(240 to 594)	55.2	(34.8 to 79.8)
			2013	820	(489 to 1,265)	57.4	(35.5 to 83.3)
		Female	1990	177	(109 to 297)	23.9	(15.0 to 38.4)
			2013	365	(221 to 579)	23.7	(14.9 to 37.0)
		Both	1990	576	(371 to 849)	37.6	(25.3 to 53.3)
			2013	1,185	(743 to 1,747)	38.4	(24.9 to 54.1)
DALYs	Ischemic Stroke	Male	1990	3,442	(1,867 to 5,243)	487.2	(289.2 to 715.3)
			2013	7,091	(3,874 to 11,477)	503.4	(291.8 to 748.5)
		Female	1990	2,399	(1,470 to 4,059)	304.6	(191.1 to 494.4)
			2013	4,864	(3,109 to 7,902)	299.2	(199.5 to 475.9)
		Both	1990	5,842	(3,612 to 8,639)	384.5	(251.6 to 543.8)
			2013	11,955	(7,530 to 18,050)	388.1	(252.3 to 560.3)
	Hemorrhagic						
	Stroke	Male	1990	14,502	(9,059 to 21,931)	1330.3	(813.3 to 1953.2)
			2013	27,727	(17,019 to 43,050)	1341.4	(820.0 to 2038.2)
		Female	1990	5,722	(3,592 to 9,547)	552.5	(344.3 to 912.4)
			2013	10,632	(6,392 to 17,081)	524.1	(321.8 to 824.8)
		Both	1990	20,225	(13,305 to 29,948)	923.6	(605.1 to 1346.2)
			2013	38,360	(24,453 to 57,608)	907.2	(578.7 to 1320.6)
YLDs	Ischemic Stroke	Male	1990	161	(101 to 229)	26.4	(17.5 to 37.9)
			2013	344	(224 to 496)	26.7	(17.5 to 37.9)
		Female	1990	138	(90 to 200)	19.8	(13.0 to 28.6)
			2013	274	(176 to 390)	17.9	(11.9 to 25.5)
		Both	1990	299	(200 to 417)	22.7	(15.4 to 31.7)

				2013	618	(412 to 851)	21.7	(14.7 to 29.5)
		Hemorrhagic						
		Stroke	Male	1990	234	(149 to 338)	31.4	(20.2 to 44.5)
				2013	483	(320 to 666)	31.4	(21.0 to 43.9)
			Female	1990	173	(110 to 240)	21.4	(14.0 to 30.5)
				2013	344	(224 to 484)	19.8	(13.1 to 28.1)
			Both	1990	408	(270 to 578)	25.8	(17.3 to 36.0)
			Both	2013	828	(563 to 1,125)	25	(17.2 to 34.0)
Paraguay	Deaths	Ischemic Stroke	Male	1990	587	(434 to 694)	84.5	(62.9 to 100.9)
				2013	1,055	(791 to 1,594)	62.9	(47.4 to 92.3)
			Female	1990	672	(447 to 771)	79.1	(53.1 to 90.8)
				2013	1,130	(802 to 1,721)	58.3	(41.4 to 87.8)
			Both	1990	1,259	(938 to 1,432)	81.7	(61.4 to 92.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	2,186	(1,703 to 3,043)	60.6	(47.5 to 83.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	340	(265 to 526)	39.3	(30.1 to 63.0)
				2013	1,035	(582 to 1,372)	53	(29.2 to 70.7)
			Female	1990	416	(353 to 609)	42.6	(35.7 to 63.6)
				2013	1,067	(674 to 1,354)	50.8	(31.6 to 64.8)
			Both	1990	756	(645 to 1,077)	41.1	(34.4 to 59.9)
				2013	2,102	(1,386 to 2,562)	52	(33.6 to 63.7)
	DALYs	Ischemic Stroke	Male	1990	10,576	(7,491 to 12,613)	1216.2	(893.8 to 1436.8)
				2013	17,218	(12,583 to 27,699)	895.1	(662.3 to 1391.0)
			Female	1990	10,501	(6,732 to 12,169)	1071.5	(701.6 to 1233.4)
								1
				2013	15,166	(10,644 to 24,464)	738.7	(521.3 to 1176.8)
			Both	2013 1990	15,166 21,077	(10,644 to 24,464) (15,059 to 24,185)	738.7 1139.6	(521.3 to 1176.8) (833.1 to 1299.7)

					•		
			2013	32,385	(24,893 to 47,722)	816.4	(632.3 to 1173.1)
	Hemorrhagic						
	Stroke	Male	1990	8,938	(7,185 to 12,924)	793.8	(629.2 to 1204.1)
			2013	24,263	(14,386 to 31,913)	1067.1	(619.6 to 1411.9)
		Female	1990	9,922	(8,577 to 13,834)	854.3	(732.2 to 1208.7)
			2013	22,277	(15,105 to 28,108)	964.4	(648.0 to 1222.3)
		Both	1990	18,860	(16,396 to 25,374)	824.7	(709.7 to 1139.4)
			2013	46,540	(32,309 to 56,257)	1016.3	(690.6 to 1233.6)
YLDs	Ischemic Stroke	Male	1990	366	(241 to 521)	37.2	(24.9 to 53.0)
			2013	773	(507 to 1,074)	36	(23.3 to 50.2)
		Female	1990	252	(170 to 372)	23.5	(15.6 to 34.9)
			2013	503	(325 to 703)	22.4	(14.5 to 31.4)
		Both	1990	619	(426 to 860)	30	(20.7 to 41.7)
			2013	1,276	(860 to 1,747)	29	(19.7 to 39.6)
	Hemorrhagic						
	Stroke	Male	1990	72	(46 to 103)	6.2	(4.0 to 8.9)
			2013	153	(94 to 226)	6.4	(3.9 to 9.5)
		Female	1990	55	(34 to 78)	4.4	(2.8 to 6.4)
			2013	113	(70 to 161)	4.6	(2.8 to 6.5)
		Both	1990	127	(84 to 177)	5.3	(3.5 to 7.3)
		Both	2013	267	(174 to 378)	5.4	(3.6 to 7.7)
Deaths	Ischemic Stroke	Male	1990	1,597	(1,225 to 1,902)	42.9	(32.2 to 51.3)
			2013	2,490	(1,872 to 3,307)	29.6	(22.5 to 39.1)
		Female	1990	1,820	(1,174 to 2,052)	42.3	(27.1 to 48.0)
			2013	2,677	(2,072 to 4,249)	25.5	(19.8 to 40.5)
		Both	1990	3,417	(2,572 to 3,841)	42.8	(31.8 to 48.2)
			2013	5,168	(4,215 to 7,026)	27.4	(22.4 to 37.0)
	Hemorrhagic			,	,		,
Deaths	Stroke	Male	1990	954	(773 to 1,233)	19.5	(15.5 to 25.9)
	Deaths	YLDs Ischemic Stroke Hemorrhagic Stroke Deaths Ischemic Stroke Hemorrhagic	Stroke Male Female Both YLDs Ischemic Stroke Male Female Both Hemorrhagic Stroke Male Female Both Female Female Female Female Female Both Both Both Both Both Both Both Hemorrhagic Stroke Male Female	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 8,938	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 8,938 (7,185 to 12,924) 793.8

Country	m 00001180	Chroka Tuna	say	Voor	Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year 2013	Mean	95% UI	Mean 19.2	95% UI
			Famala		1,877	(1,319 to 2,500)		(13.3 to 25.6)
			Female	1990	1,095	(936 to 1,548)	21	(17.2 to 29.3)
			5	2013	1,937	(1,341 to 2,656)	17.2	(11.9 to 23.6)
			Both	1990	2,049	(1,795 to 2,684)	20.4	(17.8 to 26.5)
				2013	3,814	(2,955 to 4,771)	18.1	(14.0 to 22.7)
	DALYs	Ischemic Stroke	Male	1990	33,844	(25,511 to 40,622)	654.6	(502.3 to 778.6)
				2013	41,025	(30,125 to 56,082)	413.9	(308.2 to 555.1)
			Female	1990	31,065	(20,219 to 35,192)	569.9	(370.3 to 641.7)
				2013	35,128	(26,422 to 56,538)	314.2	(238.8 to 504.5)
			Both	1990	64,909	(48,985 to 73,558)	611.8	(462.9 to 688.5)
						(60,857 to		
				2013	76,153	106,415)	361.5	(290.5 to 501.5)
		Hemorrhagic		4000	20.222	(24,402 20,022)	422.6	(252.7)
		Stroke	Male	1990	30,223	(24,403 to 39,023)	433.6	(352.7 to 557.7)
				2013	48,048	(33,961 to 64,753)	402	(286.6 to 533.1)
			Female	1990	29,253	(24,670 to 43,885)	424.5	(363.9 to 610.6)
				2013	43,291	(28,953 to 57,628)	345.8	(234.4 to 459.1)
			Both	1990	59,476	(51,109 to 79,555)	430.1	(375.0 to 566.7)
				2010		(69,718 to		(207.0 : 400.4)
			_	2013	91,340	114,733)	373	(285.0 to 466.1)
	YLDs	Ischemic Stroke	Male	1990	759	(499 to 1,078)	13.8	(9.2 to 19.3)
				2013	1,696	(1,103 to 2,398)	15.8	(10.3 to 22.1)
			Female	1990	484	(318 to 681)	8.2	(5.3 to 11.5)
				2013	914	(603 to 1,278)	7.7	(5.1 to 10.9)
			Both	1990	1,243	(854 to 1,726)	10.8	(7.4 to 14.9)
				2013	2,610	(1,774 to 3,555)	11.6	(7.9 to 15.7)

		Hana a unha aia	1 1					I
		Hemorrhagic Stroke	Male	1990	320	(206 to 472)	4.9	(3.1 to 7.2)
				2013	696	(427 to 1,054)	5.8	(3.5 to 9.0)
			Female	1990	221	(135 to 340)	3.2	(1.9 to 5.0)
				2013	447	(281 to 676)	3.5	(2.2 to 5.3)
			Both	1990	542	(354 to 760)	4	(2.6 to 5.6)
			Both	2013	1,143	(726 to 1,650)	4.6	(2.9 to 6.7)
Philippines	Deaths	Ischemic Stroke	Male	1990	5,442	(4,154 to 7,301)	67.1	(50.6 to 91.5)
				2013	16,509	(12,313 to 21,213)	106.3	(80.9 to 134.7)
			Female	1990	4,501	(3,759 to 6,230)	46.6	(38.3 to 65.6)
				2013	14,399	(9,511 to 18,529)	73.9	(49.7 to 93.9)
			Both	1990	9,943	(8,195 to 12,986)	55.6	(45.6 to 73.1)
				2013	30,908	(23,828 to 37,365)	87.6	(68.1 to 104.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	8,526	(6,675 to 10,241)	72.9	(51.8 to 91.7)
				2013	18,133	(13,122 to 24,887)	73.4	(53.2 to 103.0)
			Female	1990	6,239	(4,541 to 7,046)	50.8	(34.9 to 58.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	13,007	(9,731 to 17,363)	48.7	(36.6 to 66.9)
			Both	1990	14,766	(11,931 to 16,879)	61	(45.8 to 72.0)
				2013	31,141	(25,051 to 39,472)	60	(48.4 to 76.8)
						(88,412 to		
	DALYs	Ischemic Stroke	Male	1990	114,108	151,670)	1063	(815.6 to 1410.1)
						(244,714 to		
				2013	338,113	442,811)	1635	(1227.0 to 2089.3)
				4000	75.000	(64,740 to	60.1.0	(FOC F + OCO C)
			Female	1990	75,908	104,025)	634.9	(536.5 to 869.6)
				2013	233,054	(151,450 to	979.2	(650.8 to 1267.6)

					307,095)		
					(158,939 to		
		Both	1990	190,016	245,038)	828.3	(688.2 to 1073.8)
					(430,652 to		
			2013	571,168	702,478)	1268.3	(981.7 to 1534.4)
	Hemorrhagic				(232,987 to		
	Stroke	Male	1990	278,822	331,045)	1714.9	(1352.2 to 2043.3)
					(418,194 to		
			2013	575,817	778,187)	1810.8	(1323.1 to 2468.6)
					(133,072 to		
		Female	1990	167,911	196,131)	1036.4	(789.8 to 1180.0)
					(260,252 to		
			2013	354,102	475,174)	1066.5	(787.0 to 1427.5)
					(385,709 to		
		Both	1990	446,734	501,896)	1361.1	(1131.6 to 1541.1)
					(746,636 to		
			2013	929,919	1,165,740)	1418.7	(1148.3 to 1775.9)
YLDs	Ischemic Stroke	Male	1990	2,852	(1,874 to 4,096)	27.5	(18.4 to 39.1)
			2013	6,340	(4,089 to 8,938)	29.2	(19.0 to 40.5)
		Female	1990	2,775	(1,804 to 3,881)	22.1	(14.1 to 30.9)
			2013	5,992	(3,877 to 8,629)	22.9	(14.7 to 33.3)
		Both	1990	5,628	(3,806 to 7,665)	24.5	(16.5 to 33.1)
			2013	12,333	(8,353 to 16,887)	25.7	(17.5 to 35.4)
	Hemorrhagic			,	, , , , , , , , , , , , , , , , , , , ,		,
	Stroke	Male	1990	3,602	(2,402 to 5,090)	29.8	(19.9 to 42.1)
			2013	7,875	(5,294 to 10,926)	31.6	(21.1 to 43.5)
		Female	1990	3,030	(2,036 to 4,246)	21.6	(14.5 to 30.4)
			2013	6,163	(3,990 to 8,830)	21.4	(13.8 to 30.7)
		Both	1990	6,632	(4,565 to 9,008)	25.3	(17.3 to 34.5)
		Both	2013	14,039	(9,577 to 19,147)	25.9	(17.7 to 35.4)
l .		Dotti	2013	17,000	(3,377 (0 13,147)	23.5	(17.7 to 55.4)

Poland	Deaths	Ischemic Stroke	Male	1990	22,120	(19,194 to 26,716)	168	(145.7 to 202.4)
				2013	22,964	(18,422 to 26,915)	105.7	(84.7 to 124.0)
			Female	1990	32,205	(26,276 to 36,539)	133.1	(108.4 to 151.1)
				2013	30,555	(25,456 to 37,232)	71.2	(60.1 to 86.9)
			Both	1990	54,325	(48,617 to 60,156)	147.1	(131.9 to 163.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	53,520	(46,790 to 60,371)	84.6	(74.2 to 95.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	9,296	(7,029 to 11,463)	62.2	(45.8 to 76.4)
				2013	7,591	(5,739 to 10,475)	33	(24.8 to 46.2)
			Female	1990	12,787	(8,579 to 14,760)	54.1	(36.5 to 62.2)
				2013	7,625	(5,348 to 13,784)	20.7	(15.2 to 35.7)
			Both	1990	22,083	(16,922 to 25,183)	58.2	(44.0 to 66.6)
				2013	15,217	(11,607 to 22,470)	26.2	(20.3 to 37.5)
						(300,510 to		
	DALYs	Ischemic Stroke	Male	1990	346,298	416,317)	2347.1	(2036.2 to 2824.5)
						(248,470 to		
				2013	308,141	360,177)	1362.5	(1102.2 to 1588.0)
				1000		(300,893 to	4==0.0	(10000) (17740)
			Female	1990	373,582	417,175)	1570.2	(1266.9 to 1754.2)
				2013	279,969	(237,170 to 348,332)	724.4	(600 7 to 010 F)
				2015	279,969	(641,975 to	724.4	(609.7 to 910.5)
			Both	1990	719,880	801,605)	1893.5	(1696.2 to 2105.4)
			Dotti	1330	713,000	(518,601 to	1055.5	(1030.2 to 2103.4)
				2013	588,111	655,569)	987.1	(866.0 to 1101.3)
		Hemorrhagic				(168,074 to		
		Stroke	Male	1990	210,531	269,801)	1248	(976.3 to 1581.3)

						(124,775 to		
				2013	165,898	211,423)	700.6	(535.6 to 894.6)
						(154,504 to		
			Female	1990	208,885	236,320)	922	(690.6 to 1045.9)
						(90,080 to		
				2013	115,803	183,740)	373.2	(299.2 to 563.1)
						(352,486 to		4
			Both	1990	419,417	486,521)	1074.8	(896.8 to 1235.7)
				2042	204 702	(232,564 to	522.4	(427.21. (62.2)
				2013	281,702	363,613)	523.1	(437.2 to 663.3)
	YLDs	Ischemic Stroke	Male	1990	3,177	(1,986 to 4,616)	18.5	(11.8 to 26.9)
				2013	5,316	(3,427 to 7,912)	22.4	(14.4 to 33.0)
			Female	1990	2,732	(1,750 to 3,946)	12.2	(7.8 to 17.5)
				2013	3,987	(2,434 to 5,903)	12.8	(7.8 to 19.0)
			Both	1990	5,910	(3,891 to 8,242)	15	(10.0 to 20.8)
				2013	9,304	(6,206 to 12,927)	17.1	(11.3 to 23.8)
		Hemorrhagic						
		Stroke	Male	1990	746	(446 to 1,130)	4.2	(2.5 to 6.3)
				2013	1,120	(645 to 1,721)	4.8	(2.8 to 7.4)
			Female	1990	670	(399 to 1,022)	3.1	(1.9 to 4.7)
				2013	961	(556 to 1,522)	3.4	(2.0 to 5.2)
			Both	1990	1,417	(919 to 2,027)	3.6	(2.3 to 5.1)
			Both	2013	2,082	(1,324 to 2,964)	4	(2.6 to 5.7)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Portugal	Deaths	Ischemic Stroke	Male	1990	6,668	(5,658 to 7,519)	134.2	(113.9 to 152.0)
				2013	5,086	(3,989 to 6,349)	59.1	(46.2 to 74.0)
			Female	1990	9,321	(7,182 to 10,320)	113.5	(87.3 to 125.7)
				2013	6,535	(5,057 to 9,659)	45.2	(34.8 to 67.6)

1		1		T	_		
		Both	1990	15,989	(13,463 to 17,337)	122.6	(103.3 to 133.0)
			2013	11,621	(9,472 to 15,538)	51.1	(41.6 to 68.1)
	Hemorrhagic						
Deaths	Stroke	Male	1990	3,753	(3,030 to 4,690)	72.7	(58.3 to 91.4)
			2013	3,771	(2,585 to 4,782)	45.8	(31.7 to 57.8)
		Female	1990	4,745	(3,668 to 5,686)	60.6	(47.4 to 72.1)
			2013	3,727	(2,896 to 5,444)	28.5	(22.6 to 41.3)
		Both	1990	8,498	(7,161 to 9,812)	66.2	(56.0 to 76.5)
			2013	7,499	(6,245 to 9,468)	36	(30.3 to 45.1)
					(81,318 to		
DALYs	Ischemic Stroke	Male	1990	95,366	107,642)	1805.6	(1539.8 to 2037.0)
			2013	62,674	(50,257 to 77,482)	749	(599.4 to 931.4)
					(79,297 to		
		Female	1990	103,903	115,526)	1315.4	(1000.2 to 1463.5)
			2013	62,957	(49,663 to 91,431)	491.6	(385.6 to 716.5)
					(169,923 to		
		Both	1990	199,270	217,260)	1527.8	(1303.9 to 1665.6)
					(104,012 to		
			2013	125,632	163,136)	602.7	(497.0 to 780.0)
	Hemorrhagic						
	Stroke	Male	1990	70,582	(58,683 to 86,970)	1321.3	(1097.6 to 1624.7)
			2013	58,961	(41,896 to 72,691)	764.5	(553.7 to 934.2)
		Female	1990	68,358	(54,721 to 79,791)	954	(760.1 to 1118.8)
			2013	43,199	(35,215 to 61,686)	404.7	(332.3 to 579.9)
					(121,856 to		
		Both	1990	138,941	159,675)	1121.9	(981.7 to 1286.7)
					(87,699 to		
			2013	102,161	123,891)	568	(488.8 to 680.0)
YLDs	Ischemic Stroke	Male	1990	4,929	(3,205 to 7,035)	91	(59.8 to 130.0)
			2013	6,646	(4,351 to 9,347)	84.1	(55.5 to 118.4)

			Female	1990	5,178	(3,375 to 7,408)	70.9	(46.8 to 100.6)
				2013	8,317	(5,233 to 11,899)	77.4	(49.4 to 110.0)
			Both	1990	10,108	(6,799 to 13,946)	79.6	(53.5 to 109.7)
				2013	14,964	(10,135 to 20,236)	80.5	(54.3 to 109.3)
		Hemorrhagic						
		Stroke	Male	1990	1,266	(687 to 2,077)	23.7	(12.8 to 38.5)
				2013	1,643	(904 to 2,867)	21.3	(12.0 to 37.0)
			Female	1990	1,265	(751 to 1,961)	18.2	(10.8 to 27.9)
				2013	2,211	(1,109 to 3,667)	22	(11.5 to 35.6)
			Both	1990	2,531	(1,645 to 3,706)	20.4	(13.3 to 29.5)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	2013	3,854	(2,371 to 5,731)	21.7	(13.6 to 31.7)
Qatar	Deaths	Ischemic Stroke	Male	1990	17	(12 to 24)	38.7	(28.8 to 53.6)
				2013	30	(23 to 38)	21.7	(16.1 to 28.4)
			Female	1990	12	(9 to 18)	49.7	(35.5 to 70.8)
				2013	17	(12 to 23)	25	(18.2 to 34.6)
			Both	1990	30	(24 to 40)	43.3	(35.3 to 55.3)
				2013	47	(38 to 58)	23.5	(18.9 to 29.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	26	(20 to 35)	25.4	(17.8 to 34.1)
				2013	52	(38 to 69)	12.9	(9.1 to 17.4)
			Female	1990	22	(13 to 27)	48.8	(28.1 to 64.6)
				2013	22	(16 to 30)	19.4	(14.2 to 27.8)
			Both	1990	48	(38 to 58)	34.4	(24.9 to 42.8)
				2013	74	(58 to 92)	16.1	(12.9 to 20.4)
	DALYs	Ischemic Stroke	Male	1990	426	(302 to 625)	472.8	(355.3 to 661.0)
				2013	813	(610 to 1,074)	235.4	(181.9 to 298.2)

			Female	1990	219	(155 to 349)	596.5	(421.4 to 877.1)
				2013	277	(201 to 396)	244.9	(181.0 to 335.8)
			Both	1990	646	(497 to 919)	520.4	(425.0 to 682.4)
				2013	1,090	(876 to 1,358)	242.5	(201.1 to 296.6)
		Hemorrhagic						
		Stroke	Male	1990	1,097	(869 to 1,402)	510.5	(391.3 to 661.4)
				2013	2,053	(1,491 to 2,703)	225.6	(166.3 to 290.5)
			Female	1990	834	(494 to 1,062)	934.1	(534.5 to 1195.1)
				2013	700	(510 to 988)	274.7	(203.5 to 374.3)
			Both	1990	1,931	(1,548 to 2,290)	652.6	(497.3 to 786.8)
				2013	2,753	(2,170 to 3,425)	248.6	(199.0 to 300.9)
	YLDs	Ischemic Stroke	Male	1990	15	(9 to 22)	9.4	(6.0 to 13.9)
				2013	83	(51 to 126)	9.6	(6.3 to 13.8)
			Female	1990	3	(2 to 5)	6.8	(4.3 to 9.9)
				2013	16	(10 to 24)	7	(4.6 to 10.0)
			Both	1990	19	(12 to 27)	8.6	(5.7 to 12.3)
				2013	100	(63 to 145)	8.8	(6.0 to 12.3)
		Hemorrhagic						
		Stroke	Male	1990	2	(1 to 4)	1.3	(0.7 to 2.1)
				2013	16	(8 to 27)	1.4	(0.7 to 2.3)
			Female	1990	1	(0 to 1)	1.5	(0.8 to 2.2)
				2013	4	(2 to 7)	1.6	(0.9 to 2.6)
			Both	1990	4	(2 to 6)	1.3	(0.8 to 2.0)
			Both	2013	21	(12 to 33)	1.5	(0.9 to 2.2)
Romania	Deaths	Ischemic Stroke	Male	1990	15,309	(13,171 to 18,744)	164.4	(140.7 to 201.4)
				2013	18,097	(13,863 to 21,219)	138.2	(106.0 to 162.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

		1 1					
		Female	1990	18,952	(17,265 to 22,846)	143.3	(130.0 to 172.6)
			2013	23,928	(17,698 to 27,064)	111.6	(82.6 to 126.0)
		Both	1990	34,262	(31,279 to 40,227)	152.4	(138.7 to 178.7)
			2013	42,026	(33,300 to 46,954)	122.7	(97.4 to 137.3)
	Hemorrhagic						
Deaths	Stroke	Male	1990	10,368	(7,773 to 12,448)	98.5	(71.2 to 119.1)
			2013	6,939	(5,208 to 9,878)	52.2	(39.1 to 74.8)
		Female	1990	13,093	(9,948 to 14,552)	94.9	(71.1 to 105.6)
			2013	8,563	(6,780 to 12,952)	44	(35.2 to 65.4)
		Both	1990	23,461	(18,665 to 26,370)	97	(75.8 to 109.4)
			2013	15,502	(12,368 to 21,820)	47.9	(38.7 to 66.1)
				,	(198,104 to		,
DALYs	Ischemic Stroke	Male	1990	229,688	280,011)	2231.7	(1932.0 to 2729.5)
				-	(175,739 to		
			2013	229,645	271,579)	1717.7	(1313.5 to 2031.9)
					(215,912 to		
		Female	1990	240,817	286,003)	1750.2	(1581.7 to 2085.0)
					(182,049 to		
			2013	239,726	270,698)	1162.2	(879.4 to 1307.5)
					(431,597 to		
		Both	1990	470,505	551,624)	1963.6	(1796.3 to 2301.9)
					(373,983 to		
			2013	469,371	526,511)	1401.6	(1112.1 to 1575.2)
	Hemorrhagic		4000	246 276	(170,741 to	4070.6	/4.450.0 \
	Stroke	Male	1990	216,270	255,892)	1873.9	(1452.3 to 2229.6)
			2012	125 572	(107,509 to	1011 3	(002 C += 1222 E)
			2013	135,572	179,549)	1011.2	(803.6 to 1333.5)
		Fomala	1990	220 616	(184,593 to	1620.7	(1307.2 to 1811.7)
		Female	1330	229,616	255,051) (107,372 to	1629.7	(1307.2 (0 1811.7)
			2013	129,212	182,041)	735.7	(620.1 to 1021.2)
			2013	143,414	102,041)	/33./	(020.1 (0 1021.2)

						(373,575 to		
			Both	1990	445,887	495,562)	1749.5	(1450.1 to 1947.7)
						(223,066 to		
				2013	264,785	342,116)	864.7	(736.1 to 1111.7)
	YLDs	Ischemic Stroke	Male	1990	2,715	(1,697 to 3,829)	23.3	(14.9 to 32.9)
				2013	3,454	(2,254 to 4,972)	25.9	(17.0 to 37.2)
			Female	1990	2,188	(1,415 to 3,101)	15.8	(10.2 to 22.3)
				2013	2,413	(1,521 to 3,664)	14.3	(9.1 to 21.7)
			Both	1990	4,904	(3,293 to 6,756)	19.3	(13.0 to 26.5)
				2013	5,868	(3,884 to 8,182)	19.5	(13.0 to 27.2)
		Hemorrhagic						
		Stroke	Male	1990	572	(361 to 881)	4.9	(3.1 to 7.4)
				2013	710	(426 to 1,059)	5.5	(3.3 to 8.1)
			Female	1990	499	(282 to 784)	3.7	(2.1 to 5.7)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	559	(331 to 854)	3.6	(2.2 to 5.5)
			Both	1990	1,071	(683 to 1,548)	4.3	(2.7 to 6.1)
			Both	2013	1,269	(832 to 1,821)	4.5	(2.9 to 6.4)
Russia	Deaths	Ischemic Stroke	Male	1990	85,667	(70,113 to 98,839)	222.2	(186.3 to 251.9)
					05,007	(70,113 to 30,033)	222.2	(100.0 to 201.0)
					83,007	(100,042 to	222.2	(100.5 to 251.5)
				2013	115,629	, , , , ,	182.5	(158.8 to 214.5)
				2013	115,629	(100,042 to 136,462) (172,085 to	182.5	(158.8 to 214.5)
			Female		·	(100,042 to 136,462) (172,085 to 230,317)		,
				2013 1990	115,629 205,553	(100,042 to 136,462) (172,085 to 230,317) (165,542 to	182.5 189.7	(158.8 to 214.5) (158.2 to 213.3)
				2013	115,629	(100,042 to 136,462) (172,085 to 230,317) (165,542 to 248,975)	182.5	(158.8 to 214.5)
			Female	2013 1990 2013	115,629 205,553 187,190	(100,042 to 136,462) (172,085 to 230,317) (165,542 to 248,975) (251,325 to	182.5 189.7 126.7	(158.8 to 214.5) (158.2 to 213.3) (112.4 to 168.7)
				2013 1990	115,629 205,553	(100,042 to 136,462) (172,085 to 230,317) (165,542 to 248,975)	182.5 189.7	(158.8 to 214.5) (158.2 to 213.3)

					373,857)		
	Homorrhagia				3/3,03/1		
Deaths	Hemorrhagic Stroke	Male	1990	21,460	(12,103 to 32,283)	40	(22.8 to 60.4)
Deatils	JUOKE	IVIAIC	2013	29,237	(21,781 to 36,149)	39.9	(29.9 to 49.3)
							•
		Female	1990	34,213	(20,004 to 47,242)	32.9	(19.7 to 45.1)
			2013	28,230	(23,499 to 40,791)	22.4	(19.0 to 32.4)
		Both	1990	55,674	(34,690 to 72,420)	35.3	(22.0 to 45.8)
			2013	57,468	(49,156 to 70,248)	29.3	(25.4 to 35.3)
					(1,159,308 to		
DALYs	Ischemic Stroke	Male	1990	1,477,306	1,737,714)	3127.6	(2550.9 to 3623.7)
					(1,619,270 to		
			2013	1,881,206	2,242,051)	2714.5	(2345.0 to 3202.3)
					(2,124,018 to		
		Female	1990	2,598,496	2,940,322)	2388.1	(1953.0 to 2710.1)
					(1,857,643 to		
			2013	2,108,635	2,820,221)	1534	(1355.9 to 2050.9)
					(3,486,591 to		
		Both	1990	4,075,802	4,567,371)	2632.4	(2257.2 to 2933.4)
					(3,588,377 to		
			2013	3,989,842	4,911,632)	1971.2	(1776.4 to 2411.9)
	Hemorrhagic				(330,922 to		
	Stroke	Male	1990	569,556	844,376)	898.5	(522.4 to 1326.9)
					(587,283 to		
			2013	768,337	944,663)	975.1	(748.4 to 1185.8)
					(457,574 to		
		Female	1990	723,083	937,558)	735	(480.4 to 944.0)
					(483,138 to		
			2013	571,526	822,528)	506.6	(427.5 to 743.9)
					(845,938 to		
		Both	1990	1,292,639	1,651,989)	796.4	(528.9 to 1012.2)
			2013	1,339,864	(1,151,769 to	704.2	(614.0 to 836.7)

						1,590,066)		
	YLDs	Ischemic Stroke	Male	1990	14,986	(9,264 to 22,787)	23.9	(15.0 to 34.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	19,422	(11,603 to 30,416)	25.3	(15.4 to 39.3)
			Female	1990	18,694	(11,929 to 26,923)	18.6	(12.0 to 26.9)
				2013	19,498	(11,354 to 29,352)	16.7	(9.9 to 25.1)
			Both	1990	33,680	(22,467 to 45,757)	20.8	(13.9 to 28.4)
				2013	38,921	(25,222 to 56,256)	20.1	(13.2 to 28.8)
		Hemorrhagic						
		Stroke	Male	1990	2,048	(1,142 to 3,268)	3.1	(1.8 to 4.8)
				2013	2,329	(1,347 to 3,591)	3	(1.7 to 4.7)
			Female	1990	2,154	(1,314 to 3,454)	2.2	(1.4 to 3.6)
				2013	2,281	(1,350 to 3,571)	2.1	(1.3 to 3.2)
			Both	1990	4,202	(2,696 to 6,080)	2.6	(1.7 to 3.7)
			Both	2013	4,610	(2,919 to 6,632)	2.5	(1.6 to 3.6)
Rwanda	Deaths	Ischemic Stroke	Male	1990	846	(222 to 1,320)	123.8	(33.4 to 187.2)
				2013	803	(163 to 1,274)	65.4	(14.0 to 102.3)
			Female	1990	1,098	(406 to 1,774)	128.1	(51.0 to 201.3)
				2013	865	(285 to 1,422)	60.3	(20.4 to 99.2)
			Both	1990	1,945	(931 to 2,770)	126.9	(64.3 to 177.0)
				2013	1,669	(629 to 2,390)	62.6	(25.1 to 89.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,645	(1,146 to 2,377)	171.2	(115.9 to 261.4)
				2013	1,269	(893 to 1,994)	82	(55.6 to 138.1)
			Female	1990	1,836	(1,072 to 3,024)	164.7	(97.7 to 285.1)
				2013	1,255	(766 to 2,391)	72	(41.9 to 143.4)
			Both	1990	3,482	(2,530 to 4,901)	167.9	(119.0 to 245.7)

				2013	2,524	(1,800 to 3,932)	76.6	(52.4 to 125.8)
	DALYs	Ischemic Stroke	Male	1990	18,464	(4,607 to 29,409)	2002	(545.8 to 3091.0)
				2013	13,981	(2,858 to 22,798)	923.7	(195.4 to 1470.4)
			Female	1990	20,823	(6,840 to 36,773)	1958.5	(718.1 to 3179.1)
				2013	13,021	(4,128 to 21,771)	775.2	(256.9 to 1270.9)
			Both	1990	39,288	(18,443 to 59,720)	1989.1	(953.7 to 2829.6)
				2013	27,003	(9,578 to 39,156)	842.9	(311.0 to 1204.7)
		Hemorrhagic						
		Stroke	Male	1990	48,299	(34,542 to 66,581)	3670.4	(2545.3 to 5336.4)
				2013	31,080	(22,491 to 46,140)	1500.9	(1060.9 to 2317.4)
			Female	1990	48,959	(28,187 to 78,376)	3431.1	(1991.1 to 5628.8)
				2013	26,741	(16,796 to 47,921)	1206.6	(739.4 to 2248.6)
						(70,779 to		
			Both	1990	97,258	136,401)	3548.4	(2577.0 to 4998.4)
				2013	57,822	(42,136 to 84,042)	1342.9	(966.2 to 2056.7)
	YLDs	Ischemic Stroke	Male	1990	135	(82 to 204)	10	(5.9 to 15.1)
				2013	246	(157 to 355)	10.6	(6.8 to 15.3)
			Female	1990	112	(72 to 164)	7.9	(5.1 to 11.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	204	(122 to 304)	7.5	(4.3 to 11.1)
			Both	1990	248	(166 to 354)	8.9	(6.0 to 12.5)
				2013	451	(296 to 633)	8.9	(5.9 to 12.3)
		Hemorrhagic						
		Stroke	Male	1990	46	(27 to 70)	2.7	(1.6 to 4.1)
				2013	90	(54 to 140)	3	(1.9 to 4.5)
			Female	1990	32	(19 to 48)	1.9	(1.1 to 2.7)
				2013	63	(37 to 99)	1.9	(1.1 to 3.0)

	-	T	,					
			Both	1990	78	(50 to 113)	2.2	(1.4 to 3.2)
			Both	2013	154	(97 to 227)	2.4	(1.5 to 3.4)
Saint Lucia	Deaths	Ischemic Stroke	Male	1990	43	(35 to 52)	103.2	(86.3 to 126.1)
				2013	51	(38 to 66)	69.7	(51.9 to 89.6)
			Female	1990	56	(45 to 65)	110	(88.7 to 128.0)
				2013	70	(47 to 92)	71.3	(48.6 to 93.2)
			Both	1990	99	(87 to 113)	107.8	(93.9 to 122.5)
				2013	122	(93 to 149)	71.2	(54.3 to 86.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	22	(17 to 28)	52.2	(41.3 to 66.9)
				2013	29	(20 to 41)	38.7	(27.0 to 53.4)
			Female	1990	30	(24 to 38)	60.3	(48.4 to 76.3)
				2013	39	(26 to 53)	40.5	(28.0 to 54.9)
			Both	1990	53	(46 to 63)	56.9	(48.9 to 68.3)
				2013	68	(51 to 86)	40	(30.1 to 50.1)
	DALYs	Ischemic Stroke	Male	1990	774	(646 to 930)	1690.9	(1414.1 to 2025.3)
				2013	811	(604 to 1,054)	1059.6	(792.6 to 1375.3)
			Female	1990	858	(692 to 994)	1611.1	(1280.1 to 1865.3)
				2013	850	(586 to 1,117)	919.1	(632.8 to 1216.0)
			Both	1990	1,633	(1,427 to 1,836)	1656.9	(1445.7 to 1862.8)
				2013	1,661	(1,289 to 2,016)	990.2	(767.6 to 1205.0)
		Hemorrhagic						
		Stroke	Male	1990	531	(433 to 653)	1102.4	(890.4 to 1371.2)
				2013	618	(440 to 846)	760.3	(544.3 to 1034.0)
			Female	1990	625	(502 to 770)	1184.4	(944.3 to 1457.4)
				2013	638	(458 to 873)	697.5	(499.2 to 950.9)
			Both	1990	1,156	(1,016 to 1,347)	1148.5	(1007.6 to 1333.0)
				2013	1,257	(972 to 1,553)	730	(562.8 to 905.1)
	YLDs	Ischemic Stroke	Male	1990	20	(13 to 29)	47	(31.4 to 66.2)

				2013	36	(24 to 50)	46.6	(30.8 to 66.3)
			Female	1990	17	(10 to 25)	34	(21.4 to 48.8)
				2013	32	(19 to 44)	34.9	(21.8 to 48.8)
			Both	1990	38	(25 to 52)	40	(27.3 to 54.7)
				2013	68	(45 to 92)	40.4	(26.8 to 54.8)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	6	(4 to 10)	14.7	(8.9 to 22.6)
				2013	11	(6 to 17)	14.3	(8.4 to 22.0)
			Female	1990	5	(3 to 9)	10.9	(6.4 to 17.2)
				2013	10	(6 to 15)	11.5	(7.3 to 17.1)
			Both	1990	12	(8 to 17)	12.7	(8.2 to 17.8)
			Both	2013	22	(14 to 31)	12.8	(8.2 to 18.4)
Saint Vincent and	Deaths	Ischemic Stroke	Male	1990	25	(20 to 30)	95.9	(76.3 to 114.5)
the Grenadines				2013	25	(19 to 32)	72.5	(55.7 to 92.5)
			Female	1990	39	(30 to 45)	105.8	(83.0 to 123.8)
				2013	33	(24 to 41)	76.6	(55.6 to 93.5)
			Both	1990	64	(54 to 72)	102.1	(86.2 to 115.0)
				2013	59	(48 to 70)	75.5	(61.0 to 89.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	13	(10 to 16)	47.9	(37.8 to 59.7)
				2013	15	(10 to 19)	38.3	(27.6 to 50.4)
			Female	1990	22	(16 to 27)	61.7	(46.1 to 75.6)
				2013	19	(14 to 25)	42.7	(31.0 to 57.6)
			Both	1990	36	(30 to 42)	56	(46.4 to 65.5)
				2013	34	(27 to 42)	41	(32.6 to 51.5)
	DALYs	Ischemic Stroke	Male	1990	451	(364 to 533)	1530.2	(1243.5 to 1818.6)

				2013	452	(347 to 576)	1123.3	(869.7 to 1425.4)
			Female	1990	580	(459 to 669)	1556.7	(1220.7 to 1803.6)
				2013	462	(336 to 572)	1029.6	(743.5 to 1265.7)
			Both	1990	1,031	(871 to 1,148)	1550.6	(1306.0 to 1736.5)
				2013	915	(743 to 1,085)	1082.2	(877.9 to 1282.0)
		Hemorrhagic						
		Stroke	Male	1990	337	(266 to 417)	1018.9	(812.2 to 1246.1)
				2013	358	(265 to 461)	780.3	(572.4 to 1002.7)
			Female	1990	463	(361 to 550)	1216.8	(943.9 to 1449.5)
				2013	367	(273 to 482)	776.8	(576.3 to 1020.0)
			Both	1990	801	(675 to 915)	1130.2	(954.6 to 1291.2)
				2013	726	(593 to 882)	782.4	(638.7 to 958.2)
	YLDs	Ischemic Stroke	Male	1990	12	(7 to 18)	44	(27.7 to 64.9)
				2013	17	(10 to 24)	42.3	(27.3 to 60.1)
			Female	1990	11	(7 to 16)	31.4	(20.8 to 45.6)
				2013	14	(9 to 20)	32.7	(21.1 to 46.0)
			Both	1990	24	(15 to 33)	36.9	(24.4 to 51.3)
				2013	32	(21 to 43)	37.2	(25.0 to 50.7)
		Hemorrhagic						
		Stroke	Male	1990	4	(2 to 6)	13.8	(8.2 to 21.7)
				2013	5	(3 to 8)	13.3	(8.5 to 19.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	3	(2 to 5)	9.5	(6.0 to 14.2)
				2013	5	(3 to 8)	11.4	(6.7 to 18.5)
			Both	1990	7	(5 to 11)	11.5	(7.5 to 16.7)
			Both	2013	11	(7 to 16)	12.3	(8.0 to 17.6)
Samoa	Deaths	Ischemic Stroke	Male	1990	21	(9 to 34)	83	(38.7 to 124.4)

			2013	26	(13 to 44)	65.3	(33.5 to 102.6)
		Female	1990	11	(4 to 21)	40.7	(17.2 to 75.1)
			2013	24	(11 to 41)	42.4	(19.0 to 71.2)
		Both	1990	33	(20 to 53)	59.7	(36.8 to 91.8)
			2013	51	(33 to 78)	51.8	(34.1 to 76.9)
	Hemorrhagic						
Deaths	Stroke	Male	1990	42	(25 to 59)	127.6	(76.3 to 178.4)
			2013	41	(22 to 59)	86.4	(47.1 to 123.5)
		Female	1990	20	(11 to 36)	61.7	(34.4 to 110.6)
			2013	35	(21 to 59)	60.2	(35.8 to 99.9)
		Both	1990	62	(45 to 86)	92.3	(67.5 to 129.2)
			2013	77	(52 to 105)	72	(48.6 to 98.5)
DALYs	Ischemic Stroke	Male	1990	460	(194 to 801)	1357.3	(600.1 to 2201.7)
			2013	479	(212 to 913)	968.8	(471.4 to 1684.0)
		Female	1990	215	(90 to 429)	609.6	(262.0 to 1136.1)
			2013	370	(158 to 673)	617.1	(266.5 to 1111.9)
		Both	1990	676	(374 to 1,193)	960	(567.5 to 1575.4)
			2013	849	(522 to 1,413)	773.5	(487.3 to 1247.9)
	Hemorrhagic						
	Stroke	Male	1990	1,204	(727 to 1,738)	2795.7	(1703.6 to 3964.1)
			2013	1,045	(598 to 1,506)	1730.9	(974.6 to 2469.9)
		Female	1990	479	(278 to 857)	1154.5	(661.9 to 2069.2)
			2013	705	(430 to 1,127)	1120.2	(681.0 to 1800.9)
		Both	1990	1,684	(1,178 to 2,329)	1944.7	(1379.3 to 2690.4)
			2013	1,750	(1,215 to 2,354)	1416.8	(985.2 to 1896.0)
YLDs	Ischemic Stroke	Male	1990	8	(5 to 11)	25.1	(15.6 to 35.3)
			2013	12	(8 to 17)	25.2	(16.5 to 35.7)
		Female	1990	7	(5 to 10)	20.9	(13.8 to 30.1)
			2013	11	(7 to 16)	20.2	(12.6 to 29.0)

Both 1990 15 (10 to 21) 22.8 2013 24 (16 to 33) 22.5 Hemorrhagic	,
	(14.9 to 31.0)
Hamorrhagic	(14.5 to 51.0)
nemormagic	
Stroke Male 1990 10 (7 to 15) 29.9	(18.9 to 42.5)
2013 16 (10 to 23) 30.3	(19.2 to 42.9)
Female 1990 8 (5 to 12) 22.3	(14.4 to 31.4)
2013 13 (8 to 18) 21.7	(13.9 to 30.7)
Both 1990 19 (13 to 27) 25.8	(17.2 to 35.3)
Counts, All Ages Ra	te Per 100,000
Aę	e- standardized
Country measure Stroke Type sex year Mean 95% UI Mea	n 95% UI
Both 2013 29 (19 to 41) 25.6	(16.7 to 35.7)
Sao Tome and Deaths Ischemic Stroke Male 1990 12 (9 to 16) 53.4	(40.1 to 70.6)
Principe 2013 18 (11 to 27) 58.3	(34.9 to 85.0)
Female 1990 17 (12 to 22) 62.6	(46.5 to 82.7)
2013 24 (14 to 34) 66.5	(38.8 to 93.6)
Both 1990 29 (24 to 36) 58.6	(46.9 to 73.1)
2013 43 (28 to 57) 62.8	(40.1 to 83.3)
Hemorrhagic	
Deaths Stroke Male 1990 24 (19 to 30) 89.3	(70.7 to 111.8)
2013 35 (22 to 49) 96.3	(65.1 to 135.0)
Female 1990 30 (23 to 38) 99.4	(77.9 to 125.5)
2013 41 (27 to 55) 101.	6 (69.0 to 138.8)
Both 1990 55 (46 to 65) 95.3	(80.6 to 113.4)
2013 76 (54 to 97) 99.3	(72.9 to 125.6)
DALYs Ischemic Stroke Male 1990 288 (215 to 373) 938.	1 (718.5 to 1218.3)
2013 414 (238 to 633) 1002	8 (602.6 to 1496.7)
Female 1990 342 (240 to 452) 1044	3 (751.8 to 1363.0)
2013 477 (290 to 688) 1060	5 (633.1 to 1529.6)

			Both	1990	631	(507 to 779)	995.3	(819.5 to 1219.5)
				2013	892	(587 to 1,198)	1036.3	(673.0 to 1381.5)
		Hemorrhagic						
		Stroke	Male	1990	863	(661 to 1,077)	2005.1	(1589.5 to 2458.9)
				2013	1,087	(637 to 1,495)	1989.1	(1197.3 to 2789.9)
			Female	1990	879	(607 to 1,134)	2051.8	(1541.5 to 2583.8)
				2013	1,047	(652 to 1,431)	1900.2	(1225.9 to 2658.5)
			Both	1990	1,742	(1,359 to 2,075)	2031.8	(1689.3 to 2380.1)
				2013	2,135	(1,380 to 2,768)	1948.1	(1324.8 to 2527.6)
	YLDs	Ischemic Stroke	Male	1990	4	(2 to 6)	15.6	(10.4 to 22.1)
				2013	6	(4 to 9)	15.6	(10.2 to 22.6)
			Female	1990	3	(2 to 5)	11.4	(7.5 to 16.7)
				2013	5	(3 to 8)	11.9	(7.4 to 17.6)
			Both	1990	7	(5 to 10)	13.4	(9.2 to 18.3)
				2013	12	(8 to 16)	13.6	(9.2 to 18.6)
		Hemorrhagic						
		Stroke	Male	1990	1	(0 to 1)	4.2	(2.7 to 6.4)
				2013	1	(1 to 2)	4	(2.5 to 5.8)
			Female	1990	0	(0 to 1)	2.9	(1.8 to 4.1)
				2013	1	(0 to 2)	2.8	(1.7 to 4.0)
			Both	1990	2	(1 to 3)	3.5	(2.3 to 4.9)
			Both	2013	3	(2 to 5)	3.3	(2.2 to 4.7)
Saudi Arabia	Deaths	Ischemic Stroke	Male	1990	1,968	(1,443 to 2,677)	84.8	(63.4 to 113.5)
				2013	3,080	(2,372 to 4,094)	69	(53.5 to 90.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	1,713	(1,276 to 2,248)	78.8	(58.9 to 103.9)
				2013	2,879	(2,229 to 3,446)	59.2	(46.2 to 70.5)

		Both	1990	3,681	(3,037 to 4,543)	82	(68.1 to 100.5)
			2013	5,959	(5,057 to 6,996)	64.6	(54.9 to 75.7)
	Hemorrhagic						
Deaths	Stroke	Male	1990	1,150	(780 to 1,639)	41	(27.6 to 57.5)
			2013	2,024	(1,459 to 2,779)	34.4	(24.5 to 47.6)
		Female	1990	1,016	(682 to 1,365)	41	(27.0 to 54.4)
			2013	1,512	(1,141 to 1,956)	27.6	(20.9 to 35.8)
		Both	1990	2,166	(1,583 to 2,703)	40.9	(29.6 to 50.8)
			2013	3,537	(2,904 to 4,337)	31.2	(25.5 to 38.4)
DALYs	Ischemic Stroke	Male	1990	31,685	(22,287 to 45,291)	1116.8	(796.4 to 1566.6)
			2013	52,567	(39,534 to 71,530)	879.2	(674.3 to 1177.8)
		Female	1990	23,541	(16,736 to 31,485)	982.8	(702.7 to 1303.3)
			2013	33,052	(24,869 to 40,072)	651.5	(500.5 to 782.5)
		Both	1990	55,226	(44,307 to 71,125)	1056.6	(849.0 to 1333.6)
					(70,924 to		
			2013	85,619	104,591)	774.7	(646.3 to 939.7)
	Hemorrhagic						
	Stroke	Male	1990	28,823	(19,367 to 41,937)	720.9	(479.6 to 1056.8)
			2013	49,996	(36,249 to 68,191)	581.8	(415.1 to 801.0)
		Female	1990	24,764	(16,380 to 34,674)	748.6	(488.1 to 1037.4)
			2013	29,710	(22,732 to 37,803)	441.7	(340.2 to 560.3)
		Both	1990	53,587	(39,976 to 67,479)	729.7	(535.9 to 929.8)
			2013	79,707	(64,764 to 98,588)	517.4	(418.5 to 643.5)
YLDs	Ischemic Stroke	Male	1990	204	(129 to 296)	5.2	(3.2 to 7.8)
			2013	615	(380 to 916)	6.2	(4.0 to 9.0)
		Female	1990	70	(44 to 97)	2.6	(1.6 to 3.6)
			2013	222	(140 to 321)	3.2	(1.9 to 4.6)
		Both	1990	274	(181 to 385)	4.1	(2.7 to 5.8)
			2013	837	(544 to 1,204)	4.8	(3.3 to 6.9)

		Hemorrhagic						
		Stroke	Male	1990	18	(11 to 27)	0.4	(0.2 to 0.6)
				2013	57	(33 to 87)	0.5	(0.3 to 0.8)
			Female	1990	4	(2 to 6)	0.1	(0.1 to 0.2)
				2013	14	(8 to 22)	0.2	(0.1 to 0.3)
			Both	1990	22	(14 to 33)	0.3	(0.2 to 0.4)
			Both	2013	72	(44 to 105)	0.4	(0.2 to 0.5)
Senegal	Deaths	Ischemic Stroke	Male	1990	667	(436 to 887)	70.8	(46.6 to 92.6)
				2013	1,684	(979 to 2,237)	99.7	(61.1 to 130.2)
			Female	1990	843	(495 to 1,110)	80.7	(49.4 to 107.5)
				2013	2,577	(1,142 to 3,488)	122	(56.4 to 162.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	1,511	(1,100 to 1,841)	76.7	(56.3 to 93.8)
				2013	4,262	(2,532 to 5,339)	112.6	(67.2 to 139.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	643	(445 to 910)	52.2	(34.2 to 76.6)
				2013	921	(545 to 1,568)	42.6	(23.3 to 77.0)
			Female	1990	882	(630 to 1,241)	68.1	(46.0 to 99.4)
				2013	1,447	(929 to 2,596)	56.1	(34.3 to 108.3)
			Both	1990	1,526	(1,155 to 2,031)	61.2	(44.2 to 85.1)
				2013	2,368	(1,602 to 3,838)	50.4	(32.5 to 88.0)
	DALYs	Ischemic Stroke	Male	1990	14,818	(9,393 to 20,418)	1133	(748.8 to 1503.0)
				2013	33,028	(18,296 to 44,511)	1545.5	(883.6 to 2045.0)
			Female	1990	16,502	(9,503 to 21,683)	1182.6	(695.1 to 1545.5)
				2013	44,666	(19,108 to 62,303)	1743	(757.2 to 2381.1)
			Both	1990	31,320	(22,558 to 38,373)	1165.1	(846.8 to 1418.3)
				2013	77,694	(45,897 to 98,731)	1660.7	(985.8 to 2089.3)

		Hemorrhagic						
		Stroke	Male	1990	19,380	(14,009 to 26,578)	1065.6	(737.2 to 1503.8)
				2013	26,291	(17,154 to 40,813)	879.1	(521.0 to 1486.3)
			Female	1990	23,577	(17,168 to 32,605)	1273.5	(900.6 to 1784.1)
				2013	35,743	(24,622 to 57,313)	1078.7	(706.2 to 1865.4)
			Both	1990	42,957	(33,946 to 54,643)	1180.3	(891.3 to 1572.5)
				2013	62,034	(44,996 to 88,958)	993.3	(681.6 to 1579.7)
	YLDs	Ischemic Stroke	Male	1990	243	(156 to 348)	15.3	(9.8 to 21.8)
				2013	415	(261 to 611)	15.1	(9.6 to 22.2)
			Female	1990	196	(127 to 288)	11.3	(7.2 to 16.5)
				2013	397	(253 to 563)	11.7	(7.5 to 16.8)
			Both	1990	439	(290 to 609)	13.2	(8.7 to 18.1)
				2013	813	(539 to 1,137)	13.2	(8.9 to 18.4)
		Hemorrhagic						
		Stroke	Male	1990	77	(48 to 118)	4.1	(2.6 to 6.1)
				2013	132	(80 to 194)	3.9	(2.4 to 5.9)
			Female	1990	56	(35 to 83)	2.8	(1.8 to 4.2)
				2013	110	(67 to 164)	2.8	(1.7 to 4.2)
			Both	1990	133	(86 to 190)	3.4	(2.2 to 4.9)
			Both	2013	243	(157 to 349)	3.3	(2.1 to 4.7)
Serbia	Deaths	Ischemic Stroke	Male	1990	4,743	(3,687 to 5,932)	137.7	(107.8 to 171.5)
				2013	5,615	(4,580 to 6,553)	103.3	(84.2 to 120.9)
			Female	1990	5,565	(4,330 to 6,839)	111.2	(86.3 to 136.8)
				2013	7,575	(6,252 to 8,511)	87.7	(72.2 to 98.6)
			Both	1990	10,308	(8,605 to 12,114)	122.3	(102.0 to 143.2)
				2013	13,190	(11,332 to 14,526)	94.5	(81.2 to 104.2)
					Counts, All	٨σ٥٥	Data	Per 100 000

Counts, All Ages

Rate Per 100,000 Age- standardized

Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,641	(1,128 to 2,234)	37.3	(25.9 to 51.1)
				2013	1,443	(1,131 to 1,808)	24.8	(19.2 to 31.4)
			Female	1990	2,068	(1,327 to 2,687)	37.2	(24.0 to 48.6)
				2013	1,720	(1,431 to 2,386)	22.1	(18.4 to 30.2)
			Both	1990	3,710	(2,678 to 4,602)	37.4	(26.9 to 46.3)
				2013	3,163	(2,758 to 3,911)	23.4	(20.4 to 28.8)
	DALYs	Ischemic Stroke	Male	1990	71,413	(54,023 to 91,443)	1843.6	(1417.3 to 2329.8)
				2013	75,974	(62,020 to 88,874)	1318.5	(1082.0 to 1541.5)
			Female	1990	68,926	(53,051 to 85,488)	1330.1	(1022.1 to 1643.2)
				2013	79,390	(65,972 to 89,540)	962.4	(798.5 to 1085.7)
						(115,430 to		
			Both	1990	140,339	165,934)	1551.6	(1290.7 to 1833.6)
						(134,004 to		
				2013	155,364	171,970)	1120.8	(962.3 to 1238.8)
		Hemorrhagic	Mala	1000	44.053	(20.700 to EC.400)	0.67.6	(C12 0 to 1100 1)
		Stroke	Male	1990	41,952	(29,709 to 56,468)	867.6	(613.9 to 1180.1)
			- 1	2013	32,015	(25,764 to 39,762)	541	(436.7 to 668.9)
			Female	1990	46,188	(30,139 to 59,545)	816.5	(540.2 to 1047.1)
				2013	31,114	(26,449 to 42,067)	442.9	(378.9 to 596.9)
			Both	1990	88,141	(65,664 to 108,100)	842.2	(630.4 to 1035.3)
			DOTT	2013	63,130	(55,652 to 77,050)	489.9	(433.3 to 594.6)
	YLDs	Ischemic Stroke	Male	1990	1,068	(700 to 1,490)	22.9	(15.1 to 31.9)
	ILDS	ischennic stroke	iviale			· ' '		, , ,
			Famala	2013	1,531	(999 to 2,154)	25.8	(17.0 to 36.2)
			Female	1990	816	(503 to 1,149)	14.8	(9.3 to 20.9)
			5.1	2013	1,007	(640 to 1,434)	14.2	(9.1 to 20.4)
			Both	1990	1,885	(1,253 to 2,576)	18.5	(12.3 to 25.2)
				2013	2,539	(1,725 to 3,450)	19.5	(13.3 to 26.6)

		Hemorrhagic						
		Stroke	Male	1990	234	(134 to 360)	4.9	(2.9 to 7.5)
				2013	315	(188 to 481)	5.5	(3.3 to 8.4)
			Female	1990	184	(113 to 289)	3.4	(2.1 to 5.2)
				2013	235	(139 to 350)	3.6	(2.2 to 5.2)
			Both	1990	419	(274 to 604)	4.1	(2.7 to 5.9)
			Both	2013	551	(348 to 778)	4.5	(2.8 to 6.4)
Seychelles	Deaths	Ischemic Stroke	Male	1990	21	(16 to 27)	113.8	(83.2 to 141.4)
				2013	23	(14 to 31)	78.2	(50.1 to 105.4)
			Female	1990	18	(13 to 23)	59.8	(46.0 to 76.1)
				2013	26	(12 to 38)	52.4	(24.2 to 78.1)
			Both	1990	40	(32 to 46)	82	(66.4 to 95.6)
				2013	49	(30 to 65)	63.5	(38.9 to 83.7)
					C	•	D-4-	D 100 000
					Counts, All	Ages	Rate	Per 100,000
					Counts, All	Ages	Rate Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	Ages 95% UI		
Country		Hemorrhagic		·	Mean	95% UI	Age- Mean	standardized 95% UI
Country	measure Deaths		sex Male	1990	Mean 19	95% UI (14 to 25)	Age- Mean 93.3	standardized 95% UI (70.3 to 126.1)
Country		Hemorrhagic	Male	1990 2013	Mean 19 19	95% UI (14 to 25) (10 to 30)	Age- Mean 93.3 58.1	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4)
Country		Hemorrhagic		1990	Mean 19	95% UI (14 to 25)	Age- Mean 93.3	standardized 95% UI (70.3 to 126.1)
Country		Hemorrhagic	Male	1990 2013	Mean 19 19	95% UI (14 to 25) (10 to 30)	Age- Mean 93.3 58.1	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4)
Country		Hemorrhagic	Male	1990 2013 1990	Mean 19 19 18	95% UI (14 to 25) (10 to 30) (14 to 28)	Age- Mean 93.3 58.1 63.8	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0)
Country		Hemorrhagic	Male Female	1990 2013 1990 2013	Mean 19 19 18 23	95% UI (14 to 25) (10 to 30) (14 to 28) (14 to 39)	Age- Mean 93.3 58.1 63.8 48.6	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0) (30.0 to 82.0)
Country		Hemorrhagic	Male Female	1990 2013 1990 2013 1990	Mean 19 19 18 23 38	95% UI (14 to 25) (10 to 30) (14 to 28) (14 to 39) (30 to 52)	Age- Mean 93.3 58.1 63.8 48.6 77.4	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0) (30.0 to 82.0) (61.7 to 105.6)
Country	Deaths	Hemorrhagic Stroke	Male Female Both	1990 2013 1990 2013 1990 2013	Mean 19 19 18 23 38 43	95% UI (14 to 25) (10 to 30) (14 to 28) (14 to 39) (30 to 52) (28 to 66)	Age-Mean 93.3 58.1 63.8 48.6 77.4 53.8	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0) (30.0 to 82.0) (61.7 to 105.6) (35.6 to 83.9)
Country	Deaths	Hemorrhagic Stroke	Male Female Both	1990 2013 1990 2013 1990 2013 1990	Mean 19 19 18 23 38 43 406	95% UI (14 to 25) (10 to 30) (14 to 28) (14 to 39) (30 to 52) (28 to 66) (293 to 522)	Age-Mean 93.3 58.1 63.8 48.6 77.4 53.8 1967	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0) (30.0 to 82.0) (61.7 to 105.6) (35.6 to 83.9) (1434.3 to 2515.0)
Country	Deaths	Hemorrhagic Stroke	Male Female Both Male	1990 2013 1990 2013 1990 2013 1990 2013	Mean 19 19 18 23 38 43 406 434	95% UI (14 to 25) (10 to 30) (14 to 28) (14 to 39) (30 to 52) (28 to 66) (293 to 522) (265 to 598)	Age-Mean 93.3 58.1 63.8 48.6 77.4 53.8 1967 1227	standardized 95% UI (70.3 to 126.1) (30.1 to 96.4) (48.6 to 96.0) (30.0 to 82.0) (61.7 to 105.6) (35.6 to 83.9) (1434.3 to 2515.0) (766.5 to 1659.9)

				2013	756	(482 to 994)	931.1	(594.8 to 1220.1)
		Hemorrhagic						
		Stroke	Male	1990	483	(372 to 637)	2096.9	(1607.0 to 2773.3)
				2013	476	(290 to 704)	1170.7	(695.2 to 1765.9)
		Female	1990	335	(268 to 481)	1189.7	(952.3 to 1690.9)	
			2013	375	(256 to 589)	804.3	(550.2 to 1268.2)	
			Both	1990	819	(671 to 1,061)	1618.6	(1321.1 to 2106.3)
				2013	852	(607 to 1,188)	987.5	(694.6 to 1403.0)
	YLDs	Ischemic Stroke	Male	1990	5	(3 to 8)	29.7	(19.5 to 41.8)
				2013	10	(6 to 14)	29.7	(19.9 to 41.5)
			Female	1990	6	(4 to 8)	22.2	(14.4 to 30.9)
				2013	10	(6 to 14)	23	(15.1 to 32.9)
			Both	1990	12	(8 to 16)	25.3	(17.0 to 34.6)
				2013	20	(14 to 28)	25.8	(17.8 to 35.4)
		Hemorrhagic						
		Stroke	Male	1990	6	(4 to 9)	32	(21.0 to 44.4)
				2013	11	(7 to 16)	31.6	(20.7 to 44.4)
			Female	1990	6	(3 to 8)	21.5	(14.2 to 30.3)
				2013	10	(6 to 13)	22	(14.6 to 30.3)
			Both	1990	12	(8 to 17)	26.1	(17.7 to 36.1)
			Both	2013	21	(14 to 29)	26.2	(17.9 to 35.4)
Sierra Leone	Deaths	Ischemic Stroke	Male	1990	460	(297 to 623)	78.4	(53.1 to 103.6)
				2013	853	(358 to 1,218)	117.4	(54.1 to 163.4)
			Female	1990	361	(206 to 484)	70.2	(42.1 to 94.4)
				2013	692	(251 to 991)	103.8	(40.8 to 145.7)
			Both	1990	822	(568 to 1,024)	74.2	(54.4 to 92.3)
				2013	1,546	(732 to 2,134)	110.3	(57.1 to 146.6)
		Hemorrhagic				-		
	Deaths	Stroke	Male	1990	598	(453 to 807)	72.6	(53.9 to 99.5)

				2013	736	(432 to 1,243)	70	(37.1 to 132.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	473	(335 to 670)	63.9	(43.3 to 97.0)
				2013	627	(416 to 1,144)	62.1	(37.2 to 127.5)
			Both	1990	1,071	(862 to 1,378)	68.2	(54.5 to 90.4)
				2013	1,364	(907 to 2,201)	65.9	(39.5 to 121.9)
	DALYs	Ischemic Stroke	Male	1990	11,341	(6,984 to 15,926)	1331.5	(878.4 to 1769.6)
				2013	19,579	(7,984 to 28,216)	1922.9	(839.1 to 2690.8)
			Female	1990	8,440	(4,381 to 11,620)	1061.2	(622.7 to 1408.6)
				2013	14,614	(4,778 to 21,388)	1506.3	(568.5 to 2130.9)
			Both	1990	19,782	(13,552 to 25,398)	1195.4	(833.8 to 1486.7)
				2013	34,193	(15,490 to 47,755)	1706.1	(842.4 to 2327.2)
		Hemorrhagic						
		Stroke	Male	1990	21,220	(15,994 to 28,049)	1658.1	(1254.9 to 2239.7)
				2013	24,481	(15,529 to 37,546)	1550.3	(898.9 to 2658.6)
			Female	1990	16,004	(10,742 to 21,837)	1313.6	(919.8 to 1877.7)
				2013	19,463	(13,356 to 33,047)	1260.8	(837.3 to 2297.1)
			Both	1990	37,224	(29,621 to 46,893)	1484.5	(1187.6 to 1913.4)
				2013	43,945	(31,464 to 64,272)	1401	(920.7 to 2291.0)
	YLDs	Ischemic Stroke	Male	1990	145	(87 to 211)	15.4	(9.5 to 22.1)
				2013	202	(128 to 293)	15.6	(9.9 to 22.2)
			Female	1990	113	(73 to 165)	11.8	(7.7 to 17.4)
				2013	157	(97 to 235)	11.2	(6.8 to 16.6)
			Both	1990	258	(172 to 362)	13.6	(9.0 to 19.0)
				2013	360	(239 to 503)	13.3	(8.8 to 18.7)
		Hemorrhagic	Nala	1000	46	(20 to CO)	4.2	(2 C to C 2)
		Stroke	Male	1990	46	(28 to 69)	4.2	(2.6 to 6.3)

				2013	62	(38 to 93)	3.9	(2.5 to 5.8)
			Female	1990	33	(20 to 49)	3	(1.9 to 4.5)
				2013	45	(27 to 67)	2.7	(1.6 to 3.9)
			Both	1990	79	(51 to 113)	3.6	(2.4 to 5.1)
			Both	2013	108	(69 to 154)	3.3	(2.1 to 4.7)
Singapore	Deaths	Ischemic Stroke	Male	1990	373	(293 to 450)	56.1	(44.2 to 67.6)
				2013	589	(459 to 792)	30.7	(24.0 to 41.3)
			Female	1990	523	(374 to 604)	59.5	(42.4 to 68.6)
				2013	897	(648 to 1,317)	30.4	(22.3 to 44.7)
			Both	1990	897	(732 to 1,005)	58.6	(47.6 to 65.6)
				2013	1,487	(1,227 to 1,984)	30.9	(25.5 to 41.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	351	(284 to 435)	42.4	(33.8 to 52.8)
				2013	454	(314 to 572)	22.1	(15.2 to 28.0)
			Female	1990	431	(323 to 518)	45.8	(33.9 to 55.0)
				2013	551	(366 to 783)	19.7	(13.4 to 28.0)
			Both	1990	783	(667 to 899)	44.7	(37.7 to 52.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	1,006	(745 to 1,226)	21.1	(15.8 to 25.7)
	DALYs	Ischemic Stroke	Male	1990	6,868	(5,499 to 8,154)	864.6	(689.1 to 1024.4)
				2013	8,790	(7,158 to 11,148)	435	(354.4 to 551.9)
			Female	1990	7,179	(5,135 to 8,272)	766.9	(552.2 to 882.5)
				2013	8,713	(6,680 to 12,483)	327.3	(250.3 to 466.7)
			Both	1990	14,047	(11,732 to 15,739)	817.3	(678.4 to 911.5)
				2013	17,504	(14,830 to 22,213)	378.6	(319.9 to 479.6)
		Hemorrhagic						
		Stroke	Male	1990	8,873	(7,334 to 10,704)	890.7	(735.5 to 1086.2)

				2013	8,990	(6,943 to 11,040)	404.2	(313.5 to 493.4)
			Female	1990	8,696	(6,633 to 10,189)	843.2	(641.4 to 990.4)
				2013	7,418	(5,521 to 10,446)	291	(217.9 to 410.1)
			Both	1990	17,570	(15,243 to 19,773)	870.4	(752.7 to 982.2)
				2013	16,409	(13,399 to 19,785)	346.3	(284.0 to 414.2)
	YLDs	Ischemic Stroke	Male	1990	819	(524 to 1,166)	97.1	(62.8 to 137.5)
				2013	1,898	(1,262 to 2,622)	90	(60.4 to 123.7)
			Female	1990	561	(350 to 823)	57	(35.5 to 83.7)
				2013	1,371	(859 to 1,988)	53.8	(33.5 to 77.7)
			Both	1990	1,381	(926 to 1,868)	75.2	(50.4 to 101.8)
				2013	3,270	(2,184 to 4,514)	70.5	(46.7 to 97.1)
		Hemorrhagic						
		Stroke	Male	1990	311	(201 to 448)	33.1	(21.4 to 48.3)
				2013	704	(452 to 1,022)	32.7	(21.2 to 46.8)
			Female	1990	229	(142 to 347)	21.2	(13.2 to 32.5)
				2013	591	(371 to 883)	23.5	(14.7 to 34.9)
			Both	1990	540	(357 to 746)	26.5	(17.4 to 37.1)
			Both	2013	1,296	(839 to 1,818)	27.7	(18.1 to 39.0)
Slovakia	Deaths	Ischemic Stroke	Male	1990	3,006	(2,535 to 3,550)	158.8	(133.3 to 188.1)
				2013	2,530	(2,118 to 3,100)	100.6	(84.0 to 122.5)
			Female	1990	3,448	(2,737 to 3,992)	111.5	(88.5 to 129.6)
				2013	3,230	(2,742 to 4,080)	66.5	(56.6 to 83.4)
			Both	1990	6,454	(5,740 to 7,279)	130.4	(115.5 to 147.0)
				2013	5,760	(5,095 to 6,823)	79.3	(70.1 to 93.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	917	(643 to 1,161)	41.7	(29.1 to 52.9)
				2013	686	(469 to 862)	23.5	(16.0 to 29.8)
			Female	1990	1,079	(695 to 1,271)	35	(22.5 to 41.4)
				2013	709	(522 to 994)	15.8	(11.7 to 21.8)

			Both	1990	1,997	(1,481 to 2,346)	38.2	(28.1 to 44.9)
				2013	1,396	(1,120 to 1,704)	19.1	(15.4 to 23.4)
	DALYs	Ischemic Stroke	Male	1990	46,385	(39,440 to 55,071)	2237.5	(1900.4 to 2637.4)
				2013	37,112	(31,083 to 45,690)	1337.5	(1122.6 to 1635.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	41,801	(33,798 to 48,518)	1354	(1090.7 to 1570.2)
				2013	33,956	(29,070 to 41,939)	740.2	(635.8 to 909.5)
			Both	1990	88,187	(79,276 to 99,689)	1723.4	(1548.7 to 1939.1)
				2013	71,069	(63,031 to 83,249)	982.1	(872.0 to 1151.9)
		Hemorrhagic						
		Stroke	Male	1990	21,797	(15,803 to 27,535)	918.4	(661.0 to 1159.9)
				2013	15,441	(10,706 to 19,298)	484.5	(338.4 to 609.4)
			Female	1990	20,237	(13,768 to 23,412)	669.7	(457.9 to 777.0)
				2013	11,728	(9,059 to 15,699)	284.4	(222.3 to 374.9)
			Both	1990	42,035	(32,127 to 48,984)	785.6	(601.8 to 917.1)
				2013	27,170	(22,281 to 32,464)	375.5	(306.9 to 448.1)
	YLDs	Ischemic Stroke	Male	1990	547	(360 to 783)	23.7	(15.6 to 33.7)
				2013	784	(507 to 1,114)	24.8	(16.0 to 35.2)
			Female	1990	460	(298 to 655)	15.4	(10.0 to 21.9)
				2013	597	(373 to 866)	14.7	(9.3 to 21.3)
			Both	1990	1,008	(668 to 1,383)	19	(12.6 to 26.1)
				2013	1,381	(936 to 1,891)	19.2	(13.0 to 26.4)
		Hemorrhagic						
		Stroke	Male	1990	122	(76 to 181)	5.1	(3.2 to 7.5)
				2013	176	(109 to 260)	5.6	(3.5 to 8.2)
			Female	1990	109	(67 to 161)	3.8	(2.4 to 5.5)
				2013	146	(90 to 222)	3.9	(2.4 to 5.8)

			Both	1990	232	(153 to 326)	4.4	(2.9 to 6.1)
			Both	2013	323	(207 to 455)	4.6	(3.0 to 6.5)
Slovenia	Deaths	Ischemic Stroke	Male	1990	920	(774 to 1,071)	128.1	(106.9 to 149.8)
				2013	779	(610 to 942)	56.5	(43.9 to 68.3)
			Female	1990	1,349	(1,041 to 1,487)	95.3	(73.5 to 105.2)
				2013	1,207	(929 to 1,528)	43.7	(34.0 to 55.2)
			Both	1990	2,269	(1,931 to 2,478)	107.4	(91.3 to 117.2)
				2013	1,987	(1,654 to 2,310)	49	(40.8 to 56.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	275	(224 to 371)	31	(25.0 to 42.7)
				2013	249	(165 to 317)	17.4	(11.5 to 22.2)
			Female	1990	329	(257 to 438)	24.2	(18.9 to 32.1)
				2013	305	(202 to 406)	13.4	(9.4 to 17.3)
			Both	1990	605	(530 to 762)	27.2	(23.8 to 34.6)
				2013	554	(407 to 673)	15.3	(11.5 to 18.4)
	DALYs	Ischemic Stroke	Male	1990	13,601	(11,447 to 15,714)	1725	(1460.8 to 1996.1)
				2013	9,329	(7,394 to 11,136)	649.6	(511.4 to 776.0)
			Female	1990	15,041	(11,412 to 16,496)	1090.5	(824.1 to 1196.0)
				2013	9,796	(7,693 to 12,221)	398.2	(313.9 to 503.6)
			Both	1990	28,643	(24,386 to 31,325)	1335.4	(1135.0 to 1458.7)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	19,125	(16,159 to 21,934)	505.7	(428.3 to 581.1)
		Hemorrhagic						/
		Stroke	Male	1990	7,108	(5,849 to 9,141)	725.7	(596.4 to 938.3)
				2013	4,861	(3,410 to 6,114)	336.1	(236.7 to 422.2)
			Female	1990	6,644	(5,210 to 8,415)	514.4	(402.5 to 645.4)
				2013	4,073	(3,018 to 5,225)	225.3	(172.6 to 288.0)

		I	1		1	T .		
			Both	1990	13,753	(12,159 to 16,614)	611.2	(539.7 to 740.7)
				2013	8,934	(7,116 to 10,480)	278.5	(222.8 to 326.3)
	YLDs	Ischemic Stroke	Male	1990	226	(142 to 323)	24	(15.2 to 34.2)
			2013	343	(213 to 517)	23.7	(14.7 to 35.3)	
			Female	1990	209	(136 to 297)	16.2	(10.5 to 23.0)
	Hemorrhagic		2013	255	(164 to 362)	14.5	(9.4 to 20.7)	
		Both	1990	436	(291 to 603)	19.6	(13.1 to 27.1)	
			2013	599	(400 to 827)	18.9	(12.6 to 26.1)	
		Stroke	Male	1990	50	(29 to 77)	5.1	(3.0 to 7.8)
				2013	78	(45 to 120)	5.7	(3.3 to 8.6)
			Female	1990	47	(27 to 72)	3.9	(2.2 to 5.9)
				2013	61	(36 to 93)	4	(2.4 to 5.9)
			Both	1990	98	(62 to 142)	4.4	(2.8 to 6.4)
			Both	2013	139	(89 to 200)	4.8	(3.0 to 6.8)
Solomon	Deaths	Ischemic Stroke	Male	1990	38	(16 to 58)	77.9	(35.1 to 113.8)
Islands				2013	73	(32 to 117)	77.2	(35.5 to 118.3)
			Female	1990	20	(11 to 31)	51.3	(30.4 to 79.1)
				2013	56	(24 to 86)	62.7	(30.4 to 94.9)
			Both	1990	58	(29 to 84)	66.1	(34.4 to 91.4)
				2013	130	(62 to 195)	70.1	(35.6 to 101.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	82	(51 to 120)	134	(86.3 to 190.3)
				2013	170	(108 to 259)	140.2	(92.1 to 205.9)
			Female	1990	45	(27 to 72)	90.8	(56.0 to 146.1)
				2013	118	(68 to 188)	108.7	(63.8 to 164.5)
			Both	1990	128	(83 to 188)	113.9	(76.6 to 158.1)
				2013	289	(184 to 434)	124.6	(81.1 to 180.9)
	DALYs	Ischemic Stroke	Male	1990	858	(353 to 1,371)	1392	(611.7 to 2128.4)

				2013	1,599	(684 to 2,699)	1316.2	(596.9 to 2106.0)
			Female	1990	468	(229 to 758)	884.1	(488.5 to 1342.4)
				2013	1,165	(462 to 1,917)	1017.3	(451.5 to 1563.3)
			Both	1990	1,327	(625 to 2,077)	1158.9	(584.2 to 1689.7)
				2013	2,764	(1,258 to 4,458)	1169	(568.3 to 1776.6)
		Hemorrhagic						
		Stroke	Male	1990	2,512	(1,519 to 3,794)	3192.8	(1983.4 to 4693.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	5,103	(3,118 to 8,053)	3258.2	(2051.4 to 4974.3)
			Female	1990	1,330	(813 to 2,123)	1971	(1187.2 to 3104.8)
				2013	3,210	(1,823 to 5,247)	2265	(1300.7 to 3619.1)
			Both	1990	3,843	(2,439 to 5,854)	2607.2	(1692.3 to 3850.3)
				2013	8,314	(5,061 to 12,855)	2765.6	(1752.0 to 4174.2)
	YLDs	Ischemic Stroke	Male	1990	15	(9 to 22)	27.3	(17.2 to 40.6)
				2013	28	(19 to 39)	25.3	(17.0 to 35.2)
			Female	1990	10	(6 to 14)	21.3	(13.5 to 30.4)
				2013	22	(14 to 33)	21.1	(13.4 to 30.9)
			Both	1990	25	(16 to 35)	24.6	(16.4 to 34.3)
				2013	51	(35 to 68)	23.3	(16.0 to 31.2)
		Hemorrhagic						
		Stroke	Male	1990	20	(13 to 28)	32.2	(20.9 to 45.2)
				2013	38	(25 to 52)	30	(20.3 to 41.1)
			Female	1990	12	(8 to 18)	22.7	(14.4 to 32.2)
				2013	27	(18 to 38)	22.5	(15.1 to 31.7)
			Both	1990	32	(22 to 45)	27.8	(18.9 to 37.9)
			1				1	
			Both	2013	65	(45 to 88)	26.3	(18.2 to 35.7)

				2013	790	(232 to 1,441)	63	(19.3 to 110.4)
			Female	1990	870	(326 to 1,444)	89.6	(34.4 to 143.9)
				2013	986	(413 to 1,544)	68.6	(29.2 to 106.5)
			Both	1990	1,484	(646 to 2,494)	82.5	(35.9 to 133.7)
				2013	1,776	(771 to 2,833)	66.3	(29.7 to 103.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	942	(494 to 1,596)	85.1	(40.9 to 148.3)
				2013	1,178	(610 to 2,027)	73.7	(35.0 to 131.0)
			Female	1990	1,343	(675 to 2,136)	110.4	(57.0 to 173.8)
				2013	1,475	(799 to 2,385)	84.1	(44.6 to 139.7)
			Both	1990	2,286	(1,290 to 3,579)	99.1	(56.2 to 153.6)
				2013	2,653	(1,502 to 4,202)	79.6	(44.4 to 127.8)
	DALYs	Ischemic Stroke	Male	1990	12,351	(4,080 to 23,939)	1113.5	(368.4 to 2126.0)
				2013	15,220	(4,412 to 28,810)	962.3	(285.1 to 1752.6)
			Female	1990	15,352	(5,412 to 26,512)	1288.2	(479.5 to 2157.9)
				2013	16,648	(6,720 to 26,883)	970.6	(407.0 to 1527.8)
			Both	1990	27,703	(11,729 to 48,309)	1213.8	(529.9 to 2046.3)
				2013	31,868	(13,460 to 52,013)	970.3	(421.9 to 1550.6)
		Hemorrhagic						
		Stroke	Male	1990	26,211	(14,625 to 42,692)	1760.9	(923.1 to 2981.3)
				2013	30,856	(17,173 to 51,939)	1480.6	(765.1 to 2545.7)
			Female	1990	33,348	(15,930 to 52,848)	2171.3	(1068.3 to 3468.7)
				2013	34,118	(18,344 to 54,334)	1577.6	(847.0 to 2546.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	59,560	(33,817 to 91,766)	1984	(1114.1 to 3133.3)
						(36,663 to		
				2013	64,975	103,434)	1537	(858.9 to 2445.1)

	YLDs	Ischemic Stroke	Male	1990	153	(95 to 222)	10.4	(6.6 to 14.8)
				2013	226	(141 to 334)	10.3	(6.6 to 15.4)
			Female	1990	109	(70 to 163)	6.9	(4.3 to 10.2)
				2013	188	(119 to 278)	7.9	(4.9 to 11.7)
			Both	1990	262	(173 to 366)	8.6	(5.6 to 11.8)
				2013	415	(274 to 591)	9	(6.0 to 12.8)
		Hemorrhagic						
		Stroke	Male	1990	51	(30 to 75)	2.9	(1.8 to 4.2)
				2013	81	(49 to 124)	3	(1.8 to 4.5)
			Female	1990	31	(19 to 48)	1.8	(1.1 to 2.7)
				2013	54	(33 to 83)	1.9	(1.1 to 2.9)
			Both	1990	83	(54 to 117)	2.3	(1.5 to 3.2)
			Both	2013	136	(86 to 193)	2.4	(1.5 to 3.5)
South	Deaths	Ischemic Stroke	Male	1990	4,008	(3,108 to 5,245)	61.7	(48.0 to 79.8)
Africa				2013	5,887	(4,352 to 7,433)	52.9	(39.6 to 66.8)
			Female	1990	5,705	(4,382 to 7,046)	61.8	(47.7 to 76.2)
				2013	11,709	(8,384 to 14,884)	57.1	(40.8 to 72.7)
			Both	1990	9,714	(8,234 to 11,479)	62.4	(53.0 to 73.3)
				2013	17,596	(13,778 to 20,932)	56.5	(44.1 to 67.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,923	(3,785 to 6,373)	62.4	(48.0 to 81.6)
				2013	6,592	(4,994 to 8,518)	52.6	(39.9 to 68.1)
			Female	1990	6,722	(4,900 to 8,131)	66.7	(49.3 to 81.8)
				2013	10,606	(8,054 to 14,113)	51.4	(39.0 to 68.2)
			Both	1990	11,645	(9,411 to 13,602)	65.4	(54.0 to 77.4)
				2013	17,198	(14,369 to 20,979)	52.6	(43.9 to 64.0)
						(59,457 to		
	DALYs	Ischemic Stroke	Male	1990	79,916	106,977)	991.1	(761.0 to 1296.0)
				2013	103,722	(74,943 to	796.8	(587.3 to 999.3)

					132,785)		
					(64,005 to		
		Female	1990	89,083	114,556)	877.3	(653.5 to 1091.3)
					(108,790 to		
			2013	148,093	192,189)	730.9	(535.2 to 946.9)
					(137,131 to		
		Both	1990	168,999	209,680)	931.9	(773.5 to 1118.2)
					(199,715 to		
			2013	251,815	300,608)	767.2	(609.2 to 915.2)
	Hemorrhagic				(101,872 to		
	Stroke	Male	1990	134,423	172,904)	1296.9	(995.9 to 1682.0)
				Counts, All	Ages	Rate	Per 100,000
						Age-	standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
					(110,994 to		
			2013	145,969	188,637)	950.5	(721.3 to 1227.5)
					(94,333 to		
		Female	1990	148,448	185,280)	1274.6	(859.8 to 1554.1)
					(127,111 to		
			2013	166,631	226,703)	794.9	(606.4 to 1064.1)
					(213,738 to		
		Both	1990	282,871	340,087)	1288.5	(1013.4 to 1516.5)
					(259,044 to		
			2013	312,600	378,103)	868.7	(723.3 to 1054.5)
YLDs	Ischemic Stroke	Male	1990	1,774	(1,167 to 2,498)	18.5	(11.8 to 26.1)
			2013	3,229	(2,116 to 4,575)	20.1	(13.1 to 28.5)
		Female	1990	2,064	(1,364 to 2,953)	18.2	(12.0 to 26.1)
				· · · · · · · · · · · · · · · · · · ·	<u> </u>		·
			2013	3,896	(2,496 to 5,565)	17.8	(11.2 to 25.6)
		Both	2013 1990	3,896 3,839	(2,496 to 5,565) (2,592 to 5,247)	17.8 18.3	(11.2 to 25.6) (12.4 to 25.3)
		measure Stroke Type	Male Hemorrhagic Stroke Male Male Female Both YLDs Ischemic Stroke Male	Both 1990 2013	2013 148,093 168,999 2013 251,815	Female 1990 89,083 114,556 (108,790 to 108,790 to 120,189 (137,131 to 2013 251,815 300,608 (109,715 to 300,608 (101,872 to 134,423 172,904 (101,974 to 2013 145,969 188,637 (127,111 to 2013 166,631 226,703 (213,738 to 378,103 YLDs Ischemic Stroke Male 1990 1,774 (1,167 to 2,498) (1,167 to 2,498) (2013 3,229 (2,116 to 4,575)	Female 1990 89,083 114,556 877.3 (108,790 to 108,790 to 108,790 to 192,189 730.9 (137,131 to 2013 251,815 300,608 767.2 (101,872 to Stroke Male 1990 134,423 172,904 1296.9 (101,9715 to 300,608 767.2 (101,872 to 172,904 1296.9 (101,9715 to 300,608 767.2 (101,872 to 172,904 1296.9 (101,9715 to 188,637 950.5 (101,994 to 188,637 950.5

		Hemorrhagic						
		Stroke	Male	1990	539	(336 to 783)	4.8	(3.0 to 7.0)
				2013	958	(624 to 1,431)	5.3	(3.4 to 7.8)
			Female	1990	545	(345 to 781)	4.4	(2.8 to 6.3)
				2013	952	(597 to 1,406)	4.1	(2.6 to 6.2)
			Both	1990	1,084	(725 to 1,504)	4.6	(3.1 to 6.4)
			Both	2013	1,911	(1,300 to 2,693)	4.6	(3.1 to 6.5)
South	Deaths	Ischemic Stroke	Male	1990	10,153	(8,441 to 12,708)	146.1	(120.3 to 184.7)
Korea				2013	12,536	(9,431 to 15,166)	60.5	(45.4 to 73.4)
			Female	1990	11,511	(9,133 to 14,168)	96.2	(77.4 to 118.0)
				2013	17,986	(13,946 to 23,126)	47.9	(37.1 to 61.8)
			Both	1990	21,665	(19,497 to 25,876)	114.2	(102.5 to 136.4)
				2013	30,523	(24,705 to 35,731)	53.2	(43.1 to 62.2)
		Hemorrhagic				, , ,		· ·
	Deaths	Stroke	Male	1990	12,674	(10,148 to 14,993)	135.1	(101.4 to 160.6)
				2013	9,015	(7,175 to 11,787)	37.2	(28.9 to 49.9)
			Female	1990	14,705	(10,445 to 16,484)	109.4	(77.6 to 123.9)
				2013	10,823	(8,095 to 17,287)	29.7	(22.3 to 47.2)
			Both	1990	27,380	(22,172 to 30,450)	120.4	(94.5 to 135.5)
				2013	19,839	(16,072 to 27,053)	33.4	(26.9 to 46.1)
						(169,165 to		
	DALYs	Ischemic Stroke	Male	1990	201,716	252,135)	2205	(1843.6 to 2734.6)
						(146,845 to		
				2013	189,485	228,305)	784.2	(612.7 to 939.4)
			Female	1990	173,681	(133,257 to 215,390)	1280.2	(1006.1 to 1577.7)
			· cinaic	1000	1,0,001	(140,061 to		(1000.1 to 10.7.7)
				2013	178,914	229,721)	488.4	(381.8 to 629.9)
					Counts All	•		Per 100 000

Counts, All Ages

Rate Per 100,000 Age- standardized

Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(329,239 to		
			Both	1990	375,397	446,576)	1636.6	(1461.0 to 1941.3)
						(299,275 to		
				2013	368,400	426,518)	614.9	(499.5 to 710.9)
		Hemorrhagic				(283,382 to		
		Stroke	Male	1990	339,447	415,491)	2644.6	(2114.6 to 3124.2)
						(157,661 to		
				2013	192,666	235,196)	685.2	(559.2 to 844.3)
						(219,259 to		
			Female	1990	301,182	332,145)	1917.3	(1409.8 to 2123.3)
						(117,865 to		
				2013	149,553	229,783)	430.6	(340.9 to 662.0)
				1000		(543,898 to		(1010 = 1 01=00)
_			Both	1990	640,630	707,692)	2238.3	(1848.7 to 2479.9)
				2042	242 240	(294,211 to	554 C	(472.61607.4)
				2013	342,219	431,821)	551.6	(473.6 to 697.1)
	YLDs	Ischemic Stroke	Male	1990	9,754	(6,375 to 13,628)	94	(60.6 to 131.9)
				2013	29,297	(19,675 to 40,648)	109	(72.3 to 151.9)
			Female	1990	8,666	(5,647 to 11,937)	59.5	(38.6 to 82.0)
				2013	23,246	(15,768 to 32,594)	66.1	(44.8 to 92.6)
			Both	1990	18,420	(12,523 to 24,786)	73.1	(49.4 to 98.7)
				2013	52,543	(36,002 to 72,169)	85.1	(58.2 to 116.9)
		Hemorrhagic						,
		Stroke	Male	1990	3,505	(2,167 to 5,053)	28.8	(18.1 to 42.7)
				2013	10,034	(6,464 to 14,360)	36.6	(23.4 to 51.8)
			Female	1990	3,329	(2,151 to 4,819)	20.5	(13.2 to 29.9)
				2013	9,228	(5,877 to 12,960)	27	(17.1 to 37.9)
			Both	1990	6,835	(4,489 to 9,632)	23.7	(15.6 to 33.6)
			Both	2013	19,263	(12,629 to 26,640)	31.1	(20.5 to 43.0)
			DUIII	2013	15,203	(12,023 (0 20,040)	31.1	(20.3 to 43.0)

South	Deaths	Ischemic Stroke	Male	1990	665	(284 to 1,086)	94.9	(41.9 to 150.8)
Sudan				2013	1,489	(807 to 2,125)	90.6	(49.3 to 127.8)
			Female	1990	546	(279 to 859)	78.6	(41.1 to 118.4)
				2013	1,251	(793 to 1,796)	70.5	(44.3 to 99.5)
			Both	1990	1,211	(666 to 1,774)	86.4	(48.8 to 122.4)
				2013	2,741	(1,835 to 3,551)	79.9	(54.1 to 102.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	988	(601 to 1,506)	99.6	(59.6 to 152.6)
				2013	1,857	(1,224 to 2,603)	90.2	(56.5 to 130.0)
			Female	1990	772	(445 to 1,201)	80.4	(48.3 to 121.8)
				2013	1,439	(985 to 2,091)	67.5	(44.2 to 98.7)
			Both	1990	1,760	(1,212 to 2,439)	89.8	(61.3 to 123.8)
				2013	3,296	(2,487 to 4,354)	78.3	(56.9 to 107.0)
	DALYs	Ischemic Stroke	Male	1990	14,616	(6,484 to 23,757)	1505.3	(656.2 to 2432.0)
				2013	28,776	(15,146 to 41,157)	1413.1	(762.5 to 2021.8)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	9,981	(4,958 to 15,974)	1063.2	(547.6 to 1665.8)
				2013	20,085	(12,681 to 29,211)	943.3	(605.3 to 1361.7)
			Both	1990	24,597	(13,288 to 36,794)	1274.3	(710.4 to 1847.2)
				2013	48,861	(31,398 to 64,243)	1165.6	(774.4 to 1511.7)
		Hemorrhagic						
		Stroke	Male	1990	29,155	(18,306 to 43,631)	2132.5	(1287.6 to 3242.0)
				2013	48,766	(32,984 to 66,869)	1867.2	(1234.3 to 2606.1)
			Female	1990	19,935	(11,392 to 31,120)	1490	(853.6 to 2317.7)
				2013	32,185	(22,824 to 46,014)	1215.3	(845.2 to 1759.7)
			Both	1990	49,090	(34,089 to 67,878)	1801.7	(1237.4 to 2497.9)
				2013	80,951	(63,153 to	1529.4	(1164.2 to 2003.9)

					104 161)		
\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \			4000	400		0 -	(6.4.1.5.7)
YLDs	Ischemic Stroke	Male			` '		(6.1 to 14.5)
					· · · · · ·	10.1	(6.3 to 14.7)
		Female	1990	98	(60 to 144)	7	(4.3 to 10.1)
			2013	214	(134 to 308)	7.3	(4.7 to 10.6)
		Both	1990	226	(149 to 331)	8.2	(5.6 to 11.8)
			2013	488	(321 to 687)	8.7	(5.9 to 12.0)
	Hemorrhagic						
	Stroke	Male	1990	42	(27 to 63)	2.7	(1.7 to 3.9)
			2013	91	(53 to 138)	2.8	(1.6 to 4.2)
		Female	1990	29	(17 to 42)	1.8	(1.1 to 2.7)
			2013	62	(39 to 93)	1.8	(1.2 to 2.7)
		Both	1990	72	(47 to 103)	2.2	(1.5 to 3.2)
		Both	2013	154	(97 to 222)	2.3	(1.5 to 3.3)
Deaths	Ischemic Stroke	Male	1990	11,478	(9,220 to 13,527)	58.7	(47.1 to 69.3)
			2013	9,441	(7,226 to 12,394)	23.7	(18.3 to 31.2)
		Female	1990	18,957	(13,761 to 21,415)	56.5	(41.1 to 63.8)
			2013	15,735	(12,030 to 24,658)	23.5	(18.2 to 36.9)
		Both	1990	30,435	(24,421 to 33,959)	58	(46.5 to 64.7)
			2013	25,176	(20,607 to 34,968)	23.9	(19.6 to 33.0)
	Hemorrhagic						
Deaths	Stroke	Male	1990	6,684	(5,364 to 8,482)	32.9	(26.3 to 42.1)
			2013	7,879	(4,916 to 9,895)	21.1	(13.4 to 26.2)
		Female	1990	8,682	(6,639 to 11,239)	27.9	(21.4 to 35.6)
			2013	9,801	(6,747 to 13,759)	16.7	(11.9 to 23.0)
		Both	1990	15,366	(13,144 to 18,923)	30.4	(26.1 to 37.1)
		2013	17,681	(12,948 to 21,555)	18.7	(14.1 to 22.5)	
DALYs	Ischemic Stroke	Male	1990	160.713	(132,186 to 185.872)	780.2	(645.7 to 907.1)
		Hemorrhagic Stroke Deaths Ischemic Stroke Hemorrhagic Stroke Stroke	Female Both Hemorrhagic Stroke Female Both Both Both Both Deaths Ischemic Stroke Male Female Female Female Female Female Female Both Female Female Both Female Both Female Both Both Female Both Both Both Female Both Both Both Both Both		Female	Comparison	YLDs Ischemic Stroke Male 1990 128 (80 to 198) 9.5 2013 274 (170 to 402) 10.1 Female 1990 98 (60 to 144) 7 2013 214 (134 to 308) 7.3 Both 1990 226 (149 to 331) 8.2 Hemorrhagic Male 1990 42 (27 to 63) 2.7 Stroke Male 1990 42 (27 to 63) 2.7 Pemale 1990 42 (27 to 63) 2.7 Stroke Male 1990 42 (27 to 63) 2.7 1 2013 91 (53 to 138) 2.8 2013 62 (39 to 93) 1.8 2013 62 (39 to 93) 1.8 Both 1990 72 (47 to 103) 2.2 2013 154 (97 to 222) 2.3 Deaths Ischemic Stroke Male 1990 11,478 (9,220 to 13

						(97,989 to		
				2013	119,825	149,897)	331	(270.9 to 409.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(151,167 to		
			Female	1990	198,301	220,096)	626.3	(479.3 to 695.3)
						(110,256 to		
				2013	139,354	204,269)	253.8	(200.0 to 367.0)
						(301,246 to		
			Both	1990	359,015	397,322)	696.4	(585.5 to 771.1)
						(217,480 to		
				2013	259,179	336,618)	289.7	(243.4 to 372.2)
		Hemorrhagic				(109,000 to		
		Stroke	Male	1990	132,066	162,742)	639.1	(528.1 to 787.1)
						(80,853 to		
				2013	115,848	140,244)	347.4	(245.0 to 417.7)
						(97,350 to		
			Female	1990	123,727	153,511)	460.2	(364.7 to 569.5)
						(77,458 to		
				2013	102,972	136,426)	233.9	(182.3 to 305.5)
						(228,086 to		
			Both	1990	255,794	301,160)	544.9	(486.0 to 636.5)
						(173,863 to		(22.1.1.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.
				2013	218,820	252,414)	286.8	(231.1 to 329.0)
	YLDs	Ischemic Stroke	Male	1990	17,132	(11,547 to 23,353)	80.3	(54.2 to 109.7)
				2013	27,805	(18,938 to 39,599)	82.4	(56.0 to 117.6)
			Female	1990	20,339	(13,965 to 27,823)	69.8	(47.9 to 95.3)
				2013	26,109	(17,045 to 36,201)	60.4	(39.9 to 83.4)
			Both	1990	37,472	(25,970 to 49,546)	74.7	(51.8 to 98.8)

				,			
			2013	53,915	(36,733 to 74,274)	70.4	(47.9 to 97.3)
	Hemorrhagic						
	Stroke	Male	1990	3,350	(1,951 to 5,160)	16	(9.4 to 24.6)
			2013	6,428	(4,018 to 9,047)	19.5	(12.3 to 27.5)
		Female	1990	4,179	(2,470 to 6,226)	15.5	(9.4 to 23.2)
			2013	5,466	(3,123 to 8,524)	14.4	(8.6 to 21.6)
		Both	1990	7,530	(4,735 to 10,765)	15.6	(9.8 to 22.1)
		Both	2013	11,895	(7,676 to 16,394)	16.6	(10.8 to 22.8)
Deaths	Ischemic Stroke	Male	1990	2,296	(1,882 to 2,855)	49.3	(40.1 to 62.4)
			2013	4,397	(2,692 to 5,862)	58.8	(36.0 to 78.3)
		Female	1990	1,610	(1,223 to 1,984)	37.5	(28.4 to 46.0)
			2013	4,646	(1,513 to 6,615)	53	(17.0 to 75.7)
		Both	1990	3,906	(3,402 to 4,598)	43.4	(37.4 to 50.9)
			2013	9,043	(4,657 to 11,762)	55.8	(28.2 to 73.1)
	Hemorrhagic						
Deaths	Stroke	Male	1990	1,577	(1,281 to 2,092)	28.8	(23.1 to 39.0)
			2013	1,854	(1,240 to 3,351)	20.9	(13.6 to 40.2)
		Female	1990	1,026	(842 to 1,432)	20.2	(16.1 to 29.1)
			2013	1,440	(856 to 2,414)	14.6	(8.6 to 25.2)
				Counts, All	Ages	Rate	Per 100,000
						Age-	standardized
measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Both	1990	2,603	(2,252 to 3,327)	24.5	(20.9 to 31.9)
			2013	3,294	(2,407 to 5,500)	17.6	(12.5 to 30.6)
DALYs	Ischemic Stroke	Male	1990	47,726	(39,423 to 57,233)	860.1	(711.8 to 1038.1)
			2013	76,038	(48,047 to 99,657)	884.9	(558.1 to 1165.8)
		Female	1990	28,349	(21,798 to 34,890)	541.3	(419.6 to 660.9)
			2013	60,575	(24,113 to 84,503)	621.3	(240.9 to 867.8)
		Both	1990	76,076	(66,742 to 88,686)	701.5	(617.5 to 821.7)
	Deaths	Deaths Ischemic Stroke Hemorrhagic Stroke Stroke Measure Stroke Type	Stroke Male Female Both Both Deaths Ischemic Stroke Male Female Both Hemorrhagic Stroke Male Female Deaths Stroke Male Female Temale Female Female Female	Hemorrhagic Stroke	Hemorrhagic Stroke	Hemorrhagic Stroke Male 1990 3,350 (1,951 to 5,160)	Hemorrhagic Stroke Male 1990 3,350 (1,951 to 5,160) 16

		T			I	/76 004 :		
				2012	126 614	(76,831 to	742.0	(442.7 + - 052.0)
		Llaura a mula a aila		2013	136,614	175,337)	743.8	(412.7 to 952.9)
		Hemorrhagic Stroke	Male	1990	49,946	(41,255 to 63,477)	739.3	(607.9 to 945.8)
		Stroke	iviale					,
			- 1	2013	50,634	(36,422 to 76,858)	508.9	(362.7 to 806.8)
			Female	1990	30,638	(25,305 to 39,028)	460.2	(389.3 to 610.8)
				2013	32,352	(21,545 to 45,067)	297.1	(199.4 to 422.9)
			Both	1990	80,585	(71,029 to 97,361)	600.4	(526.8 to 735.7)
				2013	82,986	(64,792 to 116,855)	397.3	(310.6 to 582.9)
	YLDs	Ischemic Stroke	Male	1990	1,516	(991 to 2,175)	29.5	(19.3 to 42.1)
				2013	2,562	(1,652 to 3,661)	29.4	(19.0 to 41.6)
			Female	1990	1,249	(817 to 1,785)	24.7	(16.1 to 35.2)
				2013	2,673	(1,679 to 3,840)	26	(16.1 to 37.6)
			Both	1990	2,766	(1,881 to 3,813)	27.1	(18.5 to 37.1)
				2013	5,236	(3,433 to 7,264)	27.6	(18.0 to 38.2)
		Hemorrhagic						
		Stroke	Male	1990	1,719	(1,139 to 2,413)	31.3	(20.6 to 44.1)
				2013	2,874	(1,877 to 4,020)	31.5	(20.5 to 44.2)
			Female	1990	1,289	(828 to 1,810)	23.9	(15.4 to 33.6)
				2013	2,545	(1,643 to 3,540)	24.2	(15.5 to 34.0)
			Both	1990	3,008	(2,024 to 4,093)	27.6	(18.5 to 37.4)
			Both	2013	5,420	(3,641 to 7,384)	27.6	(18.4 to 37.5)
Sudan	Deaths	Ischemic Stroke	Male	1990	2,300	(1,097 to 3,444)	88.9	(43.4 to 123.9)
				2013	4,340	(2,383 to 6,287)	83.4	(47.8 to 114.6)
			Female	1990	2,678	(1,492 to 3,896)	90.5	(51.2 to 127.3)
				2013	4,735	(2,870 to 6,746)	81.1	(49.4 to 112.1)
			Both	1990	4,979	(3,259 to 6,559)	89.8	(57.4 to 113.6)
				2013	9,076	(6,004 to 11,545)	82.2	(54.4 to 102.6)

		Hemorrhagic						
	Deaths	Stroke	Male	1990	3,210	(2,171 to 4,613)	87.4	(59.5 to 128.4)
				2013	4,930	(3,399 to 6,806)	70.7	(49.0 to 98.4)
			Female	1990	4,028	(2,412 to 5,616)	101.2	(63.9 to 146.8)
				2013	5,434	(3,601 to 7,998)	73.9	(48.8 to 107.8)
			Both	1990	7,238	(5,265 to 9,506)	94.8	(69.9 to 126.5)
				2013	10,364	(7,820 to 13,717)	72.5	(55.2 to 96.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
	DALYs	Ischemic Stroke	Male	1990	50,666	(23,358 to 85,424)	1364.3	(645.3 to 2040.9)
						(43,060 to		
				2013	82,432	130,262)	1191.7	(645.4 to 1751.4)
			Female	1990	55,911	(27,353 to 97,305)	1370.9	(767.3 to 2037.7)
						(46,148 to		
				2013	80,918	121,435)	1098.1	(662.7 to 1588.3)
						(63,512 to		
			Both	1990	106,578	154,063)	1367.8	(893.8 to 1818.8)
						(103,077 to		
				2013	163,351	219,080)	1143.2	(752.3 to 1474.7)
		Hemorrhagic				(77,259 to		
		Stroke	Male	1990	115,652	165,519)	1902.4	(1287.6 to 2724.5)
						(103,327 to		,
				2013	150,264	211,357)	1421.5	(974.4 to 1983.2)
				4000	120 105	(70,573 to	2406.6	(4240 C + 2400 C)
			Female	1990	138,105	196,607)	2196.9	(1319.6 to 3109.8)
				2012	140 130	(96,417 to	1402.2	(036 F +c 3069 7)
				2013	148,126	219,014)	1402.2	(926.5 to 2068.7)
			Both	1990	253,757	(177,828 to 332,413)	2054.5	(1494.5 to 2695.1)
			DUII		-			,
1				2013	298,390	(224,074 to	1413.3	(1056.7 to 1887.4)

	1		1		_	I		
						390,087)		
	YLDs	Ischemic Stroke	Male	1990	402	(265 to 584)	9.5	(6.2 to 13.8)
				2013	860	(542 to 1,222)	9.7	(6.0 to 13.9)
			Female	1990	344	(218 to 505)	7.8	(5.0 to 11.4)
				2013	673	(417 to 990)	7.2	(4.5 to 10.6)
			Both	1990	747	(491 to 1,036)	8.6	(5.7 to 12.1)
				2013	1,533	(1,002 to 2,158)	8.4	(5.5 to 11.8)
		Hemorrhagic						
		Stroke	Male	1990	65	(39 to 102)	1.3	(0.7 to 2.0)
				2013	141	(83 to 221)	1.4	(0.8 to 2.1)
			Female	1990	66	(35 to 107)	1.3	(0.7 to 2.1)
				2013	135	(70 to 220)	1.3	(0.6 to 2.1)
			Both	1990	131	(83 to 194)	1.3	(0.8 to 2.0)
			Both	2013	276	(170 to 410)	1.3	(0.8 to 1.9)
Suriname	Deaths	Ischemic Stroke	Male	1990	85	(71 to 102)	95.5	(79.1 to 115.3)
				2013	158	(114 to 208)	102.1	(75.3 to 131.8)
			Female	1990	94	(81 to 119)	89.8	(77.6 to 114.4)
				2013	190	(140 to 243)	93.1	(68.5 to 119.1)
			Both	1990	180	(160 to 212)	92.9	(82.9 to 110.4)
				2013	348	(278 to 423)	98	(78.8 to 118.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	52	(42 to 69)	48.9	(39.4 to 65.4)
				2013	95	(61 to 131)	53.1	(33.3 to 72.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	57	(48 to 77)	48.3	(39.9 to 66.7)
				2013	111	(70 to 152)	52.2	(32.6 to 71.1)
			Both	1990	109	(95 to 139)	48.8	(42.2 to 63.6)

				2013	207	(146 to 264)	53.2	(36.7 to 67.5)
	DALYs	Ischemic Stroke	Male	1990	1,770	(1,472 to 2,105)	1634.5	(1376.0 to 1942.1)
				2013	3,064	(2,174 to 4,072)	1687.9	(1209.2 to 2231.4)
			Female	1990	1,600	(1,367 to 1,976)	1366.8	(1183.2 to 1703.4)
				2013	2,821	(2,038 to 3,697)	1304.8	(946.9 to 1701.2)
			Both	1990	3,371	(2,995 to 3,885)	1497.9	(1341.5 to 1737.3)
				2013	5,885	(4,630 to 7,246)	1487.9	(1178.1 to 1819.0)
		Hemorrhagic						
		Stroke	Male	1990	1,511	(1,229 to 1,943)	1127.6	(918.6 to 1462.9)
				2013	2,512	(1,634 to 3,476)	1159.9	(756.0 to 1590.5)
			Female	1990	1,469	(1,252 to 1,898)	1046.4	(890.7 to 1375.9)
				2013	2,295	(1,528 to 3,102)	984.1	(647.6 to 1326.9)
			Both	1990	2,981	(2,642 to 3,624)	1086.9	(958.9 to 1345.2)
				2013	4,808	(3,562 to 6,139)	1074.1	(781.7 to 1367.2)
	YLDs	Ischemic Stroke	Male	1990	47	(31 to 66)	46.7	(30.9 to 64.7)
				2013	83	(54 to 118)	47	(30.4 to 65.8)
			Female	1990	37	(24 to 53)	32.7	(21.4 to 46.5)
				2013	72	(45 to 105)	32.9	(20.4 to 48.2)
			Both	1990	85	(58 to 116)	39.2	(26.8 to 53.5)
				2013	155	(104 to 217)	39.1	(26.4 to 54.4)
		Hemorrhagic						
		Stroke	Male	1990	16	(10 to 26)	14.5	(8.7 to 22.6)
				2013	28	(17 to 43)	14.2	(8.3 to 21.9)
			Female	1990	12	(8 to 18)	10	(6.4 to 14.6)
				2013	24	(15 to 37)	10.7	(6.6 to 16.0)
			Both	1990	29	(19 to 41)	12.1	(8.0 to 17.3)
			Both	2013	53	(34 to 76)	12.3	(7.9 to 17.6)
Swaziland	Deaths	Ischemic Stroke	Male	1990	53	(34 to 73)	58.2	(38.1 to 78.6)
				2013	141	(94 to 191)	80.2	(55.1 to 105.7)

			Female	1990	82	(53 to 110)	69	(45.2 to 92.9)
				2013	181	(127 to 250)	81.2	(57.2 to 110.1)
			Both	1990	136	(97 to 170)	65.1	(47.5 to 81.6)
				2013	323	(247 to 411)	81.5	(62.8 to 102.3)
	Deaths	Hemorrhagic Stroke	Male	1990	83	(59 to 109)	73.1	(51.4 to 97.1)
	Deatils	Stroke	iviale			, , ,		· · · · · · · · · · · · · · · · · · ·
				2013	176	(126 to 236)	79.2	(57.6 to 107.6)
			Female	1990	114	(77 to 144)	80.6	(52.4 to 102.3)
				2013	191	(136 to 273)	69.5	(48.6 to 99.0)
			Both	1990	197	(151 to 239)	77.9	(58.3 to 94.5)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
•				2013	367	(290 to 479)	74	(58.3 to 97.4)
	DALYs	Ischemic Stroke	Male	1990	1,144	(707 to 1,578)	940.3	(604.0 to 1286.2)
				2013	3,025	(1,884 to 4,389)	1362.4	(892.2 to 1877.7)
			Female	1990	1,491	(912 to 2,024)	1006.2	(643.6 to 1345.0)
				2013	3,144	(2,181 to 4,464)	1156.4	(812.5 to 1595.2)
			Both	1990	2,636	(1,806 to 3,315)	984.2	(698.1 to 1228.2)
				2013	6,169	(4,617 to 8,027)	1251.6	(959.5 to 1594.2)
		Hemorrhagic		4000		(4 -00) 0 00-1		(10010: 10010)
		Stroke	Male	1990	2,257	(1,599 to 2,927)	1429.4	(1021.2 to 1871.9)
				2013	5,256	(3,834 to 6,920)	1768.7	(1265.9 to 2367.1)
			Female	1990	2,725	(1,931 to 3,421)	1455.1	(1003.6 to 1841.5)
				2013	5,162	(3,721 to 7,718)	1408.5	(1003.7 to 2035.2)
			Both	1990	4,983	(3,868 to 5,998)	1448.6	(1112.2 to 1754.2)
				2013	10,419	(8,247 to 13,508)	1567.9	(1234.5 to 2049.6)
	YLDs	Ischemic Stroke	Male	1990	29	(19 to 41)	19.3	(12.7 to 27.1)
				2013	46	(29 to 69)	17.6	(10.7 to 26.2)

			Female	1990	34	(22 to 40)	17.8	(11 6 to 25 2)
			remale		1	(22 to 49)		(11.6 to 25.3)
				2013	58	(38 to 83)	18.1	(11.5 to 25.8)
			Both	1990	64	(43 to 89)	18.5	(12.5 to 25.4)
				2013	105	(69 to 147)	17.9	(11.8 to 25.0)
		Hemorrhagic			_	4-		.
		Stroke	Male	1990	9	(5 to 13)	5.1	(3.1 to 7.5)
				2013	14	(9 to 22)	4.7	(2.9 to 7.1)
			Female	1990	9	(6 to 14)	4.4	(2.7 to 6.3)
				2013	15	(9 to 23)	4.3	(2.6 to 6.4)
			Both	1990	19	(12 to 26)	4.7	(3.0 to 6.6)
			Both	2013	30	(20 to 42)	4.5	(3.0 to 6.3)
Sweden	Deaths	Ischemic Stroke	Male	1990	2,923	(2,356 to 3,741)	47.9	(38.4 to 61.6)
				2013	2,508	(1,808 to 3,179)	30.3	(21.9 to 38.3)
			Female	1990	4,037	(3,245 to 4,965)	40.2	(32.2 to 49.4)
				2013	3,592	(2,691 to 4,674)	27.6	(21.1 to 36.0)
			Both	1990	6,960	(6,073 to 8,141)	43.7	(38.2 to 51.2)
				2013	6,100	(4,865 to 7,271)	29	(23.2 to 34.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,711	(1,283 to 2,220)	28.8	(21.6 to 37.1)
				2013	1,488	(1,014 to 1,939)	18.6	(12.8 to 24.0)
			Female	1990	2,227	(1,589 to 2,823)	25.1	(18.4 to 31.4)
				2013	1,839	(1,316 to 2,638)	15.9	(11.8 to 22.6)
			Both	1990	3,939	(3,219 to 4,731)	26.9	(22.0 to 32.1)
				2013	3,328	(2,641 to 4,241)	17.2	(13.7 to 21.7)
	DALYs	Ischemic Stroke	Male	1990	40,618	(34,097 to 50,192)	668.9	(561.3 to 827.0)
				2013	30,870	(24,023 to 37,652)	390.3	(304.4 to 473.6)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

			Female	1990	41,494	(34,261 to 49,561)	461	(384.8 to 557.1)
				2013	30,973	(24,589 to 38,951)	281.6	(226.9 to 351.1)
			Both	1990	82,113	(72,868 to 94,077)	554.1	(491.0 to 635.5)
				2013	61,843	(51,852 to 72,359)	332.8	(279.3 to 389.9)
		Hemorrhagic						
		Stroke	Male	1990	29,304	(23,510 to 36,907)	517.5	(418.3 to 652.4)
				2013	21,923	(15,586 to 27,093)	296.4	(210.8 to 367.2)
			Female	1990	29,206	(22,326 to 36,030)	404.8	(317.9 to 497.7)
				2013	19,849	(15,491 to 26,859)	220.6	(173.9 to 288.5)
			Both	1990	58,511	(50,304 to 68,705)	456.9	(397.4 to 535.8)
				2013	41,773	(33,567 to 49,780)	256.8	(206.9 to 301.1)
	YLDs	Ischemic Stroke	Male	1990	5,330	(3,422 to 7,763)	89.9	(58.2 to 131.3)
				2013	6,156	(3,859 to 8,891)	82.2	(51.7 to 117.5)
			Female	1990	4,650	(3,019 to 6,564)	60.7	(40.0 to 84.8)
				2013	4,947	(3,313 to 6,837)	56	(37.6 to 77.8)
			Both	1990	9,981	(6,649 to 13,987)	73.8	(49.6 to 103.6)
				2013	11,104	(7,315 to 15,293)	68.6	(45.8 to 94.5)
		Hemorrhagic						
		Stroke	Male	1990	1,617	(930 to 2,489)	28	(16.3 to 42.9)
				2013	1,786	(1,085 to 2,576)	24.9	(15.3 to 35.5)
			Female	1990	1,303	(768 to 1,989)	18.7	(11.5 to 27.6)
				2013	1,459	(917 to 2,185)	18.4	(11.8 to 26.9)
			Both	1990	2,921	(1,906 to 4,165)	22.8	(14.9 to 32.0)
			Both	2013	3,246	(2,091 to 4,517)	21.5	(14.0 to 29.7)
Switzerland	Deaths	Ischemic Stroke	Male	1990	1,704	(1,145 to 2,323)	45.7	(30.6 to 62.3)
				2013	1,600	(1,041 to 2,126)	25.5	(16.5 to 33.9)
			Female	1990	2,896	(2,241 to 3,484)	40.8	(31.5 to 48.9)
				2013	2,653	(1,962 to 3,802)	24.6	(18.2 to 35.1)
			Both	1990	4,601	(3,641 to 5,410)	43.1	(34.3 to 51.0)

				2013	4,253	(3,398 to 5,380)	25.2	(20.2 to 31.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,003	(662 to 1,599)	26.6	(17.5 to 42.6)
				2013	1,054	(631 to 1,503)	17.1	(10.3 to 24.2)
			Female	1990	1,373	(906 to 1,796)	21.3	(14.3 to 27.2)
				2013	1,423	(957 to 2,081)	14.3	(10.0 to 20.2)
			Both	1990	2,376	(1,807 to 3,212)	23.6	(18.1 to 31.6)
				2013	2,477	(1,797 to 3,327)	15.6	(11.5 to 20.5)
	DALYs	Ischemic Stroke	Male	1990	23,096	(16,316 to 30,278)	606.9	(429.6 to 794.3)
				2013	19,871	(14,279 to 24,946)	328.1	(236.7 to 411.7)
			Female	1990	28,154	(22,738 to 33,313)	444.1	(359.1 to 523.1)
				2013	22,551	(17,367 to 30,811)	247.3	(190.3 to 337.9)
			Both	1990	51,251	(41,731 to 60,593)	513.5	(416.2 to 603.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	42,423	(35,155 to 51,229)	284.5	(234.3 to 343.3)
		Hemorrhagic						
		Stroke	Male	1990	16,547	(11,843 to 24,510)	433.4	(310.2 to 643.5)
				2013	13,996	(9,172 to 18,326)	237.4	(156.6 to 308.7)
			Female	1990	16,903	(11,925 to 20,821)	316.2	(227.3 to 382.6)
				2013	13,580	(10,142 to 18,324)	172.8	(131.8 to 226.3)
			Both	1990	33,450	(26,995 to 43,155)	366.8	(298.8 to 468.2)
				2013	27,577	(21,312 to 34,346)	202.4	(157.9 to 247.7)
	YLDs	Ischemic Stroke	Male	1990	3,288	(2,058 to 4,704)	85.7	(53.6 to 122.3)
					-,	, , ,		
				2013	5,110	(3,322 to 7,066)	87.5	(57.0 to 121.3)
			Female	2013 1990			87.5 62.4	(57.0 to 121.3) (41.3 to 86.6)
					5,110	(3,322 to 7,066)		` '

		ı	1 -		1		T	T
				2013	9,648	(6,577 to 12,988)	73.9	(50.3 to 99.6)
		Hemorrhagic						
		Stroke	Male	1990	657	(386 to 1,060)	17.3	(10.2 to 27.6)
				2013	1,013	(528 to 1,781)	17.9	(9.5 to 31.1)
			Female	1990	680	(369 to 1,072)	13.6	(7.5 to 21.7)
				2013	947	(539 to 1,718)	14	(8.0 to 24.7)
			Both	1990	1,337	(847 to 1,977)	15.2	(9.7 to 22.5)
			Both	2013	1,960	(1,167 to 3,029)	15.8	(9.7 to 24.2)
Syria	Deaths	Ischemic Stroke	Male	1990	1,154	(719 to 1,745)	63.8	(39.5 to 97.4)
				2013	1,857	(715 to 2,971)	46	(17.7 to 74.1)
			Female	1990	1,144	(779 to 1,967)	57.6	(39.0 to 99.3)
				2013	1,877	(976 to 2,621)	43.6	(22.6 to 61.1)
			Both	1990	2,298	(1,700 to 3,380)	60.6	(44.9 to 89.5)
				2013	3,735	(2,133 to 5,107)	44.8	(25.3 to 61.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,159	(3,206 to 5,400)	165.6	(126.3 to 213.8)
				2013	5,487	(3,580 to 7,498)	113.7	(71.8 to 156.6)
			Female	1990	2,503	(1,845 to 3,708)	111.7	(81.6 to 170.9)
				2013	3,959	(2,682 to 5,209)	87.3	(57.5 to 115.2)
			Both	1990	6,663	(5,388 to 8,425)	138.4	(112.2 to 177.5)
				2013	9,446	(7,134 to 11,925)	100.4	(74.9 to 130.2)
	DALYs	Ischemic Stroke	Male	1990	28,058	(17,424 to 43,257)	1073.7	(671.7 to 1606.8)
				2013	33,728	(13,465 to 54,795)	665	(264.9 to 1070.5)
			Female	1990	30,195	(21,001 to 49,402)	1010.7	(686.0 to 1746.6)
				2013	33,426	(16,705 to 47,697)	602.9	(312.8 to 848.5)
			Both	1990	58,253	(43,839 to 85,340)	1041.1	(770.9 to 1519.1)
				2013	67,154	(38,583 to 91,753)	633	(367.1 to 867.3)
		Hemorrhagic			, -	(121,342 to		,,
		Stroke	Male	1990	158,616	205,125)	3951.6	(3007.4 to 5148.4)

					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(94,109 to		
				2013	139,530	193,197)	2130.2	(1398.2 to 2919.9)
						(57,897 to		
			Female	1990	82,109	118,405)	2245.4	(1624.8 to 3365.2)
						(54,851 to		(
				2013	79,477	105,639)	1334.6	(900.1 to 1765.5)
			Doth	1000	240.726	(189,777 to	2402.2	(2400 1 += 2022 0)
			Both	1990	240,726	301,335) (170,266 to	3103.3	(2490.1 to 3933.9)
				2013	219,007	277,326)	1731.6	(1309.0 to 2196.8)
	YLDs	Ischemic Stroke	Male	1990	234	(155 to 337)	9	(6.0 to 13.2)
		isoneime serone	Maic	2013	598	(392 to 836)	9.5	(6.3 to 13.3)
			Female	1990	189	(125 to 277)	7.4	(4.9 to 10.8)
				2013	460	(294 to 657)	7.4	(4.8 to 10.4)
			Both	1990	423	(286 to 584)	8.2	(5.6 to 11.2)
				2013	1,058	(706 to 1,421)	8.4	(5.7 to 11.3)
		Hemorrhagic						
		Stroke	Male	1990	41	(24 to 64)	1.4	(0.8 to 2.1)
				2013	97	(58 to 152)	1.4	(0.8 to 2.1)
			Female	1990	39	(21 to 60)	1.4	(0.8 to 2.1)
				2013	91	(44 to 151)	1.3	(0.7 to 2.2)
			Both	1990	81	(51 to 118)	1.4	(0.9 to 2.0)
			Both	2013	189	(117 to 286)	1.3	(0.8 to 2.0)
Taiwan	Deaths	Ischemic Stroke	Male	1990	2,838	(2,150 to 3,598)	53.4	(40.0 to 67.9)
				2013	3,770	(3,030 to 4,707)	26.3	(21.3 to 33.0)
			Female	1990	2,169	(1,467 to 2,912)	44.4	(30.3 to 59.6)
				2013	3,045	(2,342 to 4,054)	19.5	(15.0 to 25.9)

			Both	1990	5,008	(4,001 to 6,005)	49.4	(39.5 to 59.0)
				2013	6,816	(5,851 to 8,123)	22.6	(19.6 to 27.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	4,144	(3,300 to 5,210)	62.4	(49.0 to 78.5)
				2013	4,386	(3,610 to 5,311)	30.3	(24.9 to 36.4)
			Female	1990	2,865	(1,927 to 3,608)	50.4	(33.1 to 64.1)
				2013	2,998	(2,366 to 4,009)	19.5	(15.5 to 25.9)
			Both	1990	7,009	(5,692 to 8,317)	57	(45.1 to 68.3)
				2013	7,385	(6,422 to 8,848)	24.7	(21.5 to 29.5)
	DALYs	Ischemic Stroke	Male	1990	53,223	(40,418 to 67,020)	803.8	(621.4 to 1009.4)
				2013	54,481	(45,536 to 66,295)	392.6	(329.1 to 476.1)
			Female	1990	31,659	(21,731 to 42,056)	563.6	(390.6 to 742.9)
				2013	34,272	(28,033 to 43,499)	224.1	(183.8 to 286.0)
						(68,053 to		
			Both	1990	84,883	103,166)	692.9	(559.7 to 829.8)
						(78,094 to		
				2013	88,754	103,986)	303.8	(267.7 to 356.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic				(87,358 to		
		Stroke	Male	1990	108,451	131,172)	1339.1	(1081.2 to 1645.6)
						(82,476 to		
				2013	96,943	113,804)	673.3	(572.4 to 783.8)
			Female	1990	63,163	(45,518 to 78,777)	919.6	(647.9 to 1143.7)
				2013	50,855	(43,119 to 65,378)	340.3	(290.6 to 440.2)
						(143,926 to		
			Both	1990	171,614	199,058)	1141.9	(946.6 to 1330.2)
						(131,553 to		

	YLDs	Ischemic Stroke	Male	1990	3,976	(2,465 to 5,759)	57	(35.5 to 81.2)
				2013	6,669	(3,914 to 9,722)	47.8	(27.8 to 69.7)
			Female	1990	2,200	(1,370 to 3,183)	35.2	(22.0 to 51.3)
				2013	4,813	(3,038 to 7,035)	31.6	(19.9 to 46.5)
			Both	1990	6,176	(4,070 to 8,707)	46.4	(30.9 to 64.4)
				2013	11,482	(7,505 to 16,092)	39.3	(25.7 to 55.2)
		Hemorrhagic						
		Stroke	Male	1990	3,696	(2,474 to 5,179)	50.9	(33.7 to 72.4)
				2013	6,939	(4,283 to 10,244)	49.2	(30.3 to 72.5)
			Female	1990	2,400	(1,548 to 3,482)	36.2	(23.7 to 52.1)
				2013	4,763	(3,088 to 6,876)	31.3	(20.3 to 44.9)
			Both	1990	6,096	(4,188 to 8,373)	43.6	(29.9 to 59.8)
			Both	2013	11,702	(7,795 to 16,447)	39.8	(26.4 to 56.3)
Tajikistan	Deaths	Ischemic Stroke	Male	1990	877	(539 to 1,392)	103.9	(65.7 to 159.0)
				2013	1,140	(684 to 1,707)	91.7	(57.3 to 130.1)
			Female	1990	1,082	(823 to 1,515)	87.7	(67.1 to 121.9)
				2013	1,301	(751 to 1,699)	84.8	(48.7 to 110.2)
			Both	1990	1,959	(1,557 to 2,680)	94.2	(75.4 to 126.6)
				2013	2,442	(1,572 to 3,012)	88.2	(56.9 to 107.9)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,023	(552 to 1,363)	106.4	(56.1 to 144.2)
				2013	1,273	(768 to 1,814)	87.8	(51.8 to 127.5)
			Female	1990	1,218	(847 to 1,523)	92.9	(63.4 to 116.1)
				2013	1,457	(1,071 to 1,940)	84.3	(60.9 to 114.1)
			Both	1990	2,241	(1,537 to 2,654)	98.2	(66.0 to 117.1)
				2013	2,731	(2,087 to 3,492)	86	(65.2 to 112.1)
	DALYs	Ischemic Stroke	Male	1990	14,666	(8,448 to 25,292)	1506.6	(898.1 to 2476.6)
				2013	18,770	(10,535 to 30,389)	1272.4	(744.0 to 1965.2)
			Female	1990	15,690	(11,633 to 22,752)	1188.2	(888.3 to 1701.9)

				2013	18,083	(10,497 to 24,112)	1071.4	(617.1 to 1438.2)
			Both	1990	30,357	(23,270 to 43,829)	1319.5	(1025.8 to 1861.9)
				2013	36,853	(23,516 to 48,038)	1167.2	(748.0 to 1468.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	22,535	(12,540 to 29,376)	1960.7	(1067.4 to 2580.1)
				2013	28,588	(17,485 to 40,093)	1570.6	(940.8 to 2213.5)
			Female	1990	24,586	(18,338 to 30,491)	1724.6	(1259.9 to 2121.6)
				2013	30,767	(23,018 to 40,243)	1496.7	(1106.3 to 1983.9)
			Both	1990	47,121	(34,019 to 55,043)	1817.5	(1280.1 to 2136.7)
				2013	59,355	(46,138 to 73,790)	1530.7	(1174.0 to 1934.3)
	YLDs	Ischemic Stroke	Male	1990	104	(65 to 156)	8.7	(5.4 to 12.9)
				2013	168	(105 to 250)	8.5	(5.2 to 12.6)
			Female	1990	112	(70 to 160)	7.8	(4.9 to 11.3)
				2013	170	(106 to 248)	8	(5.1 to 11.5)
			Both	1990	217	(140 to 306)	8.2	(5.4 to 11.6)
				2013	338	(228 to 470)	8.2	(5.6 to 11.3)
		Hemorrhagic						
		Stroke	Male	1990	116	(72 to 168)	8.3	(5.2 to 11.9)
				2013	188	(121 to 274)	7.8	(5.0 to 11.3)
			Female	1990	74	(45 to 108)	4.7	(2.9 to 7.0)
				2013	117	(72 to 173)	4.8	(3.0 to 7.1)
			Both	1990	190	(124 to 263)	6.4	(4.2 to 8.8)
			Both	2013	306	(203 to 425)	6.2	(4.2 to 8.5)
Tanzania	Deaths	Ischemic Stroke	Male	1990	1,003	(267 to 1,542)	38.3	(10.3 to 58.6)
				2013	2,477	(849 to 3,683)	42.1	(14.9 to 61.7)

			Female	1990	1,086	(459 to 1,560)	33.5	(13.7 to 49.5)
				2013	2,536	(1,163 to 3,808)	34.1	(15.8 to 51.9)
			Both	1990	2,090	(897 to 2,931)	35.6	(15.5 to 50.0)
				2013	5,013	(2,398 to 7,038)	37.6	(18.2 to 52.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,430	(948 to 2,002)	40.1	(24.0 to 60.1)
				2013	3,004	(2,017 to 4,518)	41.1	(25.9 to 65.5)
			Female	1990	1,583	(1,099 to 2,263)	37.7	(24.3 to 56.1)
				2013	3,148	(2,111 to 4,572)	35.4	(22.7 to 53.4)
			Both	1990	3,014	(2,280 to 3,892)	38.8	(26.9 to 52.0)
				2013	6,152	(4,534 to 8,434)	37.9	(26.2 to 52.9)
	DALYs	Ischemic Stroke	Male	1990	19,243	(5,896 to 28,872)	537.7	(154.0 to 819.2)
				2013	41,285	(14,263 to 62,043)	560.4	(196.6 to 832.4)
			Female	1990	18,641	(8,509 to 26,611)	453.2	(201.7 to 645.2)
				2013	39,533	(18,515 to 58,705)	446.7	(209.8 to 666.3)
			Both	1990	37,884	(17,400 to 51,913)	491.7	(220.4 to 683.2)
						(39,023 to		
				2013	80,819	113,084)	497.5	(241.3 to 697.0)
		Hemorrhagic						
		Stroke	Male	1990	39,248	(28,641 to 51,108)	753.7	(500.7 to 1052.8)
				2013	71,625	(50,449 to 99,711)	713.1	(485.6 to 1061.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	39,659	(28,627 to 55,048)	693.9	(486.6 to 989.0)
						(50,953 to		
				2013	71,793	101,755)	627.4	(426.2 to 898.8)
			Both	1990	78,908	(63,915 to 99,053)	722.2	(550.5 to 932.3)
					440	(111,785 to		/ = 00.0 / 555.5
				2013	143,418	182,026)	666.3	(503.9 to 899.5)

	YLDs	Ischemic Stroke	Male	1990	688	(421 to 1,018)	14	(8.8 to 20.6)
				2013	1,300	(799 to 1,971)	12.1	(7.3 to 18.5)
			Female	1990	573	(350 to 862)	10.4	(6.4 to 15.4)
				2013	1,049	(656 to 1,568)	9	(5.7 to 13.6)
			Both	1990	1,262	(819 to 1,810)	12	(7.9 to 17.1)
				2013	2,350	(1,528 to 3,371)	10.4	(6.8 to 15.0)
		Hemorrhagic						
	Stroke	Stroke	Male	1990	240	(148 to 349)	3.9	(2.5 to 5.9)
				2013	457	(280 to 693)	3.5	(2.0 to 5.3)
			Female	1990	180	(109 to 276)	2.8	(1.7 to 4.2)
				2013	326	(195 to 495)	2.4	(1.4 to 3.7)
			Both	1990	421	(277 to 599)	3.3	(2.2 to 4.7)
			Both	2013	784	(502 to 1,144)	2.9	(1.8 to 4.3)
Thailand	Deaths	Ischemic Stroke	Male	1990	2,783	(2,132 to 4,033)	26.4	(20.1 to 38.1)
				2013	9,007	(5,485 to 12,501)	32	(19.2 to 44.8)
			Female	1990	3,874	(3,037 to 5,122)	29.3	(23.1 to 38.8)
				2013	12,045	(7,433 to 16,783)	33.7	(20.8 to 46.9)
			Both	1990	6,657	(5,701 to 8,726)	28.4	(24.1 to 37.3)
				2013	21,052	(14,733 to 27,158)	33.1	(23.2 to 43.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	12,432	(10,545 to 14,982)	99.5	(84.5 to 116.1)
				2013	19,574	(14,442 to 26,585)	65.4	(47.7 to 90.1)
			Female	1990	14,033	(10,059 to 15,769)	100.7	(72.8 to 113.4)
				2013	21,073	(14,225 to 35,291)	58.1	(39.0 to 97.9)
			Both	1990	26,466	(22,190 to 29,292)	101.7	(83.7 to 111.9)
				2013	40,647	(30,836 to 58,425)	61.9	(46.6 to 90.2)
	DALYs	Ischemic Stroke	Male	1990	56,089	(43,270 to 81,767)	394.9	(308.5 to 564.2)
				2013	145,097	(93,598 to 195,725)	460.1	(297.8 to 619.4)

Female 1990 55,801 (44,670 to 73,184) 359.7 (294.0 to 476.9)									
Both 1990 111,891 144,911 378.9 (324.8 to 489.4)				Female	1990	55,801	(44,670 to 73,184)	359.7	(294.0 to 476.9)
Both 1990 111,891 (95,057 to 144,911) 378.9 (324.8 to 489.4)							(95,284 to		
Both 1990 111,891 144,911 378.9 (324.8 to 489.4)					2013	146,257	193,084)	391	(255.5 to 521.7)
Hemorrhagic Stroke Male 1990 327,815 359,999 424.7 (314.9 to 527.3)							(95,057 to		
Hemorrhagic Stroke Male 1990 327,815 359,999 424.7 (314.9 to 527.3)				Both	1990	111,891	144,911)	378.9	(324.8 to 489.4)
Hemorrhagic Stroke Male 1990 327,815 427,660) 1824.5 (1545.7 to 2247.3)							(215,920 to		
Stroke Male 1990 327,815 427,660) 1824.5 (1545.7 to 2247.3)					2013	291,355	359,999)	424.7	(314.9 to 527.3)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI (305,221 to 536,209) 1188.8 (897.1 to 1565.6) (163,869 to 163,869 to 164,369 to 164,3705) (214,572 to 164,3705) (775.4 (555.7 to 1228.7) (800			Hemorrhagic				(271,320 to		
Age Standardized Country measure Stroke Type Sex year Mean 95% Ul Mean 95% Ul			Stroke	Male	1990	327,815	427,660)	1824.5	(1545.7 to 2247.3)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI 2013 405,289 536,209 1188.8 (897.1 to 1565.6) 1 1990 240,201 272,101 1384.7 (978.9 to 1549.6) 2013 296,690 463,705 775.4 (555.7 to 1228.7) 80th 1990 568,017 653,827) 1606.7 (1363.0 to 1793.9) (553,477 to 2013 701,979 919,152) 975.6 (766.6 to 1298.3) YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) YLDs Ischemic Stroke Male 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) Emale 1990 7,655 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 <td></td> <td></td> <td></td> <td></td> <td></td> <td>Counts, All</td> <td>Ages</td> <td>Rate</td> <td>Per 100,000</td>						Counts, All	Ages	Rate	Per 100,000
2013 405,289 536,209 1188.8 (897.1 to 1565.6) Female 1990 240,201 272,101 1384.7 (978.9 to 1549.6) 2013 296,690 463,705 775.4 (555.7 to 1228.7) 8								Age-	standardized
2013 405,289 536,209 1188.8 (897.1 to 1565.6)	Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
Female 1990 240,201 272,101 1384.7 (978.9 to 1549.6) (214,572 to 2013 296,690 463,705) 775.4 (555.7 to 1228.7) (482,759 to 63,827) 1606.7 (1363.0 to 1793.9) (553,477 to 919,152) 975.6 (766.6 to 1298.3) (2712 to 5,506) 28.6 (19.3 to 40.5) (2013 9,638 (6,464 to 13,453) 30 (20.1 to 41.8) (2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) (2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) (20.9 to 43.1) (20.9 to 43.1)							(305,221 to		
Female 1990 240,201 272,101) 1384.7 (978.9 to 1549.6) 2013 296,690 463,705) 775.4 (555.7 to 1228.7) Both 1990 568,017 653,827) 1606.7 (1363.0 to 1793.9) 2013 701,979 919,152) 975.6 (766.6 to 1298.3) YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) 2013 9,638 (6,464 to 13,453) 30 (20.1 to 41.8) Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)					2013	405,289	536,209)	1188.8	(897.1 to 1565.6)
2013 296,690 463,705) 775.4 (555.7 to 1228.7) (482,759 to 653,827) 1606.7 (1363.0 to 1793.9) (553,477 to 919,152) 975.6 (766.6 to 1298.3) (176.6 to 1298.3) (176							(163,869 to		
2013 296,690 463,705 775.4 (555.7 to 1228.7)				Female	1990	240,201	272,101)	1384.7	(978.9 to 1549.6)
Both 1990 568,017 653,827) 1606.7 (1363.0 to 1793.9)							(214,572 to		
Both 1990 568,017 653,827) 1606.7 (1363.0 to 1793.9) YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) YLDs Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) YLDs Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) YLDs Both 1990 7,655 (5,813 to 12,755) 23 (15.0 to 33.0) YLDs Hemorrhagic 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0)					2013	296,690	463,705)	775.4	(555.7 to 1228.7)
2013 701,979 919,152 975.6 (766.6 to 1298.3)							(482,759 to		
YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) 2013 9,638 (6,464 to 13,453) 30 (20.1 to 41.8) Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) 2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) 2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)				Both	1990	568,017	653,827)	1606.7	(1363.0 to 1793.9)
YLDs Ischemic Stroke Male 1990 3,981 (2,712 to 5,506) 28.6 (19.3 to 40.5) 2013 9,638 (6,464 to 13,453) 30 (20.1 to 41.8) Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) 2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) 2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)							(553,477 to		
2013 9,638 (6,464 to 13,453) 30 (20.1 to 41.8)					2013	701,979	919,152)	975.6	(766.6 to 1298.3)
Female 1990 3,673 (2,461 to 5,228) 22.6 (14.8 to 32.0) 2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) 2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)		YLDs	Ischemic Stroke	Male	1990	3,981	(2,712 to 5,506)	28.6	(19.3 to 40.5)
2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) 2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)					2013	9,638	(6,464 to 13,453)	30	(20.1 to 41.8)
2013 8,870 (5,813 to 12,755) 23 (15.0 to 33.0) Both 1990 7,655 (5,322 to 10,435) 25.3 (17.7 to 34.4) 2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)				Female	1990	3,673	(2,461 to 5,228)	22.6	(14.8 to 32.0)
2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)					2013	8,870	(5,813 to 12,755)	23	(15.0 to 33.0)
2013 18,508 (12,663 to 25,591) 26.1 (17.8 to 36.0) Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)				Both	1990	7,655	(5,322 to 10,435)	25.3	(17.7 to 34.4)
Hemorrhagic Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)					2013	18,508	(12,663 to 25,591)	26.1	(17.8 to 36.0)
Stroke Male 1990 4,845 (3,222 to 6,833) 31.1 (20.9 to 43.1)			Hemorrhagic						
2013 10,715 (7,232 to 14,894) 32 (21.7 to 44.4)			_	Male	1990	4,845	(3,222 to 6,833)	31.1	(20.9 to 43.1)
					2013	10,715	(7,232 to 14,894)	32	(21.7 to 44.4)

			Female	1990	3,896	(2,557 to 5,447)	22	(14.0 to 31.1)
				2013	8,562	(5,498 to 12,186)	21.8	(14.1 to 31.0)
			Both	1990	8,742	(5,932 to 11,964)	26.1	(17.7 to 35.7)
			Both	2013	19,277	(13,223 to 26,427)	26.5	(18.2 to 36.2)
The	Deaths	Ischemic Stroke	Male	1990	45	(37 to 55)	99.8	(79.5 to 122.2)
Bahamas				2013	84	(59 to 110)	73.6	(52.9 to 95.8)
			Female	1990	58	(44 to 67)	92.8	(71.8 to 107.5)
				2013	109	(77 to 143)	64.4	(45.7 to 83.7)
			Both	1990	104	(88 to 117)	96.4	(80.9 to 108.6)
				2013	194	(154 to 233)	68.8	(55.2 to 82.6)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	29	(23 to 37)	52.4	(41.1 to 67.5)
				2013	57	(37 to 78)	42.6	(27.5 to 58.0)
			Female	1990	35	(28 to 45)	51.1	(40.8 to 65.5)
				2013	70	(46 to 95)	39.8	(26.4 to 53.8)
			Both	1990	65	(56 to 80)	52	(44.1 to 63.8)
				2013	128	(95 to 159)	41.4	(31.0 to 51.5)
	DALYs	Ischemic Stroke	Male	1990	949	(762 to 1,147)	1635.9	(1316.5 to 1980.1)
				2013	1,607	(1,111 to 2,160)	1154.8	(822.3 to 1514.9)
			Female	1990	946	(733 to 1,105)	1328.7	(1034.5 to 1545.8)
				2013	1,564	(1,089 to 2,086)	871.7	(611.0 to 1164.0)
			Both	1990	1,896	(1,626 to 2,132)	1470.2	(1259.7 to 1650.1)
				2013	3,171	(2,505 to 3,858)	1000.8	(795.9 to 1210.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	855	(686 to 1,057)	1138.7	(907.3 to 1430.4)

				2013	1,406	(902 to 1,968)	864.2	(567.8 to 1186.2)
			Female	1990	855	(685 to 1,055)	1008.5	(811.9 to 1263.4)
				2013	1,365	(882 to 1,868)	714	(470.3 to 962.7)
			Both	1990	1,710	(1,480 to 2,050)	1071.3	(930.7 to 1301.0)
				2013	2,771	(2,068 to 3,497)	786.4	(592.2 to 983.4)
	YLDs	Ischemic Stroke	Male	1990	26	(17 to 37)	47.3	(31.5 to 67.7)
				2013	67	(43 to 95)	48.4	(31.4 to 69.5)
			Female	1990	23	(15 to 33)	32.5	(21.7 to 46.1)
				2013	58	(38 to 85)	32.1	(20.9 to 47.0)
			Both	1990	49	(34 to 68)	38.8	(26.5 to 53.7)
				2013	125	(86 to 172)	39	(27.0 to 53.6)
		Hemorrhagic						
		Stroke	Male	1990	9	(5 to 14)	14.4	(8.7 to 22.5)
				2013	22	(14 to 34)	14.6	(9.2 to 21.7)
			Female	1990	8	(4 to 12)	10.1	(6.0 to 15.2)
				2013	21	(12 to 38)	11.1	(6.5 to 20.6)
			Both	1990	17	(11 to 25)	12	(7.6 to 17.5)
			Both	2013	43	(28 to 64)	12.6	(8.1 to 19.0)
The	Deaths	Ischemic Stroke	Male	1990	80	(45 to 133)	68.7	(42.7 to 106.9)
Gambia				2013	219	(127 to 331)	104.1	(65.0 to 149.9)
			Female	1990	80	(40 to 127)	79.3	(44.3 to 119.0)
				2013	220	(97 to 341)	112.1	(53.6 to 167.6)
			Both	1990	160	(99 to 248)	73.9	(49.0 to 108.1)
				2013	439	(247 to 668)	108.2	(64.6 to 155.3)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	81	(47 to 134)	52	(29.8 to 88.1)
				2013	131	(68 to 252)	47.1	(23.5 to 93.8)
			Female	1990	87	(52 to 135)	66.1	(40.5 to 102.9)
				2013	138	(77 to 265)	52.2	(28.5 to 106.2)

			Both	1990	169	(106 to 267)	58.7	(36.8 to 92.4)
				2013	270	(153 to 469)	49.7	(27.6 to 93.7)
	DALYs	Ischemic Stroke	Male	1990	1,749	(947 to 2,947)	1090.5	(623.2 to 1810.5)
				2013	4,393	(2,406 to 6,887)	1602.8	(928.7 to 2415.8)
			Female	1990	1,621	(722 to 2,609)	1183.4	(584.2 to 1862.3)
				2013	4,042	(1,671 to 6,562)	1606.2	(703.1 to 2514.8)
			Both	1990	3,371	(1,996 to 5,230)	1136	(699.6 to 1745.3)
				2013	8,435	(4,410 to 13,169)	1605	(892.0 to 2448.6)
		Hemorrhagic						
		Stroke	Male	1990	2,719	(1,671 to 4,339)	1089.4	(619.5 to 1805.2)
				2013	3,916	(2,135 to 6,785)	963.1	(498.5 to 1861.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	2,656	(1,555 to 3,912)	1297.5	(774.7 to 2051.2)
				2013	3,928	(2,269 to 6,765)	1030.6	(578.6 to 1971.3)
			Both	1990	5,375	(3,478 to 8,097)	1186	(732.2 to 1901.8)
				2013	7,845	(4,625 to 12,863)	996.5	(562.6 to 1748.5)
	YLDs	Ischemic Stroke	Male	1990	29	(19 to 42)	15.7	(10.2 to 22.6)
				2013	55	(36 to 77)	15.6	(10.2 to 21.8)
			Female	1990	19	(12 to 29)	11.9	(7.7 to 17.7)
				2013	44	(28 to 65)	12.3	(8.1 to 17.6)
			Both	1990	49	(33 to 69)	13.9	(9.2 to 19.8)
				2013	100	(68 to 138)	14	(9.6 to 18.9)
		Hemorrhagic						
		Stroke	Male	1990	9	(6 to 13)	4.2	(2.7 to 6.2)
				2013	17	(11 to 25)	4	(2.6 to 5.9)
			Female	1990	6	(3 to 8)	3	(1.9 to 4.4)
				2013	12	(7 to 18)	2.8	(1.7 to 4.1)

			Both	1990	15	(10 to 21)	3.6	(2.4 to 5.1)
			Both	2013	30	(20 to 42)	3.4	(2.3 to 4.8)
Timor-Leste	Deaths	Ischemic Stroke	Male	1990	53	(25 to 91)	69.6	(33.8 to 112.9)
				2013	115	(55 to 185)	82.9	(43.2 to 126.1)
			Female	1990	54	(28 to 109)	78.9	(43.7 to 140.4)
				2013	138	(69 to 237)	88.9	(46.4 to 145.6)
			Both	1990	107	(68 to 165)	74	(48.0 to 105.8)
				2013	254	(168 to 352)	86.5	(58.8 to 116.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	86	(44 to 122)	65.7	(26.1 to 102.6)
				2013	120	(67 to 189)	60.3	(32.0 to 100.5)
			Female	1990	76	(40 to 113)	64.6	(29.8 to 100.5)
				2013	108	(59 to 172)	52.3	(27.1 to 89.9)
			Both	1990	163	(106 to 215)	65.1	(38.4 to 88.6)
				2013	229	(151 to 311)	56.2	(36.0 to 79.5)
	DALYs	Ischemic Stroke	Male	1990	1,406	(759 to 2,441)	1082.9	(529.7 to 1834.4)
				2013	2,252	(1,052 to 3,793)	1170.3	(575.7 to 1883.3)
			Female	1990	1,379	(636 to 3,104)	1194.8	(642.9 to 2308.3)
				2013	2,373	(1,163 to 4,392)	1169.5	(591.9 to 2024.5)
			Both	1990	2,786	(1,821 to 4,783)	1136	(730.0 to 1705.1)
				2013	4,626	(2,995 to 6,749)	1173.6	(782.6 to 1641.6)
		Hemorrhagic						
		Stroke	Male	1990	3,483	(2,024 to 4,870)	1525.4	(732.3 to 2179.5)
				2013	3,781	(2,378 to 5,516)	1303.6	(748.5 to 2001.4)
			Female	1990	2,884	(1,540 to 4,130)	1445	(748.7 to 2158.2)
				2013	2,890	(1,694 to 4,386)	1040.1	(588.5 to 1611.6)
			Both	1990	6,367	(4,194 to 8,475)	1485.4	(949.4 to 1962.3)
					Counts, All	Λαος	Pato	Per 100.000

Counts, All Ages

Rate Per 100,000 Age- standardized

Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	6,672	(4,672 to 8,813)	1171	(787.9 to 1570.3)
	YLDs	Ischemic Stroke	Male	1990	30	(20 to 43)	28.3	(18.5 to 39.3)
				2013	57	(36 to 82)	28.8	(18.6 to 41.2)
			Female	1990	23	(14 to 33)	22.7	(14.6 to 32.8)
				2013	54	(35 to 78)	25.1	(16.5 to 36.0)
			Both	1990	54	(37 to 75)	25.6	(17.7 to 35.2)
				2013	112	(74 to 157)	26.9	(18.0 to 37.6)
		Hemorrhagic						
		Stroke	Male	1990	41	(26 to 58)	30.8	(20.0 to 43.4)
				2013	69	(44 to 98)	31.1	(20.2 to 43.7)
			Female	1990	28	(18 to 40)	22.4	(14.8 to 31.8)
				2013	56	(36 to 78)	23.6	(15.5 to 33.5)
			Both	1990	70	(46 to 95)	26.7	(18.0 to 36.9)
			Both	2013	125	(83 to 171)	27.2	(18.1 to 37.4)
Togo	Deaths	Ischemic Stroke	Male	1990	336	(224 to 453)	72.4	(48.9 to 95.7)
				2013	734	(330 to 1,031)	97.9	(48.8 to 132.7)
			Female	1990	444	(281 to 605)	80.5	(53.4 to 112.6)
				2013	699	(338 to 966)	83.7	(43.0 to 111.8)
			Both	1990	781	(587 to 951)	77.3	(58.2 to 95.8)
				2013	1,433	(750 to 1,869)	90.2	(49.3 to 114.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	378	(277 to 503)	59.2	(40.9 to 79.9)
				2013	578	(338 to 963)	54.3	(29.6 to 98.9)
			Female	1990	466	(343 to 656)	67	(47.7 to 97.4)
				2013	524	(342 to 1,003)	45.1	(27.7 to 95.6)
			Both	1990	845	(649 to 1,073)	63.8	(47.7 to 85.4)
				2013	1,102	(732 to 1,858)	49.3	(30.6 to 94.1)
	DALYs	Ischemic Stroke	Male	1990	7,651	(5,109 to 10,406)	1160.8	(777.3 to 1562.9)

Female 1990 15,344 (6,683 to 22,079) 1482.2 (675.6 to 2070.6)									•
Both 1990 12,724 (9,517 to 16,529) 1267.7 (926.1 to 1689.0)					2013	15,344	(6,683 to 22,079)	1482.2	(675.6 to 2070.6)
Both 1990 16,445 (12,598 to 20,082) 1183.8 (886.5 to 1437.2)				Female	1990	8,793	(5,353 to 11,926)	1192.7	(753.6 to 1616.6)
Hemorrhagic Stroke Male 1990 12,724 (9,517 to 16,529) 1267.7 (926.1 to 1689.0)					2013	12,817	(5,806 to 18,251)	1129.1	(554.5 to 1553.7)
Hemorrhagic Stroke Male 1990 12,724 (9,517 to 16,529) 1267.7 (926.1 to 1689.0)				Both	1990	16,445	(12,598 to 20,082)	1183.8	(886.5 to 1437.2)
Stroke Male 1990 12,724 (9,517 to 16,529) 1267.7 (926.1 to 1689.0) 1 2013 18,324 (11,552 to 27,261) 1120.5 (654.2 to 1888.1) 1 Female 1990 13,420 (9,765 to 18,327) 1314.4 (967.9 to 1832.7) 2013 14,990 (10,176 to 25,755) 859.4 (562.7 to 1635.6) (562.7 to 1635.6) 3 2013 33,314 (23,523 to 49,677) 980.8 (650.1 to 1647.8) 4 1 2013 33,314 (23,523 to 49,677) 980.8 (650.1 to 1647.8) 5 1 2013 216 (139 to 312) 15.4 (9.9 to 22.3) 6 2013 216 (139 to 312) 15.4 (9.9 to 22.3) (9.9 to 22.3) 7 80 8x year Mean 95% UI Mean 95% UI 8 1 9x 4x Mean 95% UI Mean 95% UI 9 2013 186 (119 to 259) 11.5					2013	28,161	(13,972 to 37,650)	1290.3	(680.0 to 1680.6)
Stroke Type Sex Year Mean Year Stroke Type Sex Year Mean Year Stroke Type Sex Year Mean Year Yea			Hemorrhagic						
Female 1990			Stroke	Male	1990	12,724	(9,517 to 16,529)	1267.7	(926.1 to 1689.0)
Stroke Type Sex Year Mean 95% U (67 to 155) 11.9 (7.5 to 17.8)					2013	18,324	(11,552 to 27,261)	1120.5	(654.2 to 1888.1)
Both 1990 26,145 (20,502 to 31,784) 1298.3 (999.3 to 1653.9)				Female	1990	13,420	(9,765 to 18,327)	1314.4	(967.9 to 1832.7)
YLDs Ischemic Stroke Male 1990 114 (69 to 169) 14.5 (8.9 to 21.3)					2013	14,990	(10,176 to 25,755)	859.4	(562.7 to 1635.6)
YLDs Ischemic Stroke Male 1990 114 (69 to 169) 14.5 (8.9 to 21.3) 2013 216 (139 to 312) 15.4 (9.9 to 22.3) Counts, All Ages Rate Age-Age-Age-Age-Age-Age-Age-Age-Age-Age-				Both	1990	26,145	(20,502 to 31,784)	1298.3	(999.3 to 1653.9)
Country measure Stroke Type sex year Mean 95% Ul Mean 95% Ul					2013	33,314	(23,523 to 49,677)	980.8	(650.1 to 1647.8)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Female 1990 104 (67 to 155) 11.9 (7.5 to 17.8) Both 1990 219 (145 to 310) 13.1 (8.5 to 18.6) Hemorrhagic 2013 403 (268 to 551) 13.3 (8.9 to 18.2) Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) Stroke Male 1990 29 (18 to 44) 3 (1.9 to 4.5) Stroke Female 1990 29 (18 to 44) 3 (1.9 to 4.5) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)		YLDs	Ischemic Stroke	Male	1990	114	(69 to 169)	14.5	(8.9 to 21.3)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI 1 Female 1990 104 (67 to 155) 11.9 (7.5 to 17.8) 1 2013 186 (119 to 259) 11.5 (7.4 to 16.0) 2 Both 1990 219 (145 to 310) 13.1 (8.5 to 18.6) 3 2013 403 (268 to 551) 13.3 (8.9 to 18.2) Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)					2013	216	(139 to 312)	15.4	(9.9 to 22.3)
Country measure Stroke Type sex year Mean 95% UI Mean 95% UI Female 1990 104 (67 to 155) 11.9 (7.5 to 17.8) 1 2013 186 (119 to 259) 11.5 (7.4 to 16.0) Both 1990 219 (145 to 310) 13.1 (8.5 to 18.6) 2013 403 (268 to 551) 13.3 (8.9 to 18.2) Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)						Counts All	Δσρς	Rate	Per 100 000
Female 1990 104 (67 to 155) 11.9 (7.5 to 17.8) 2013 186 (119 to 259) 11.5 (7.4 to 16.0) Both 1990 219 (145 to 310) 13.1 (8.5 to 18.6) 2013 403 (268 to 551) 13.3 (8.9 to 18.2) Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)						Courtes, 7 til	ABC3	Hate	1 C1 100,000
2013 186 (119 to 259) 11.5 (7.4 to 16.0)						Courtes, 7 m	nges		
Both 1990 219 (145 to 310) 13.1 (8.5 to 18.6) 2013 403 (268 to 551) 13.3 (8.9 to 18.2) Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Stroke Type	sex	year			Age-	standardized
Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9)	Country	measure	Stroke Type			Mean	95% UI	Age- Mean	standardized 95% UI
Hemorrhagic Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Stroke Type		1990	Mean 104	95% UI (67 to 155)	Age- Mean 11.9	standardized 95% UI (7.5 to 17.8)
Stroke Male 1990 37 (23 to 55) 3.9 (2.5 to 5.9) 2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Stroke Type	Female	1990 2013	Mean 104 186	95% UI (67 to 155) (119 to 259)	Age- Mean 11.9 11.5	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0)
2013 66 (41 to 100) 3.9 (2.4 to 6.0) Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Stroke Type	Female	1990 2013 1990	Mean 104 186 219	95% UI (67 to 155) (119 to 259) (145 to 310)	Age- Mean 11.9 11.5 13.1	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6)
Female 1990 29 (18 to 44) 3 (1.9 to 4.5) 2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure		Female	1990 2013 1990	Mean 104 186 219	95% UI (67 to 155) (119 to 259) (145 to 310)	Age- Mean 11.9 11.5 13.1	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6)
2013 52 (32 to 76) 2.7 (1.7 to 3.9) Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Hemorrhagic	Female Both	1990 2013 1990 2013	Mean 104 186 219 403	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551)	Age-Mean 11.9 11.5 13.1 13.3	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6) (8.9 to 18.2)
Both 1990 66 (43 to 95) 3.4 (2.3 to 4.9) Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Hemorrhagic	Female Both	1990 2013 1990 2013	Mean 104 186 219 403	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551) (23 to 55)	Age-Mean 11.9 11.5 13.1 13.3	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6) (8.9 to 18.2) (2.5 to 5.9)
Both 2013 119 (78 to 169) 3.2 (2.1 to 4.6)	Country	measure	Hemorrhagic	Both Male	1990 2013 1990 2013 1990 2013	Mean 104 186 219 403 37 66	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551) (23 to 55) (41 to 100)	Age-Mean 11.9 11.5 13.1 13.3 3.9 3.9	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6) (8.9 to 18.2) (2.5 to 5.9) (2.4 to 6.0)
	Country	measure	Hemorrhagic	Both Male	1990 2013 1990 2013 1990 2013 1990	Mean 104 186 219 403 37 66 29	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551) (23 to 55) (41 to 100) (18 to 44)	Age-Mean 11.9 11.5 13.1 13.3 3.9 3.9 3.9	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6) (8.9 to 18.2) (2.5 to 5.9) (2.4 to 6.0) (1.9 to 4.5)
Tonga Deaths Ischemic Stroke Male 1990 8 (4 to 11) 44.9 (28.4 to 64.7)	Country	measure	Hemorrhagic	Both Male Female	1990 2013 1990 2013 1990 2013 1990 2013	Mean 104 186 219 403 37 66 29 52	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551) (23 to 55) (41 to 100) (18 to 44) (32 to 76)	Age-Mean 11.9 11.5 13.1 13.3 3.9 3.9 3.9 3.9	standardized 95% UI (7.5 to 17.8) (7.4 to 16.0) (8.5 to 18.6) (8.9 to 18.2) (2.5 to 5.9) (2.4 to 6.0) (1.9 to 4.5) (1.7 to 3.9)
	Country	measure	Hemorrhagic	Both Male Female Both	1990 2013 1990 2013 1990 2013 1990 2013 1990	Mean 104 186 219 403 37 66 29 52 66	95% UI (67 to 155) (119 to 259) (145 to 310) (268 to 551) (23 to 55) (41 to 100) (18 to 44) (32 to 76) (43 to 95)	Age-Mean 11.9 11.5 13.1 13.3 3.9 3.9 3.9 3.9 3.0	\$\text{standardized} \text{95\% UI} \tag{7.5 to 17.8} \tag{7.4 to 16.0} \tag{8.5 to 18.6} \tag{8.9 to 18.2} \tag{2.5 to 5.9} \tag{2.4 to 6.0} \tag{1.9 to 4.5} \tag{1.7 to 3.9} \tag{2.3 to 4.9}

			2013	12	(7 to 17)	46.9	(29.8 to 66.6)
		Female	1990	10	(7 to 13)	51.2	(36.2 to 68.6)
			2013	17	(10 to 24)	42.8	(25.4 to 62.6)
		Both	1990	18	(14 to 24)	48.6	(37.1 to 63.0)
			2013	29	(21 to 40)	45.1	(33.0 to 61.9)
	Hemorrhagic						
Deaths	Stroke	Male	1990	12	(8 to 16)	58.8	(39.0 to 80.1)
			2013	16	(10 to 22)	58.9	(37.5 to 80.7)
		Female	1990	15	(10 to 21)	68	(48.2 to 93.4)
			2013	21	(14 to 29)	54.4	(36.5 to 76.0)
		Both	1990	27	(21 to 34)	64.1	(49.5 to 80.6)
			2013	37	(27 to 47)	57.3	(41.7 to 72.9)
DALYs	Ischemic Stroke	Male	1990	148	(84 to 233)	690.5	(414.0 to 1025.5)
			2013	205	(124 to 300)	722.6	(446.5 to 1043.0)
		Female	1990	165	(110 to 232)	705.5	(492.1 to 963.5)
			2013	216	(126 to 329)	564.4	(326.1 to 861.4)
		Both	1990	313	(227 to 425)	701.2	(526.5 to 928.3)
			2013	422	(303 to 614)	639.7	(461.5 to 919.0)
	Hemorrhagic						
	Stroke	Male	1990	317	(210 to 436)	1189.8	(782.8 to 1635.7)
			2013	401	(259 to 560)	1222.7	(785.2 to 1688.7)
		Female	1990	321	(225 to 455)	1190.9	(853.0 to 1665.0)
			2013	357	(242 to 498)	920.7	(614.0 to 1291.2)
		Both	1990	638	(481 to 834)	1195.9	(907.3 to 1541.3)
			2013	759	(562 to 969)	1069.6	(792.0 to 1363.1)
YLDs	Ischemic Stroke	Male	1990	5	(3 to 7)	24.7	(15.7 to 35.2)
			2013	7	(4 to 10)	26.7	(17.2 to 38.9)
		Female	1990	4	(2 to 6)	18.9	(11.5 to 28.1)
			2013	7	(4 to 10)	20.5	(12.7 to 29.6)

			Both	1990	10	(6 to 13)	21.6	(14.4 to 29.8)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	14	(10 to 20)	23.1	(15.7 to 32.0)
		Hemorrhagic						
		Stroke	Male	1990	6	(4 to 9)	29.3	(19.1 to 41.9)
				2013	9	(6 to 13)	31.6	(20.2 to 44.4)
			Female	1990	5	(3 to 7)	20.8	(13.0 to 30.7)
				2013	8	(5 to 11)	22.1	(14.2 to 31.7)
			Both	1990	12	(8 to 16)	24.7	(17.0 to 33.9)
			Both	2013	17	(11 to 24)	26.2	(17.6 to 36.1)
Trinidad	Deaths	Ischemic Stroke	Male	1990	319	(258 to 374)	111.2	(89.5 to 131.4)
And				2013	449	(352 to 569)	92.2	(73.4 to 116.5)
Tobago			Female	1990	333	(253 to 383)	87.7	(66.8 to 100.1)
				2013	395	(304 to 536)	59.7	(46.1 to 80.5)
			Both	1990	653	(554 to 719)	98.1	(83.3 to 108.0)
				2013	845	(718 to 1,039)	73.7	(63.1 to 91.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	158	(128 to 202)	48.4	(38.9 to 63.5)
				2013	266	(170 to 341)	47.7	(30.3 to 61.8)
			Female	1990	181	(149 to 237)	44.8	(36.7 to 58.7)
				2013	227	(162 to 300)	32.8	(23.3 to 43.0)
			Both	1990	339	(298 to 414)	46.4	(40.4 to 57.6)
				2013	493	(366 to 586)	39.5	(29.1 to 47.0)
	DALYs	Ischemic Stroke	Male	1990	5,881	(4,786 to 6,864)	1774.1	(1442.8 to 2073.0
				2013	8,322	(6,529 to 10,689)	1448.2	(1149.8 to 1831.2
			Female	1990	5,204	(3,960 to 5,899)	1282.9	(981.5 to 1460.1)
				2013	5,621	(4,356 to 7,632)	804.9	(624.6 to 1094.6)

			Both	1990	11,085	(9,395 to 12,238)	1501.9	(1275.6 to 1650.5)
				2013	13,944	(11,630 to 17,123)	1092.7	(920.2 to 1338.6)
		Hemorrhagic						
		Stroke	Male	1990	3,970	(3,261 to 4,924)	1013.6	(831.8 to 1259.5)
				2013	6,328	(4,177 to 8,142)	975.1	(645.5 to 1243.5)
			Female	1990	4,211	(3,471 to 5,303)	943.9	(782.1 to 1199.9)
				2013	4,556	(3,381 to 6,049)	624.6	(461.5 to 823.8)
			Both	1990	8,181	(7,249 to 9,680)	973	(859.9 to 1165.1)
				2013	10,885	(8,316 to 13,015)	787.2	(602.1 to 940.4)
	YLDs	Ischemic Stroke	Male	1990	163	(106 to 231)	49.8	(32.5 to 71.4)
				2013	277	(174 to 404)	47.6	(29.8 to 70.5)
			Female	1990	130	(84 to 182)	32.3	(21.2 to 45.5)
				2013	232	(155 to 323)	32.9	(22.1 to 45.9)
			Both	1990	293	(193 to 407)	40	(26.5 to 55.5)
				2013	509	(340 to 707)	39.6	(26.4 to 55.0)
		Hemorrhagic						
		Stroke	Male	1990	59	(35 to 86)	16.1	(9.5 to 23.9)
		_	Male	1990	59 Counts, All	(35 to 86) Ages	16.1 Rate Age-	(9.5 to 23.9) Per 100,000 standardized
Country	measure	_	Male	1990 year			Rate	Per 100,000
Country	measure	Stroke			Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke		year	Counts, All Mean	Ages 95% UI	Rate Age- Mean	Per 100,000 standardized 95% UI
Country	measure	Stroke	sex	year 2013	Counts, All Mean 91	95% UI (56 to 133)	Rate Age- Mean 14.8	Per 100,000 standardized 95% UI (9.2 to 22.1)
Country	measure	Stroke	sex	year 2013 1990	Mean 91 44	95% UI (56 to 133) (26 to 66)	Rate Age- Mean 14.8 10.3	Per 100,000 standardized 95% UI (9.2 to 22.1) (6.2 to 15.7)
Country	measure	Stroke	sex Female	year 2013 1990 2013	Mean 91 44 79	95% UI (56 to 133) (26 to 66) (48 to 118)	Rate Age- Mean 14.8 10.3	Per 100,000 standardized 95% UI (9.2 to 22.1) (6.2 to 15.7) (6.7 to 16.5)
Country	measure	Stroke	sex Female Both	year 2013 1990 2013 1990	Mean 91 44 79 103	95% UI (56 to 133) (26 to 66) (48 to 118) (66 to 146)	Rate Age- Mean 14.8 10.3 11 12.9	Per 100,000 standardized 95% UI (9.2 to 22.1) (6.2 to 15.7) (6.7 to 16.5) (8.3 to 18.2)
		Stroke Stroke Type	sex Female Both Both	year 2013 1990 2013 1990 2013	Mean 91 44 79 103 170	95% UI (56 to 133) (26 to 66) (48 to 118) (66 to 146) (111 to 245)	Rate Age- Mean 14.8 10.3 11 12.9 12.7	Per 100,000 standardized 95% UI (9.2 to 22.1) (6.2 to 15.7) (6.7 to 16.5) (8.3 to 18.2) (8.3 to 18.2)
		Stroke Stroke Type	sex Female Both Both	year 2013 1990 2013 1990 2013 1990	Mean 91 44 79 103 170 927	95% UI (56 to 133) (26 to 66) (48 to 118) (66 to 146) (111 to 245) (681 to 1,232)	Rate Age- Mean 14.8 10.3 11 12.9 12.7 67.1	Per 100,000 standardized 95% UI (9.2 to 22.1) (6.2 to 15.7) (6.7 to 16.5) (8.3 to 18.2) (8.3 to 18.2) (49.2 to 88.8)

		Both	1990	1,967	(1,498 to 2,405)	71.4	(54.1 to 87.9)
			2013	4,234	(2,802 to 5,429)	55.8	(36.8 to 71.9)
	Hemorrhagic						
Deaths	Stroke	Male	1990	865	(634 to 1,190)	49.4	(34.9 to 68.9)
			2013	1,368	(837 to 2,141)	36.4	(21.9 to 57.4)
		Female	1990	1,108	(769 to 1,454)	64.7	(43.6 to 88.6)
			2013	1,967	(1,146 to 2,719)	44.8	(25.9 to 62.8)
		Both	1990	1,974	(1,571 to 2,494)	57.2	(44.2 to 74.4)
			2013	3,336	(2,214 to 4,488)	41.2	(26.8 to 55.6)
DALYs	Ischemic Stroke	Male	1990	16,321	(11,580 to 22,665)	886.6	(647.2 to 1180.5)
			2013	24,519	(15,641 to 36,844)	632.8	(408.7 to 929.5)
		Female	1990	17,362	(10,492 to 23,118)	951.5	(609.0 to 1233.8)
			2013	29,870	(14,707 to 42,811)	664.8	(332.5 to 953.2)
		Both	1990	33,684	(24,930 to 42,830)	919.4	(702.6 to 1130.7)
			2013	54,390	(35,319 to 72,353)	651.5	(428.1 to 857.4)
	Hemorrhagic						
	Stroke	Male	1990	24,379	(17,791 to 34,050)	926.1	(679.8 to 1278.6)
			2013	26,999	(17,359 to 41,175)	606.7	(383.4 to 935.5)
		Female	1990	29,574	(20,059 to 38,940)	1154.9	(810.6 to 1519.2)
			2013	32,883	(19,525 to 46,944)	676.7	(396.3 to 954.8)
		Both	1990	53,954	(42,614 to 67,529)	1040.9	(835.9 to 1302.1)
			2013	59,882	(42,085 to 80,163)	645.1	(445.9 to 862.8)
YLDs	Ischemic Stroke	Male	1990	268	(172 to 379)	11.9	(7.8 to 16.8)
			2013	456	(298 to 668)	9.7	(6.4 to 14.2)
		Female	1990	206	(130 to 294)	9.3	(5.9 to 13.3)
			2013	386	(250 to 560)	7.6	(4.9 to 11.0)
		Both	1990	474	(314 to 635)	10.6	(7.1 to 14.2)
			2013	842	(567 to 1,191)	8.6	(5.8 to 12.2)
	Hemorrhagic	Male	1990	36	(20 to 55)	1.4	(0.8 to 2.2)
	DALYS	DALYS Ischemic Stroke Hemorrhagic Stroke YLDs Ischemic Stroke	Hemorrhagic Stroke Male Female Both DALYS Ischemic Stroke Male Female Both Hemorrhagic Stroke Male Female Both Hemorrhagic Stroke Male Female Female Both Female Both Female Both Female Both	Hemorrhagic Stroke Male 1990 2013	Deaths	Deaths Hemorrhagic Stroke Male 1990 865 (634 to 1,190)	Deaths Stroke Male 1990 865 (634 to 1,190) 49.4

					T	1	1	T
		Stroke						
				2013	65	(36 to 102)	1.3	(0.8 to 2.1)
			Female	1990	41	(24 to 63)	1.7	(1.0 to 2.6)
				2013	75	(43 to 117)	1.4	(0.8 to 2.2)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	78	(49 to 112)	1.6	(1.0 to 2.3)
			Both	2013	141	(89 to 207)	1.4	(0.9 to 2.0)
Turkey	Deaths	Ischemic Stroke	Male	1990	8,293	(6,472 to 10,703)	91.4	(69.7 to 117.2)
				2013	9,935	(7,763 to 13,088)	49.3	(38.5 to 64.2)
			Female	1990	9,414	(7,374 to 12,779)	70.5	(54.6 to 95.2)
				2013	15,381	(11,484 to 18,592)	49.2	(36.5 to 59.6)
			Both	1990	17,707	(14,915 to 22,186)	78.9	(66.3 to 98.0)
				2013	25,317	(21,024 to 29,574)	49.5	(41.2 to 58.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	8,975	(6,969 to 11,274)	71	(52.3 to 90.8)
				2013	8,433	(6,465 to 11,106)	36.1	(27.1 to 47.7)
			Female	1990	9,283	(7,261 to 13,405)	59.3	(46.0 to 84.6)
				2013	11,227	(8,274 to 13,846)	34.5	(25.4 to 42.7)
			Both	1990	18,258	(15,472 to 22,493)	64.4	(54.1 to 80.8)
				2013	19,661	(16,170 to 23,318)	35.2	(28.9 to 42.0)
						(108,392 to		
	DALYs	Ischemic Stroke	Male	1990	141,289	189,194)	1208.6	(948.1 to 1572.6)
						(106,597 to		
				2013	137,842	182,213)	584.9	(455.4 to 767.6)
			Famala	1000	120 624	(98,100 to	0505	(672.0 to 1170.0)
			Female	1990	128,634	178,884) (121,721 to	858.5	(673.0 to 1179.6)
				2013	168,681	200,287)	522.5	(378.2 to 620.0)
				2013	100,001	200,207	322.3	(370.2 to 020.0)

9 (839.2 to 1285.3)
6 (452.0 to 642.4)
6 (1167.3 to 1887.8)
(491.1 to 827.6)
7 (908.7 to 1735.9)
(421.8 to 697.7)
(1116.1 to 1663.1)
(499.9 to 709.5)
(6.4 to 14.4)
(6.0 to 13.5)
(4.6 to 10.6)
(4.6 to 10.4)
(5.7 to 12.1)
(5.6 to 11.5)
te Per 100,000
e- standardized
n 95% UI
(0.9 to 2.5)
(0.9 to 2.2)
(1.0 to 2.5)
(0.8 to 2.1)
(1.0 to 2.3)
1

		I	1			1		
			Both	2013	969	(595 to 1,391)	1.5	(0.9 to 2.1)
Turkmenistan	Deaths	Ischemic Stroke	Male	1990	57	(44 to 79)	9.7	(7.3 to 13.2)
				2013	106	(64 to 173)	9.9	(5.9 to 15.8)
			Female	1990	63	(50 to 84)	6.7	(5.3 to 9.1)
				2013	92	(59 to 140)	6.1	(3.9 to 9.3)
			Both	1990	121	(102 to 155)	7.8	(6.6 to 10.2)
				2013	198	(142 to 276)	7.6	(5.5 to 10.5)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,228	(1,063 to 1,437)	194.5	(167.3 to 227.4)
				2013	2,246	(1,684 to 3,040)	196.6	(152.3 to 258.6)
			Female	1990	1,494	(1,334 to 1,772)	160	(143.9 to 191.0)
				2013	2,136	(1,686 to 2,922)	142.1	(113.2 to 192.0)
			Both	1990	2,723	(2,510 to 3,035)	174.1	(160.2 to 195.4)
				2013	4,382	(3,612 to 5,447)	166.1	(139.7 to 203.3)
	DALYs	Ischemic Stroke	Male	1990	1,471	(1,109 to 2,053)	184.1	(140.6 to 250.8)
				2013	2,663	(1,634 to 4,315)	185.1	(114.2 to 298.5)
			Female	1990	1,428	(1,125 to 1,865)	131.9	(105.5 to 171.5)
				2013	1,961	(1,296 to 3,022)	110	(73.5 to 169.9)
			Both	1990	2,900	(2,474 to 3,694)	152.5	(129.9 to 191.6)
				2013	4,624	(3,381 to 6,437)	142.4	(104.2 to 196.6)
		Hemorrhagic			•	, ,		,
		Stroke	Male	1990	31,672	(27,254 to 37,129)	3826.6	(3294.5 to 4469.0)
				2013	57,246	(41,173 to 78,434)	3831.2	(2841.2 to 5217.9)
			Female	1990	31,683	(27,521 to 37,144)	2939.4	(2569.4 to 3446.2)
				2013	43,151	(32,369 to 60,177)	2433	(1880.1 to 3368.2)
			Both	1990	63,356	(57,588 to 70,765)	3315.5	(3033.1 to 3683.5)
						(80,065 to		•
				2013	100,397	127,459)	3060.1	(2484.9 to 3860.3)
	YLDs	Ischemic Stroke	Male	1990	56	(36 to 83)	7.2	(4.6 to 10.6)

				2013	107	(68 to 151)	7.2	(4.6 to 10.3)
			Female	1990	70	(44 to 102)	6.8	(4.3 to 9.9)
				2013	114	(70 to 169)	6.4	(3.9 to 9.6)
			Both	1990	127	(85 to 180)	7	(4.7 to 9.8)
				2013	222	(144 to 313)	6.8	(4.4 to 9.6)
		Hemorrhagic						
		Stroke	Male	1990	89	(57 to 127)	9.4	(6.1 to 13.2)
				2013	157	(97 to 231)	8.8	(5.6 to 12.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	62	(39 to 92)	5.5	(3.4 to 8.2)
				2013	107	(65 to 161)	5.4	(3.2 to 8.1)
			Both	1990	152	(100 to 211)	7.2	(4.8 to 10.0)
			Both	2013	264	(172 to 374)	6.9	(4.5 to 9.9)
Uganda	Deaths	Ischemic Stroke	Male	1990	1,007	(187 to 1,745)	52.4	(9.6 to 89.7)
				2013	1,790	(641 to 2,755)	46.6	(17.1 to 69.9)
			Female	1990	1,496	(262 to 2,400)	69.6	(12.9 to 110.0)
				2013	2,752	(808 to 4,137)	63.1	(19.2 to 94.6)
			Both	1990	2,504	(587 to 3,837)	62.1	(15.1 to 94.6)
				2013	4,543	(1,749 to 6,467)	55.9	(21.9 to 78.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	1,941	(1,324 to 2,803)	74.2	(48.9 to 113.5)
				2013	2,572	(1,762 to 3,982)	53	(35.4 to 86.6)
			Female	1990	2,640	(1,220 to 3,638)	96.4	(42.7 to 143.4)
,				2013	3,455	(1,886 to 5,478)	67.4	(35.3 to 111.7)
			Both	1990	4,582	(2,930 to 6,124)	86.6	(53.0 to 123.5)
				2013	6,028	(4,201 to 9,046)	61	(41.1 to 96.9)
	DALYs	Ischemic Stroke	Male	1990	20,531	(4,306 to 36,113)	792.1	(156.3 to 1365.1)

				2013	32,092	(11,643 to 50,660)	670.7	(240.5 to 1035.6)
			Female	1990	25,999	(4,420 to 43,465)	967.4	(173.5 to 1559.8)
				2013	41,215	(11,914 to 63,228)	817.4	(236.9 to 1234.9)
			Both	1990	46,530	(11,390 to 72,178)	891.3	(216.0 to 1362.6)
						(28,608 to		
				2013	73,308	105,973)	753.6	(296.0 to 1077.5)
		Hemorrhagic						
		Stroke	Male	1990	55,475	(38,379 to 78,116)	1536	(1044.1 to 2216.0)
				2013	65,860	(45,695 to 97,623)	1006.7	(689.7 to 1555.6)
			Female	1990	64,329	(30,725 to 84,771)	1802.7	(832.3 to 2475.6)
						(42,658 to		
				2013	75,620	112,292)	1157.7	(639.3 to 1758.5)
						(80,354 to		(
			Both	1990	119,805	155,793)	1684.8	(1077.5 to 2238.5)
				2012	141 400	(101,863 to	1000.6	/770 C to 1505 0\
	V/I D	lanka ata Guarka	0.0.1.	2013	141,480	196,734)	1090.6	(770.6 to 1595.8)
	YLDs	Ischemic Stroke	Male	1990	366	(236 to 527)	10.1	(6.4 to 14.6)
				2013	751	(479 to 1,113)	10.7	(6.8 to 16.1)
			Female	1990	278	(175 to 397)	7.3	(4.6 to 10.5)
				2013	542	(347 to 815)	7.2	(4.6 to 10.9)
			Both	1990	645	(432 to 897)	8.6	(5.8 to 12.0)
				2013	1,293	(844 to 1,847)	8.9	(5.8 to 12.7)
		Hemorrhagic						
		Stroke	Male	1990	131	(78 to 196)	3	(1.8 to 4.3)
				2013	280	(174 to 430)	3.1	(1.9 to 4.7)
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Female	1990	86	(51 to 130)	1.9	(1.2 to 2.8)
				2013	175	(104 to 261)	1.9	(1.2 to 2.8)

			Both	1990	217	(138 to 311)	2.4	(1.5 to 3.4)
			Both	2013	455	(288 to 665)	2.5	(1.6 to 3.6)
Ukraine	Deaths	Ischemic Stroke	Male	1990	26,657	(21,127 to 31,815)	148.4	(117.5 to 175.0)
				2013	32,153	(26,584 to 38,997)	128.7	(106.7 to 154.5)
			Female	1990	58,127	(44,220 to 66,563)	133.7	(102.1 to 153.4)
				2013	50,499	(43,308 to 63,804)	92.5	(79.3 to 117.7)
			Both	1990	84,784	(69,711 to 95,779)	139.5	(115.0 to 157.5)
	Deaths			2013	82,652	(73,718 to 96,167)	105.9	(94.1 to 123.7)
		Hemorrhagic						,
		Stroke	Male	1990	12,355	(8,358 to 16,377)	57.3	(38.1 to 76.1)
				2013	12,344	(8,365 to 17,382)	48	(32.2 to 68.5)
			Female	1990	21,120	(13,628 to 25,730)	50.4	(33.0 to 61.5)
				2013	15,364	(11,634 to 24,582)	32.3	(25.0 to 50.0)
			Both	1990	33,475	(23,908 to 40,061)	53.6	(38.3 to 64.3)
				2013	27,708	(21,692 to 39,950)	38.8	(30.4 to 54.9)
						(352,678 to		
	DALYs	Ischemic Stroke	Male	1990	447,765	550,844)	2163.5	(1722.4 to 2616.7)
						(418,968 to		
				2013	512,002	623,398)	1953.4	(1615.1 to 2356.7)
			5 1-	1000	724 604	(552,224 to	1601.6	(4200 C to 4072 4)
			Female	1990	724,691	847,016)	1681.6	(1280.6 to 1972.1)
				2013	578,674	(500,777 to 727,450)	1139.7	(973.6 to 1429.9)
				2013	370,074	(973,550 to	1133.7	(373.0 to 1423.3)
			Both	1990	1,172,456	1,362,689)	1865.6	(1560.6 to 2163.2)
						(968,704 to		,
				2013	1,090,677	1,266,969)	1452.4	(1290.2 to 1688.9)
		Hemorrhagic				(205,699 to		
		Stroke	Male	1990	295,846	384,231)	1216.7	(848.3 to 1588.5)
				2013	285,091	(198,637 to	1082.5	(762.8 to 1482.3)

						387,814)		
						(253,841 to		
			Female	1990	376,371	449,182)	951.5	(656.3 to 1125.9)
						(203,614 to		
				2013	257,577	384,362)	623.3	(503.0 to 905.9)
						(501,096 to		
			Both	1990	672,217	799,148)	1067.8	(803.2 to 1266.6)
						(434,914 to		
				2013	542,669	733,023)	820.3	(660.8 to 1078.8)
	YLDs	Ischemic Stroke	Male	1990	11,221	(7,497 to 15,929)	46.7	(31.6 to 65.0)
				2013	13,848	(9,129 to 19,470)	52.7	(34.5 to 73.8)
			Female	1990	11,159	(7,446 to 15,623)	29.2	(19.5 to 40.4)
				2013	11,456	(7,707 to 16,206)	28	(18.5 to 39.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
			Both	1990	22,380	(15,114 to 30,441)	36.2	(24.5 to 49.0)
				2013	25,305	(17,532 to 34,184)	38.1	(26.4 to 51.5)
		Hemorrhagic						
		Stroke	Male	1990	691	(379 to 1,199)	2.8	(1.5 to 4.7)
				2013	723	(371 to 1,338)	2.8	(1.5 to 5.0)
			Female	1990	709	(378 to 1,207)	2	(1.1 to 3.4)
				2013	677	(390 to 1,095)	1.9	(1.1 to 3.0)
			Both	1990	1,401	(831 to 2,179)	2.3	(1.4 to 3.6)
			Both	2013	1,401	(835 to 2,113)	2.3	(1.4 to 3.4)
Unitad	Deaths	Ischemic Stroke	Male	1990	120	(72 to 199)	85.6	(56.2 to 130.5)
United Arab				2013	217	(121 to 334)	65.1	(39.7 to 90.7)
Emirates			Female	1990	100	(44 to 146)	101.8	(47.5 to 145.5)
Limates			1 Ciriaic	2013	97	(38 to 147)	63.3	(28.2 to 91.4)
			Doth			,		, , , , , , , , , , , , , , , , , , , ,
			Both	1990	220	(137 to 300)	93.9	(59.7 to 121.5)

			2013	315	(186 to 441)	65	(42.4 to 84.6)
	Hemorrhagic						
Deaths	Stroke	Male	1990	154	(82 to 242)	55.6	(26.9 to 89.6)
			2013	368	(224 to 578)	43.8	(25.4 to 69.8)
		Female	1990	103	(36 to 178)	76.5	(26.4 to 131.1)
			2013	146	(70 to 241)	46.4	(23.4 to 73.8)
		Both	1990	258	(157 to 377)	65.5	(35.4 to 95.8)
			2013	514	(354 to 749)	45.2	(31.2 to 67.5)
DALYs	Ischemic Stroke	Male	1990	3,198	(1,799 to 5,562)	1194.9	(751.4 to 1912.6)
			2013	6,357	(3,329 to 10,332)	836.2	(505.7 to 1176.8)
		Female	1990	1,776	(691 to 2,757)	1376.7	(604.5 to 2025.8)
			2013	2,072	(739 to 3,326)	732.7	(320.0 to 1091.6)
		Both	1990	4,974	(2,912 to 7,320)	1282.5	(805.0 to 1705.8)
			2013	8,430	(4,688 to 12,792)	803.8	(497.2 to 1077.6)
	Hemorrhagic						
	Stroke	Male	1990	5,805	(3,233 to 9,130)	1050.1	(547.8 to 1679.7)
			2013	15,027	(9,059 to 24,037)	740	(446.4 to 1176.0)
		Female	1990	3,061	(1,114 to 5,258)	1388.9	(473.5 to 2412.1)
			2013	5,112	(2,449 to 8,595)	713.9	(351.2 to 1154.3)
		Both	1990	8,866	(5,540 to 12,778)	1188.5	(684.3 to 1737.1)
			2013	20,139	(13,771 to 29,630)	734.8	(510.3 to 1088.3)
YLDs	Ischemic Stroke	Male	1990	49	(31 to 72)	9.4	(6.1 to 13.0)
			2013	239	(142 to 347)	9.3	(5.9 to 12.8)
		Female	1990	14	(9 to 21)	7	(4.5 to 10.1)
			2013	71	(45 to 106)	7.9	(5.1 to 11.4)
		Both	1990	64	(41 to 90)	8.5	(5.7 to 11.5)
			2013	311	(195 to 447)	8.8	(5.8 to 12.0)
				Counts, All	Ages	Rate	Per 100,000

							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	9	(5 to 15)	1.2	(0.7 to 1.8)
				2013	49	(24 to 83)	1.3	(0.7 to 2.0)
			Female	1990	3	(1 to 5)	1.2	(0.7 to 2.1)
				2013	18	(9 to 30)	1.3	(0.8 to 2.2)
			Both	1990	12	(7 to 19)	1.2	(0.7 to 1.8)
			Both	2013	67	(39 to 106)	1.3	(0.8 to 1.9)
United	Deaths	Ischemic Stroke	Male	1990	17,411	(14,055 to 21,066)	54.6	(44.0 to 65.8)
Kingdom				2013	13,099	(9,601 to 16,846)	25.7	(18.9 to 33.0)
			Female	1990	31,017	(22,969 to 34,287)	50.7	(37.4 to 56.0)
				2013	21,170	(18,009 to 28,522)	26.9	(23.0 to 36.5)
			Both	1990	48,429	(39,939 to 53,162)	52.9	(43.6 to 58.1)
				2013	34,269	(29,644 to 42,655)	26.7	(23.1 to 33.1)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	10,428	(8,380 to 13,716)	32	(25.7 to 41.9)
				2013	9,102	(5,552 to 12,013)	18.7	(11.6 to 24.4)
			Female	1990	17,079	(13,066 to 21,248)	31.6	(24.5 to 39.4)
				2013	13,773	(9,921 to 17,873)	19.5	(14.4 to 25.2)
			Both	1990	27,508	(23,766 to 33,273)	32.1	(27.9 to 38.7)
				2013	22,876	(17,286 to 26,965)	19.3	(14.8 to 22.7)
						(210,272 to		
	DALYs	Ischemic Stroke	Male	1990	256,351	307,474)	766.5	(627.4 to 918.3)
						(136,249 to		(22.2.2
				2013	175,279	215,136)	364.5	(284.0 to 446.5)
			Female	1990	329,188	(248,692 to 362,174)	596.5	(451.9 to 661.6)
			· cinaic	1330	323,100	(169,184 to	330.3	(131.3 to 001.0)
				2013	197,881	256,980)	296.2	(252.2 to 385.5)

						(496,562 to		
			Both	1990	585,540	645,218)	672	(573.2 to 742.0)
						(321,750 to		
				2013	373,161	447,585)	329.7	(281.7 to 393.1)
		Hemorrhagic				(158,263 to		
		Stroke	Male	1990	193,014	245,458)	584.9	(480.1 to 740.1)
						(93,344 to		
				2013	136,343	170,874)	305.3	(212.0 to 378.9)
						(187,586 to		
			Female	1990	238,120	294,320)	532.3	(428.2 to 661.0)
						(121,176 to		
				2013	153,476	193,959)	276.4	(224.3 to 347.6)
						(382,222 to		
			Both	1990	431,135	515,658)	556.5	(494.8 to 662.9)
						(231,307 to		
				2013	289,820	331,741)	291	(235.7 to 331.9)
	YLDs	Ischemic Stroke	Male	1990	28,636	(19,631 to 38,572)	84.8	(58.3 to 114.0)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	o				0=0(N 4	050/ 111
	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
	measure	Stroke Type	sex	year 2013	Mean 45,284	95% UI (31,590 to 59,273)	98.1	(68.4 to 128.3)
	illeasure	Stroke Type	Female					
	measure	Stroke Type		2013	45,284	(31,590 to 59,273)	98.1	(68.4 to 128.3)
	measure	Stroke Type		2013 1990	45,284 30,037	(31,590 to 59,273) (21,126 to 40,006)	98.1 62.6	(68.4 to 128.3) (43.7 to 83.5)
	measure	Stroke Type	Female	2013 1990 2013	45,284 30,037 38,473	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704)	98.1 62.6 68.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9)
	measure	Stroke Type	Female	2013 1990 2013	45,284 30,037 38,473	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169)	98.1 62.6 68.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9)
	measure	Hemorrhagic	Female	2013 1990 2013 1990	45,284 30,037 38,473 58,674	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169) (58,977 to	98.1 62.6 68.2 72.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9) (50.3 to 96.3)
	measure		Female	2013 1990 2013 1990	45,284 30,037 38,473 58,674	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169) (58,977 to	98.1 62.6 68.2 72.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9) (50.3 to 96.3)
	Illeasure	Hemorrhagic	Female Both	2013 1990 2013 1990 2013	45,284 30,037 38,473 58,674 83,757	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169) (58,977 to 110,025)	98.1 62.6 68.2 72.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9) (50.3 to 96.3) (57.9 to 108.3)
	Illeasure	Hemorrhagic	Female Both	2013 1990 2013 1990 2013 1990	45,284 30,037 38,473 58,674 83,757 6,356	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169) (58,977 to 110,025) (4,445 to 8,526)	98.1 62.6 68.2 72.2 82.2	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9) (50.3 to 96.3) (57.9 to 108.3) (13.4 to 25.7)
	Illeasure	Hemorrhagic	Female Both Male	2013 1990 2013 1990 2013 1990 2013	45,284 30,037 38,473 58,674 83,757 6,356 9,635	(31,590 to 59,273) (21,126 to 40,006) (27,208 to 50,704) (40,972 to 78,169) (58,977 to 110,025) (4,445 to 8,526) (6,604 to 12,967)	98.1 62.6 68.2 72.2 82.2 19.2 21.6	(68.4 to 128.3) (43.7 to 83.5) (48.1 to 89.9) (50.3 to 96.3) (57.9 to 108.3) (13.4 to 25.7) (14.8 to 29.2)

			Both	1990	12,580	(8,733 to 16,714)	16.3	(11.3 to 21.7)
			Both	2013	17,722	(12,330 to 23,550)	18.4	(12.8 to 24.4)
United	Deaths	Ischemic Stroke	Male	1990	34,042	(26,947 to 43,130)	30.4	(24.0 to 38.6)
States				2013	34,261	(24,840 to 44,793)	17.4	(12.7 to 22.7)
			Female	1990	51,546	(39,805 to 66,103)	25.7	(19.9 to 32.9)
				2013	52,552	(38,834 to 74,671)	17.2	(12.8 to 24.0)
						(72,854 to		· · · · · · · · · · · · · · · · · · ·
			Both	1990	85,588	102,433)	27.7	(23.6 to 33.2)
						(70,233 to		
				2013	86,814	110,540)	17.5	(14.2 to 22.0)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	27,563	(21,808 to 35,788)	23.7	(18.7 to 30.9)
				2013	33,966	(22,876 to 43,736)	17.3	(11.8 to 22.3)
			Female	1990	36,584	(29,748 to 49,687)	20.5	(17.0 to 27.8)
				2013	43,452	(29,990 to 57,032)	15.9	(11.4 to 20.3)
			Both	1990	64,148	(56,359 to 81,232)	21.9	(19.2 to 27.5)
				2013	77,418	(56,812 to 92,237)	16.6	(12.3 to 19.6)
						(506,353 to		
	DALYs	Ischemic Stroke	Male	1990	614,425	754,159)	520.3	(428.7 to 639.6)
						(580,591 to		
				2013	729,573	888,987)	375.7	(299.7 to 457.8)
						(555,553 to		
			Female	1990	678,586	826,290)	376.7	(310.7 to 459.8)
						(648,311 to		(222.2)
				2013	812,046	1,013,324)	325.2	(260.3 to 404.4)
			Dotle	1000	1 202 011	(1,125,068 to	427.6	(270 7 to 544 4)
			Both	1990	1,293,011	1,507,585)	437.6	(379.7 to 511.4)
				2013	1,541,619	(1,272,995 to 1,854,470)	348.5	(287.0 to 421.9)
		Hemorrhagic		2013	1,341,013	(482,248 to	340.3	(207.0 (0 421.9)
		Stroke	Male	1990	584,795	727,997)	482	(397.0 to 601.6)

						(466,025 to		
				2013	629,382	776,903)	329.6	(244.7 to 405.6)
								,
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
						(523,943 to		
			Female	1990	614,995	808,836)	399	(337.7 to 525.1)
						(512,565 to		
				2013	666,125	807,151)	295.5	(227.7 to 350.3)
						(1,074,283 to		
			Both	1990	1,199,791	1,439,026)	435.6	(390.3 to 521.8)
						(1,036,041 to		
				2013	1,295,507	1,481,442)	311.2	(249.6 to 355.2)
						(101,969 to		
	YLDs	Ischemic Stroke	Male	1990	154,842	215,960)	129.9	(86.1 to 181.1)
						(239,197 to		
				2013	341,067	449,654)	175.8	(123.1 to 231.8)
						(113,293 to		
			Female	1990	175,958	249,169)	103.2	(65.9 to 146.9)
				2010	.=	(264,899 to	460.0	(117.0)
				2013	378,932	509,498)	163.8	(115.0 to 220.8)
			Dath	1000	220.004	(222,886 to	1111	(7C 0 to 4FC 7)
			Both	1990	330,801	453,557)	114.4	(76.8 to 156.7)
				2013	710 000	(511,420 to 965,127)	169	(110 9 to 226 0)
		Hemorrhagic		2013	719,999	903,127)	109	(119.8 to 226.9)
		Stroke	Male	1990	23,632	(15,006 to 34,670)	19.5	(12.4 to 28.6)
		Stille	iviale		•	, , ,		,
				2013	53,721	(35,922 to 74,177)	29	(19.4 to 39.8)
			Female	1990	31,655	(20,075 to 46,696)	19.8	(12.5 to 29.3)
				2012	02.275	(57,455 to	20.2	(27.0 + - 52.0)
				2013	83,275	113,967)	39.2	(27.0 to 53.8)

			Both	1990	55,288	(37,616 to 78,894)	19.6	(13.3 to 27.9)
						(96,067 to		
			Both	2013	136,997	184,826)	34.2	(23.9 to 46.2)
Uruguay	Deaths	Ischemic Stroke	Male	1990	1,209	(976 to 1,426)	89.5	(72.5 to 105.8)
				2013	1,085	(861 to 1,383)	57.3	(45.5 to 73.2)
			Female	1990	1,601	(1,325 to 1,925)	76.3	(62.9 to 91.5)
				2013	1,545	(1,181 to 2,261)	42.5	(33.0 to 62.1)
			Both	1990	2,810	(2,457 to 3,145)	82.3	(71.8 to 92.5)
				2013	2,631	(2,197 to 3,494)	48.6	(40.5 to 64.2)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	501	(402 to 683)	35.2	(28.1 to 48.4)
				2013	641	(400 to 834)	34.5	(21.5 to 44.8)
			Female	1990	607	(510 to 885)	30.6	(25.9 to 44.3)
				2013	840	(524 to 1,090)	26.9	(17.2 to 34.3)
			Both	1990	1,108	(947 to 1,472)	32.7	(27.9 to 43.3)
				2013	1,481	(1,033 to 1,781)	30.2	(21.3 to 36.2)
	DALYs	Ischemic Stroke	Male	1990	19,196	(15,525 to 22,700)	1318.6	(1068.8 to 1556.9)
				2013	14,572	(11,603 to 18,834)	783.4	(621.8 to 1013.5)
			Female	1990	19,097	(15,752 to 22,553)	942.2	(771.0 to 1109.9)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	14,143	(11,147 to 20,539)	457.6	(358.1 to 673.5)
			Both	1990	38,293	(33,479 to 42,723)	1108.8	(970.6 to 1236.1)
				2013	28,715	(24,086 to 37,392)	594.9	(496.7 to 775.8)
		Hemorrhagic						
		Stroke	Male	1990	10,894	(8,885 to 14,255)	727.8	(595.3 to 957.8)
				2013	11,786	(7,869 to 15,278)	650.1	(435.7 to 842.8)
			Female	1990	11,014	(9,621 to 15,555)	602.6	(529.0 to 842.1)

				2013	11,272	(7,402 to 13,967)	458.2	(308.1 to 563.7)
			Both	1990	21,908	(18,989 to 28,082)	658.5	(572.6 to 842.7)
				2013	23,058	(16,826 to 27,560)	542.7	(397.3 to 651.1)
	YLDs	Ischemic Stroke	Male	1990	512	(335 to 717)	33.8	(22.5 to 47.6)
				2013	561	(366 to 808)	30.8	(20.1 to 44.2)
			Female	1990	380	(250 to 534)	19.7	(13.2 to 27.6)
				2013	450	(288 to 622)	18.4	(12.0 to 25.5)
			Both	1990	892	(603 to 1,218)	25.9	(17.5 to 35.4)
				2013	1,011	(688 to 1,361)	23.7	(16.1 to 32.0)
		Hemorrhagic						
		Stroke	Male	1990	167	(104 to 239)	11.1	(7.0 to 15.9)
				2013	220	(141 to 315)	12.3	(7.9 to 17.5)
			Female	1990	105	(66 to 156)	5.7	(3.6 to 8.5)
				2013	137	(87 to 196)	6.1	(4.0 to 8.7)
			Both	1990	272	(177 to 377)	8.1	(5.3 to 11.2)
			Both	2013	357	(239 to 491)	8.8	(6.0 to 12.1)
Uzbekistan	Deaths	Ischemic Stroke	Male	1990	3,910	(3,107 to 4,783)	113.5	(90.5 to 139.1)
				2013	7,420	(5,603 to 9,574)	123.5	(95.1 to 156.3)
			Female	1990	5,051	(4,301 to 6,130)	92.2	(78.4 to 111.4)
				2013	7,549	(5,543 to 9,554)	93.7	(68.7 to 118.8)
			Both	1990	8,961	(7,856 to 10,457)	100.9	(88.3 to 117.6)
				2013	14,970	(12,470 to 17,906)	107.4	(89.3 to 126.8)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	2,944	(2,268 to 3,729)	75.7	(57.5 to 96.6)
				2013	5,466	(3,912 to 7,391)	80.2	(58.1 to 106.8)
			Female	1990	3,680	(2,946 to 4,676)	64.7	(51.5 to 82.5)
				2013	5,236	(3,656 to 7,254)	61.2	(42.8 to 83.1)
			Both	1990	6,625	(5,606 to 7,973)	69.1	(58.3 to 84.1)
				2013	10,703	(8,705 to 13,448)	69.9	(56.4 to 88.2)

	DALYs	Ischemic Stroke	Male	1990	73,913	(58,709 to 91,686)	1839	(1465.9 to 2264.4)
						(104,963 to		
				2013	144,434	193,337)	1979	(1471.8 to 2578.4)
			Female	1990	73,219	(61,008 to 89,756)	1293.8	(1082.7 to 1586.0)
						(78,687 to		
				2013	106,255	140,602)	1239.1	(915.2 to 1609.4)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
·		•		•		(126,398 to		
			Both	1990	147,133	173,453)	1524.9	(1319.6 to 1787.9)
						(204,664 to		
				2013	250,690	308,290)	1580.5	(1297.2 to 1916.4)
		Hemorrhagic						
		Stroke	Male	1990	70,205	(55,703 to 87,549)	1517	(1189.3 to 1915.5)
						(94,005 to		
				2013	134,038	186,930)	1585.3	(1133.0 to 2159.0)
			Female	1990	75,122	(60,074 to 90,600)	1255.8	(1003.1 to 1519.1)
						(74,504 to		
				2013	104,916	149,646)	1095.3	(781.0 to 1520.8)
						(124,012 to		
			Both	1990	145,327	170,168)	1365.9	(1162.6 to 1603.6)
						(190,173 to		
				2013	238,955	304,975)	1323	(1059.2 to 1679.1)
	YLDs	Ischemic Stroke	Male	1990	406	(258 to 591)	8.8	(5.6 to 12.8)
				2013	752	(476 to 1,074)	8.9	(5.7 to 12.7)
			Female	1990	478	(301 to 678)	8	(5.1 to 11.5)
				2013	754	(475 to 1,135)	7.7	(4.9 to 11.5)
			Both	1990	884	(579 to 1,237)	8.4	(5.5 to 11.7)
				2013	1,506	(1,007 to 2,155)	8.3	(5.6 to 11.8)

			,					
		Hemorrhagic						
		Stroke	Male	1990	442	(276 to 642)	8	(5.1 to 11.6)
				2013	782	(498 to 1,112)	7.8	(5.0 to 11.2)
			Female	1990	298	(188 to 436)	4.6	(2.8 to 6.8)
				2013	498	(313 to 738)	4.6	(2.8 to 6.9)
			Both	1990	741	(481 to 1,036)	6.1	(4.0 to 8.6)
			Both	2013	1,281	(845 to 1,805)	6.1	(4.1 to 8.7)
Vanuatu	Deaths	Ischemic Stroke	Male	1990	24	(12 to 35)	81.9	(41.6 to 118.4)
				2013	39	(22 to 56)	81.5	(48.4 to 112.8)
			Female	1990	11	(7 to 18)	51.1	(31.8 to 79.3)
				2013	29	(14 to 48)	61.8	(33.7 to 98.3)
			Both	1990	36	(21 to 52)	68.3	(42.6 to 96.1)
				2013	68	(39 to 100)	71.6	(43.5 to 102.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	42	(26 to 61)	123.4	(78.7 to 176.7)
				2013	78	(49 to 117)	128.4	(82.3 to 185.4)
			Female	1990	21	(13 to 33)	80.5	(50.8 to 125.8)
				2013	54	(31 to 85)	96.8	(58.0 to 148.2)
			Both	1990	63	(41 to 90)	103.9	(69.5 to 145.2)
				2013	133	(85 to 197)	112.6	(74.7 to 158.4)
	DALYs	Ischemic Stroke	Male	1990	487	(237 to 784)	1426.1	(715.1 to 2204.5)
				2013	836	(445 to 1,288)	1353.1	(767.3 to 1965.3)
			Female	1990	236	(131 to 377)	849.6	(498.0 to 1330.1)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	585	(261 to 997)	967	(479.7 to 1584.3)
			Both	1990	723	(403 to 1,113)	1167.1	(686.1 to 1733.6)
				2013	1,421	(758 to 2,186)	1159.8	(662.4 to 1693.7)

		Hana a mila a di	 				ı I	
		Hemorrhagic Stroke	Male	1990	1,184	(728 to 1,764)	2885.7	(1792.4 to 4286.1)
		Stroke	iviaic	2013	2,322	(1,429 to 3,637)	2946.2	(1837.7 to 4465.9)
			Famala		558		1682.9	· , , , , , , , , , , , , , , , , , , ,
			Female	1990		(334 to 911)		(1034.5 to 2678.3)
				2013	1,425	(804 to 2,338)	1956.8	(1135.9 to 3096.1)
			Both	1990	1,742	(1,108 to 2,588)	2325.8	(1506.0 to 3400.9)
				2013	3,748	(2,348 to 5,730)	2451.1	(1553.2 to 3677.7)
	YLDs	Ischemic Stroke	Male	1990	8	(5 to 12)	27	(17.8 to 38.2)
				2013	15	(10 to 21)	26.7	(17.5 to 37.9)
			Female	1990	5	(3 to 7)	19.5	(12.8 to 27.7)
				2013	12	(7 to 17)	20.6	(13.2 to 30.0)
			Both	1990	13	(9 to 19)	23.7	(16.0 to 32.6)
				2013	27	(18 to 37)	23.7	(16.1 to 32.5)
		Hemorrhagic						
		Stroke	Male	1990	11	(7 to 15)	31.3	(20.3 to 43.7)
				2013	20	(13 to 29)	31.8	(20.5 to 45.7)
			Female	1990	6	(4 to 8)	21.1	(14.0 to 29.1)
				2013	14	(9 to 20)	22.1	(14.0 to 31.1)
			Both	1990	17	(11 to 23)	26.7	(17.7 to 36.5)
			Both	2013	34	(22 to 48)	26.9	(17.6 to 37.0)
Venezuela	Deaths	Ischemic Stroke	Male	1990	1,843	(1,124 to 2,390)	58.7	(35.8 to 76.5)
				2013	1,903	(1,246 to 3,221)	23.4	(15.0 to 39.5)
			Female	1990	1,999	(1,192 to 2,323)	53.6	(32.3 to 62.4)
				2013	1,970	(1,371 to 3,329)	20.5	(14.2 to 35.0)
			Both	1990	3,843	(2,515 to 4,629)	56.1	(37.0 to 67.5)
				2013	3,874	(2,776 to 6,195)	22	(15.6 to 35.1)
		Hemorrhagic			·	,		· ,
	Deaths	Stroke	Male	1990	1,385	(984 to 2,174)	36.7	(24.7 to 59.8)
				2013	4,554	(3,034 to 5,822)	50.5	(32.9 to 65.2)

			Female	1990	1,533	(1,220 to 2,521)	35.1	(26.7 to 61.0)
				2013	4,562	(2,885 to 5,563)	44.6	(27.4 to 55.0)
			Both	1990	2,919	(2,250 to 4,450)	35.8	(26.5 to 57.8)
				2013	9,117	(6,366 to 10,924)	47.6	(32.1 to 57.6)
	DALYs	Ischemic Stroke	Male	1990	35,516	(21,096 to 46,077)	867.7	(526.2 to 1126.5)
				2013	31,490	(21,551 to 52,164)	323.5	(216.2 to 539.2)
			Female	1990	32,819	(18,630 to 38,269)	717.6	(421.2 to 833.7)
				2013	25,031	(17,998 to 43,950)	235.8	(169.2 to 407.6)
			Both	1990	68,335	(43,032 to 83,174)	788.7	(508.3 to 953.6)
				2013	56,522	(41,900 to 91,203)	277.8	(204.1 to 443.7)
<u>'</u>					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country m	easure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
		Hemorrhagic						
		Stroke	Male	1990	36,171	(27,233 to 53,559)	714.4	(518.6 to 1097.4)
						(69,042 to		
				2013	97,820	124,233)	886.7	(617.5 to 1123.3)
			Female	1990	37,585	(32,146 to 56,397)	672.5	(556.4 to 1052.6)
				2013	79,826	(56,359 to 95,528)	678.3	(468.7 to 817.0)
						(60,454 to		
			Both	1990	73,757	105,427)	691.3	(546.8 to 1019.7)
						(133,781 to		
				2013	177,647	209,597)	779.2	(574.5 to 923.9)
,	YLDs	Ischemic Stroke	Male	1990	326	(198 to 487)	7.7	(4.8 to 11.5)
				2013	773	(476 to 1,146)	7.5	(4.6 to 11.2)
			Female	1990	277	(171 to 413)	5.9	(3.6 to 8.8)
				2013	677	(425 to 1,005)	6	(3.7 to 8.9)
			Both	1990	603	(395 to 830)	6.8	(4.4 to 9.2)
				2013	1,451	(945 to 2,078)	6.7	(4.3 to 9.6)
		Hemorrhagic	Male	1990	992	(649 to 1,426)	19.9	(13.1 to 28.5)

		Stroke						
				2013	2,317	(1,481 to 3,292)	20.5	(13.3 to 29.0)
			Female	1990	693	(463 to 1,001)	12.9	(8.5 to 18.7)
				2013	1,564	(979 to 2,249)	12.8	(8.0 to 18.2)
			Both	1990	1,686	(1,134 to 2,295)	16.2	(11.1 to 22.2)
			Both	2013	3,882	(2,582 to 5,394)	16.4	(10.9 to 22.8)
Vietnam	Deaths	Ischemic Stroke	Male	1990	20,525	(12,432 to 26,365)	143.8	(86.7 to 182.7)
				2013	35,001	(23,192 to 45,848)	140.6	(93.4 to 183.8)
			Female	1990	29,102	(19,969 to 38,427)	122.5	(83.9 to 161.8)
				2013	53,074	(35,862 to 64,983)	107.6	(72.6 to 133.1)
			Both	1990	49,627	(36,418 to 60,366)	132.4	(98.1 to 161.0)
					,	(62,709 to		,
				2013	88,075	105,439)	120.8	(85.5 to 145.4)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	26,858	(19,802 to 35,337)	164.7	(120.9 to 218.7)
				2013	28,145	(19,795 to 40,024)	97.8	(68.2 to 141.2)
			Female	1990	28,145	(19,548 to 37,955)	119.7	(83.3 to 161.2)
				2013	29,513	(21,296 to 45,127)	63	(45.3 to 95.4)
			Both	1990	55,003	(43,896 to 67,359)	140.2	(111.4 to 172.6)
				2013	57,658	(44,919 to 79,896)	77.9	(60.5 to 108.3)
						(213,082 to		
	DALYs	Ischemic Stroke	Male	1990	368,688	484,990)	2277.1	(1341.9 to 2953.6)
						(329,538 to		
				2013	507,828	703,840)	1882	(1229.4 to 2572.7)
			F	4000	202.422	(254,735 to	4625.3	/4076 F L. 2400 F\
			Female	1990	383,433	515,318)	1625.3	(1076.5 to 2190.5)
					Counts All	Λ σ ο ο	Doto	Dov. 100,000
					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI

1				1	1		
					(340,216 to		
			2013	514,796	657,908)	1156.5	(763.2 to 1517.9)
					(540,071 to		
		Both	1990	752,121	929,524)	1914.3	(1380.3 to 2355.7)
					(722,434 to		
			2013	1,022,625	1,293,542)	1454.4	(1021.7 to 1844.4)
	Hemorrhagic				(522,123 to		
	Stroke	Male	1990	689,528	883,739)	3657.2	(2738.3 to 4728.7)
					(457,313 to		
			2013	650,507	908,347)	1917.4	(1361.5 to 2710.9)
					(358,655 to		
		Female	1990	507,657	675,885)	2178.4	(1539.7 to 2894.5)
					(310,971 to		
			2013	435,037	644,826)	998.5	(706.3 to 1499.5)
					(976,733 to		
		Both	1990	1,197,186	1,433,140)	2857.1	(2317.3 to 3437.8)
					(832,966 to		
			2013	1,085,545	1,398,783)	1408.6	(1081.2 to 1850.6)
YLDs	Ischemic Stroke	Male	1990	4,978	(3,341 to 6,952)	30.1	(20.2 to 42.0)
			2013	7,949	(5,294 to 11,628)	27.8	(18.2 to 40.6)
		Female	1990	5,458	(3,712 to 7,535)	23.1	(15.7 to 32.0)
			2013	9,513	(6,286 to 13,747)	22.3	(14.8 to 31.9)
		Both	1990	10,436	(7,147 to 14,243)	26	(17.8 to 35.4)
			2013	17,462	(12,040 to 24,329)	24.6	(16.9 to 34.3)
	Hemorrhagic						
	Stroke	Male	1990	5,881	(3,975 to 8,213)	33.3	(22.3 to 46.4)
			2013	9,429	(6,238 to 13,236)	29.8	(19.4 to 41.8)
		Female	1990	5,444	(3,630 to 7,299)	22.8	(15.1 to 30.5)
			2013	9,118	(5,983 to 12,927)	21	(13.8 to 29.8)
		Both	1990	11,325	(7,794 to 15,095)	27.1	(18.8 to 36.1)
		Both	2013	18,547	(12,690 to 25,319)	24.7	(17.0 to 34.0)

Yemen	Deaths	Ischemic Stroke	Male	1990	1,288	(681 to 2,080)	104	(55.2 to 161.5)
				2013	2,988	(1,579 to 4,553)	100.2	(54.6 to 146.5)
			Female	1990	1,702	(894 to 2,736)	99.8	(56.4 to 153.4)
				2013	3,600	(1,811 to 5,571)	108.6	(59.6 to 159.6)
			Both	1990	2,990	(1,812 to 4,470)	101.5	(64.8 to 146.8)
				2013	6,589	(3,909 to 9,750)	104.6	(65.8 to 143.8)
		Hemorrhagic						
De	Deaths	Stroke	Male	1990	1,984	(1,264 to 3,050)	102.3	(59.3 to 167.0)
				2013	3,807	(2,256 to 6,297)	92.7	(55.2 to 148.0)
			Female	1990	2,771	(1,675 to 4,462)	117.3	(74.1 to 195.4)
				2013	4,910	(2,780 to 8,281)	111.5	(67.1 to 182.4)
			Both	1990	4,755	(3,041 to 7,162)	111	(71.9 to 172.8)
				2013	8,718	(5,325 to 13,697)	102.6	(65.1 to 157.4)
					Counts, All	Ages	Rate	Per 100,000
							A	ata a da udina d
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Age- Mean	95% UI
Country	measure DALYs	Stroke Type Ischemic Stroke	sex Male	year 1990	Mean 30,827	95% UI (16,482 to 52,929)		
Country		• • • • • • • • • • • • • • • • • • • •					Mean	95% UI
Country		• • • • • • • • • • • • • • • • • • • •		1990	30,827	(16,482 to 52,929)	Mean 1632.7	95% UI (849.2 to 2667.7)
Country		• • • • • • • • • • • • • • • • • • • •	Male	1990 2013	30,827 59,530	(16,482 to 52,929) (30,172 to 98,287)	Mean 1632.7 1466.6	95% UI (849.2 to 2667.7) (778.7 to 2261.7)
Country		• • • • • • • • • • • • • • • • • • • •	Male	1990 2013	30,827 59,530	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302)	Mean 1632.7 1466.6	95% UI (849.2 to 2667.7) (778.7 to 2261.7)
Country		• • • • • • • • • • • • • • • • • • • •	Male Female	1990 2013 1990 2013	30,827 59,530 37,258 70,837	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to	Mean 1632.7 1466.6 1561.9	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5)
Country		• • • • • • • • • • • • • • • • • • • •	Male	1990 2013 1990	30,827 59,530 37,258	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719)	Mean 1632.7 1466.6 1561.9	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0)
Country		• • • • • • • • • • • • • • • • • • • •	Male Female	1990 2013 1990 2013	30,827 59,530 37,258 70,837 68,085	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719) (69,703 to	Mean 1632.7 1466.6 1561.9 1621.7 1592.3	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5) (956.3 to 2418.0)
Country		Ischemic Stroke	Male Female	1990 2013 1990 2013	30,827 59,530 37,258 70,837	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719) (69,703 to 209,027)	Mean 1632.7 1466.6 1561.9	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5)
Country		Ischemic Stroke Hemorrhagic	Male Female Both	1990 2013 1990 2013 1990 2013	30,827 59,530 37,258 70,837 68,085 130,367	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719) (69,703 to 209,027) (51,213 to	Mean 1632.7 1466.6 1561.9 1621.7 1592.3	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5) (956.3 to 2418.0) (897.5 to 2336.2)
Country		Ischemic Stroke	Male Female	1990 2013 1990 2013	30,827 59,530 37,258 70,837 68,085	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719) (69,703 to 209,027) (51,213 to 125,671)	Mean 1632.7 1466.6 1561.9 1621.7 1592.3	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5) (956.3 to 2418.0)
Country		Ischemic Stroke Hemorrhagic	Male Female Both	1990 2013 1990 2013 1990 2013	30,827 59,530 37,258 70,837 68,085 130,367	(16,482 to 52,929) (30,172 to 98,287) (16,991 to 67,733) (33,809 to 118,302) (38,085 to 111,719) (69,703 to 209,027) (51,213 to	Mean 1632.7 1466.6 1561.9 1621.7 1592.3	95% UI (849.2 to 2667.7) (778.7 to 2261.7) (820.0 to 2528.0) (809.5 to 2534.5) (956.3 to 2418.0) (897.5 to 2336.2)

-					1			
						157,811)		
						(82,020 to		
				2013	145,951	247,841)	2328.5	(1318.0 to 3932.0)
						(115,049 to		
			Both	1990	183,090	265,819)	2490.4	(1534.3 to 3907.0)
						(163,187 to		
				2013	266,753	425,141)	2130.5	(1283.9 to 3407.2)
	YLDs	Ischemic Stroke	Male	1990	199	(132 to 280)	10	(6.7 to 14.2)
				2013	501	(318 to 706)	9.6	(6.0 to 13.5)
			Female	1990	179	(114 to 257)	7.3	(4.6 to 10.5)
				2013	408	(260 to 582)	7.4	(4.8 to 10.5)
			Both	1990	379	(251 to 526)	8.5	(5.7 to 11.8)
				2013	909	(612 to 1,246)	8.4	(5.7 to 11.6)
		Hemorrhagic						
		Stroke	Male	1990	31	(18 to 47)	1.3	(0.8 to 1.9)
				2013	77	(43 to 123)	1.3	(0.7 to 2.0)
			Female	1990	37	(21 to 59)	1.3	(0.8 to 2.2)
				2013	79	(44 to 128)	1.2	(0.7 to 2.1)
			Both	1990	68	(43 to 100)	1.3	(0.8 to 1.9)
			Both	2013	156	(98 to 232)	1.3	(0.8 to 1.9)
Zambia	Deaths	Ischemic Stroke	Male	1990	214	(102 to 321)	27.4	(11.2 to 43.4)
				2013	1,026	(584 to 1,471)	67.6	(38.2 to 96.1)
			Female	1990	930	(601 to 1,212)	93	(60.1 to 123.8)
				2013	2,183	(1,329 to 2,894)	119.7	(73.6 to 157.8)
			Both	1990	1,144	(780 to 1,512)	64.3	(43.0 to 86.4)
				2013	3,210	(2,182 to 3,972)	97	(64.9 to 121.6)
		Hemorrhagic			-, -	(, = ===,===	-	(1
	Deaths	Stroke	Male	1990	316	(207 to 425)	26.1	(12.6 to 40.8)
				2013	1,189	(812 to 1,645)	62.8	(39.6 to 91.5)
			Female	1990	1,069	(779 to 1,417)	85.1	(59.4 to 117.0)

					Counts, All	Ages	Rate Age-	Per 100,000 standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	2,347	(1,754 to 3,318)	110.9	(81.3 to 159.5)
			Both	1990	1,386	(1,049 to 1,741)	58.8	(41.1 to 77.2)
				2013	3,536	(2,816 to 4,668)	89.3	(68.3 to 120.5)
	DALYs	Ischemic Stroke	Male	1990	4,316	(2,473 to 5,861)	338.9	(172.1 to 500.3)
				2013	17,762	(10,338 to 25,224)	944.2	(538.4 to 1346.5)
			Female	1990	15,209	(9,633 to 19,840)	1225.9	(788.0 to 1591.2)
				2013	33,378	(19,722 to 44,404)	1602.4	(968.4 to 2118.9)
			Both	1990	19,526	(13,341 to 24,677)	830.8	(566.2 to 1090.3)
				2013	51,140	(33,402 to 63,378)	1309.3	(871.1 to 1613.8)
		Hemorrhagic Stroke	Male	1990	10,446	(7,725 to 13,846)	528.6	(356.7 to 712.0)
		Stroke	Widic	2013	29,310	(20,922 to 39,872)	1156.4	(785.5 to 1604.5)
			Female	1990	25,199	(19,080 to 33,393)	1529	(1123.5 to 2029.5)
			remare	2013	50,653	(37,555 to 70,907)	1970.5	(1470.0 to 2760.7)
			Both	1990	35,646	(28,451 to 44,102)	1073.4	(820.6 to 1351.0)
				2013	79,964	(64,668 to 103,723)	1593.1	(1274.3 to 2117.1)
	YLDs	Ischemic Stroke	Male	1990	160	(97 to 236)	9.9	(5.9 to 14.9)
				2013	285	(186 to 431)	9.9	(6.4 to 15.0)
			Female	1990	128	(82 to 190)	7.3	(4.6 to 10.7)
				2013	216	(134 to 331)	7.1	(4.3 to 11.2)
			Both	1990	289	(188 to 412)	8.5	(5.6 to 12.3)
				2013	502	(330 to 709)	8.4	(5.6 to 11.9)
		Hemorrhagic						
		Stroke	Male	1990	62	(39 to 95)	3.1	(2.0 to 4.8)
				2013	108	(68 to 166)	2.9	(1.8 to 4.3)

			Female	1990	37	(23 to 56)	1.8	(1.1 to 2.7)
				2013	62	(36 to 97)	1.6	(1.0 to 2.6)
			Both	1990	100	(66 to 145)	2.4	(1.6 to 3.6)
			Both	2013	171	(112 to 246)	2.3	(1.5 to 3.2)
Zimbabwe	Deaths	Ischemic Stroke	Male	1990	568	(437 to 768)	43	(33.1 to 59.2)
				2013	1,231	(790 to 1,713)	53.7	(34.2 to 74.8)
			Female	1990	1,213	(936 to 1,476)	81.6	(63.3 to 99.8)
				2013	2,126	(1,384 to 3,217)	72.3	(47.6 to 108.5)
			Both	1990	1,782	(1,501 to 2,145)	64.4	(54.4 to 77.8)
				2013	3,357	(2,444 to 4,475)	64.7	(47.2 to 85.7)
		Hemorrhagic						
	Deaths	Stroke	Male	1990	653	(513 to 887)	35.9	(27.6 to 50.2)
				2013	1,505	(919 to 2,069)	48.6	(27.7 to 67.9)
			Female	1990	1,187	(924 to 1,432)	62.4	(47.5 to 74.2)
				2013	2,153	(1,467 to 3,139)	56.2	(37.2 to 87.0)
			Both	1990	1,840	(1,574 to 2,186)	50.2	(41.9 to 59.3)
					Counts, All	Ages	Rate	Per 100,000
							Age-	standardized
Country	measure	Stroke Type	sex	year	Mean	95% UI	Mean	95% UI
				2013	3,658	(2,588 to 4,731)	53	(37.7 to 70.7)
	DALYs	Ischemic Stroke	Male	1990	11,169	(8,683 to 15,185)	658.9	(510.8 to 893.3)
				2013	22,406	(14,542 to 31,355)	822.6	(526.3 to 1141.3)
			Female	1990	20,091	(14,962 to 24,937)	1133.4	(864.4 to 1386.4)
				2013	31,298	(19,731 to 49,730)	951	(609.8 to 1468.5)
			Both	1990	31,260	(25,987 to 37,757)	918	(769.7 to 1097.1)
				2013	53,704	(39,268 to 72,628)	899.8	(651.7 to 1210.0)
		Hemorrhagic						
		Stroke	Male	1990	20,897	(16,287 to 28,296)	805.1	(626.9 to 1092.7)
				2013	52,446	(34,197 to 71,022)	1154.6	(717.0 to 1599.2)

		Female	1990	32,053	(25,089 to 39,804)	1315.5	(1005.3 to 1586.7)
			2013	72,051	(44,845 to 98,819)	1306.6	(885.0 to 1921.9)
		Both	1990	52,950	(45,282 to 64,807)	1074.5	(913.8 to 1272.8)
					(83,329 to		
			2013	124,497	158,841)	1237.8	(896.4 to 1623.8)
YLDs	Ischemic Stroke	Male	1990	362	(240 to 502)	17.5	(11.5 to 24.4)
			2013	513	(328 to 731)	17.4	(11.0 to 24.8)
		Female	1990	391	(254 to 566)	17.4	(11.4 to 25.0)
			2013	638	(412 to 914)	18	(11.6 to 25.9)
		Both	1990	754	(516 to 1,038)	17.4	(12.1 to 24.4)
			2013	1,151	(774 to 1,591)	17.8	(11.9 to 24.6)
	Hemorrhagic						
	Stroke	Male	1990	116	(71 to 170)	4.7	(2.8 to 6.9)
			2013	164	(104 to 245)	4.7	(2.9 to 7.1)
		Female	1990	114	(72 to 165)	4.4	(2.8 to 6.3)
			2013	170	(106 to 248)	4.3	(2.7 to 6.3)
		Both	1990	231	(151 to 321)	4.5	(2.9 to 6.3)
			2013	335	(224 to 467)	4.5	(2.9 to 6.4)