Meredith C. Foster¹, Josef Coresh²⁻⁴, Joseph V. Bonventre⁵, Venkata S. Sabbisetti⁵, Sushrut S. Waikar⁵, Theodore E. Mifflin⁶, Robert G. Nelson⁷, Morgan Grams²⁻⁴, Harold I. Feldman⁸, Vasan S. Ramachandran⁹, Paul L. Kimmel¹⁰, Chi-yuan Hsu¹¹, and Kathleen D. Liu¹¹ for the CKD Biomarkers Consortium

- Division of Nephrology, Department of Medicine, Tufts Medical Center, Boston, MA
- 2. Departments of Epidemiology and Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
- 3. Department of Medicine, Johns Hopkins Hospital, Baltimore, MD
- 4. The Welch Center for Prevention, Epidemiology and Clinical Research, Johns Hopkins Medical Institutions, Baltimore, MD
- 5. Divison of Nephrology, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA
- 6. Department of Biostatistics and Epidemiology, Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, Philadelphia, PA
- 7. National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Phoenix, AZ
- 8. Department of Biostatistics and Epidemiology, Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, Philadelphia, PA
- 9. Section of Preventive Medicine and Epidemiology, Department of Medicine, Boston University, Boston, MA
- 10. National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD
- 11. Division of Nephrology, Department of Medicine, University of California, San Francisco, San Francisco, CA