

SUPPORTING INFORMATION

Microbial immigration across the Mediterranean via airborne dust

Riccardo Rosselli¹, Maura Fiamma², Massimo Deligios², Gabriella Pintus³, Grazia Pellizzaro³

Annalisa Canu³, Pierpaolo Duce³, Andrea Squartini^{5*}, Rosella Muresu⁴, Pietro Cappuccinelli²

¹Department of Biology, University of Padova, Via Ugo Bassi 58/b, 35131 Padova, Italy

²Department of Biomedical Sciences-University of Sassari, Italy, ³Institute of Biometeorology-

National Research Council (IBIMET-CNR), Italy, ⁴Institute of Animal Production Systems in

Mediterranean Environments-National Research Council (ISPAAM-CNR),Italy. ⁵Department of

Agronomy Animals, Food, Natural Resources and Environment, DAFNAE, University of Padova,

Viale dell'Università 16, 35020 Legnaro (Padova) Italy *Corresponding author: squart@unipd.it




Figure S1 Daily PM₁₀ values recorded during the period 10-25 February 2014 at the monitoring stations located in Northern Sardinia (40.72°N, 8.55°E) and Southern Sardinia (39.23°N, 9.00°E).




Figure S2. Satellite images (MODIS), 19th and 20th February 2014 (credit to: <http://modis.gsfc.nasa.gov/data/>)