

377      **Supplemental Table 1: Antimicrobials used for Outpatient Parenteral Antibiotic Therapy (OPAT)**  
 378      patients at any time stratified by presence or absence of eosinophilia.

	All (N=824)	Eosinophilia (N=210)	No Eosinophilia (N=614)	P-value <sup>†</sup>
<i>Antibiotic<sup>‡</sup>, N (%)</i>				
Cephalosporins	377 (46)	86 (41)	291 (47)	0.11
<b>Vancomycin</b>	<b>332 (40)</b>	<b>105 (50)</b>	<b>227 (37)</b>	<b>0.001</b>
Penicillins	219 (27)	63 (30)	156 (25)	0.21
<b>Metronidazole</b>	<b>135 (16)</b>	<b>24 (11)</b>	<b>111 (18)</b>	<b>0.02</b>
Flouroquinolones	130 (16)	32 (15)	98 (16)	0.91
<b>Rifampin</b>	<b>121 (15)</b>	<b>45 (21)</b>	<b>76 (12)</b>	<b>0.002</b>
Daptomycin	74 (9)	26 (12)	48 (8)	0.05 <sup>§</sup>
Carbapenems	69 (8)	21 (10)	48 (8)	0.32
Fluconazole	47 (6)	12 (6)	35 (6)	>0.50
Linezolid	47 (6)	18 (9)	29 (5)	0.06
Aminoglycosides	30 (4)	8 (4)	22 (4)	>0.50
Tetracyclines	21 (3)	5 (2)	16 (3)	>0.50
Clindamycin	9 (1)	4 (2)	5 (<1)	0.24
Micafungin	7 (<1)	1 (<1)	6 (1)	>0.50
Voriconazole	7 (<1)	2 (1)	5 (<1)	>0.50
Aztreonam	6 (<1)	2 (1)	4 (<1)	>0.50
Amphotericin B	6 (<1)	2 (1)	4 (<1)	>0.50
Macrolides	6 (<1)	2 (1)	4 (<1)	>0.50
TMP/SMX	5 (<1)	1 (<1)	4 (<1)	>0.50
Flucytosine	2 (<1)	1 (<1)	1 (<1)	0.45

	All (N=824)	Eosinophilia (N=210)	No Eosinophilia (N=614)	P-value <sup>†</sup>
Ganciclovir	2 (<1)	0 (0)	2 (<1)	>0.50
Cefoxotin	1 (<1)	0 (0)	1 (<1)	>0.50
Ethambutol	1 (<1)	1 (<1)	0 (0)	0.25
Foscarnet	1 (<1)	0 (0)	1 (<1)	>0.50
Posaconazole	1 (<1)	0 (0)	1 (<1)	>0.50

379 *Abbreviations:* TMP/SMX: trimethoprim/sulfamethoxazole

380 <sup>†</sup>Fisher's exact test.

381 <sup>‡</sup>Proportion of patients using antibiotics anytime during OPAT follow-up

382 <sup>§</sup>For Daptomycin, P=0.0508

383

384

385

386

387

388

389

390

391

392

393

394

395

396  
397