

S6 Table – Anti-TP53 in diagnostic (D) and post-therapy (P) draws

Patient #	Kind	Interval D to P	Anti-TP53 (D)	Anti-TP53 (P)	P / D
1	Cancer	76w 4d	0.91	0.50	0.55
3	Cancer	103w 0d	0.07	0.11	1.52
4	Cancer	48w 4d	1.25	1.11	0.89
5	Cancer	81w 4d	22.45	0.92	0.04
10	Cancer	56w 0d	0.15	0.13	0.87
11	Cancer	57w 2d	1.03	0.90	0.87
12	Cancer	118w 1d	0.90	0.91	1.01
14	Cancer	109w 0d	0.55	0.52	0.93
16	Cancer	25w 5d	0.54	0.62	1.15
17	Cancer	53w 4d	0.51	0.29	0.57
20	Cancer	60w 1d	0.70	0.15	0.22
21	Cancer	51w 1d	3.81	2.78	0.73
25	Cancer	48w 1d	0.53	0.53	1.00
27	Cancer	61w 3d	0.72	0.85	1.17
29	Cancer	48w 3d	0.89	0.47	0.52
30	Cancer	80w 3d	1.18	0.78	0.66
31	Cancer	52w 1d	0.76	0.58	0.76
32	Cancer	70w 1d	0.68	0.48	0.71
34	Cancer	52w 0d	0.73	0.69	0.94
35	Cancer	53w 5d	0.39	0.29	0.74
36	Cancer	54w 6d	0.79	0.80	1.01
44	Control	107w 3d	0.27	0.03	0.11
52	Control	114w 4d	0.17	0.20	1.19
53	Control	83w 4d	0.57	0.55	0.97
57	Control	59w 2d	0.30	0.22	0.76
77	Control	53w 6d	0.38	0.44	1.18
90	Control	118w 2d	0.22	0.16	0.73
99	Control	118w 4d	0.51	0.53	1.04
104	Control	107w 2d	0.47	0.56	1.20
124	Control	172w 6d	0.54	0.16	0.30
131	Control	16w 6d	0.65	0.56	0.86

Comparison of two blood draws from the same cancer patient or healthy control. For cancers, D is the diagnostic draw and P is the post-therapeutic draw. For the controls, D and P are two blood draws with intervals similar to those for the 21 cancer patients.