

Variable	Responses					Mean (SD)
	Male	Female				
<b>Gender</b>	1073 (72%)	421 (28%)				
	<b>18 – 29</b>	<b>30 – 39</b>	<b>40 – 49</b>	<b>50 – 59</b>	<b>60+</b>	
<b>Age</b>	416 (28%)	360 (24%)	367 (25%)	249 (17%)	104 (7%)	
	<b>First Responder</b>	<b>EMT</b>	<b>AEMT</b>	<b>Paramedic</b>	<b>Physician</b>	
<b>Training</b>	41 (3%)	667 (45%)	64 (4%)	705 (47%)	16 (1%)	
	<b>Volunteer</b>	<b>Paid</b>	<b>Both</b>			
<b>Paid/Volunteer</b>	386 (26%)	810 (56%)	272 (18%)			
	<b>1 – 5</b>	<b>6 – 10</b>	<b>10 – 19</b>	<b>20 – 29</b>	<b>30+</b>	
<b>Years of Experience</b>	366 (25%)	256 (17%)	447 (30%)	276 (18%)	155 (10%)	
	<b>Once</b>	<b>Multiple</b>	<b>Never</b>			
<b>BBP Training</b>	129 (9%)	1336 (89%)	26 (2%)			
	<b>Once</b>	<b>Multiple</b>	<b>Never</b>			
<b>BSI Training</b>	106 (7%)	1370 (92%)	14 (1%)			
	<b>Yes</b>	<b>No</b>				
<b>Sanitizer in ambulance</b>	1387 (94%)	95 (6%)				
	<b>Yes</b>	<b>No</b>				
<b>Sanitizer in station</b>	1365 (92%)	126 (8%)				
	<b>Yes</b>	<b>No</b>				
<b>Brings own Sanitizer</b>	380 (25%)	1113 (75%)				
	<b>Never</b>	<b>Rarely</b>	<b>Sometimes</b>	<b>Most of the time</b>	<b>Every time</b>	
<b>Cleans before patient</b>	175 (12%)	394 (26%)	343 (23%)	394 (26%)	190 (13%)	3.0 (1.2)
<b>Cleans after skin contact</b>	21 (1%)	66 (4%)	165 (11%)	430 (29%)	813 (54%)	4.3 (0.9)
<b>Cleans when contact finishes</b>	2 (<1%)	6 (<1%)	76 (5%)	412 (28%)	997 (67%)	4.6 (0.6)
<b>Uses gloves</b>	1 (<1%)	16 (1%)	167 (11%)	522 (35%)	779 (52%)	4.4 (0.7)
<b>Uses gloves with equipment</b>	40 (3%)	252 (17%)	579 (39%)	454 (31%)	159 (11%)	3.3 (1.0)
<b>Cleans after using equipment</b>	28 (2%)	199 (13%)	435 (24%)	492 (33%)	332 (22%)	3.6 (1.0)
<b>Cleans after driving</b>	134 (9%)	341 (23%)	397 (27%)	329 (22%)	264 (18%)	3.2 (1.2)

<b>Cleans after invasive procedures</b>	228 (16%)	273 (19%)	227 (16%)	218 (16%)	456 (33%)	3.3 (1.5)
<b>Cleans stethoscope</b>	99 (7%)	223 (16%)	490 (34%)	426 (30%)	186 (13%)	3.3 (1.1)
	<b>Soap</b>	<b>Sanitizer</b>	<b>Do not clean after</b>			
<b>Cleans after hands visibly contaminated</b>	1258 (84%)	239 (16%)				
	<b>Soap</b>	<b>Sanitizer</b>	<b>No preference</b>			
<b>Soap or Sanitizer preference</b>	1059 (71%)	255 (17%)	185 (12%)			
	<b>Yes</b>	<b>No</b>	<b>I don't know</b>			
<b>Must use soap with GI illness</b>	839 (56%)	534 (36%)	124 (8%)			