SUPPLEMENTARY TABLE. Virulence of *Salmonella* serovars for mice.

Subspecies	Serogroup	Serovar	Virulence*	Ref
enterica	O:2 (A)	Paratyphi A	-	(1, 2)
	O:4 (B)	Abortusovis	-	(2)
		Bredeney	-	(3)
		Derby	-	(1)
		Heidelberg	-	(1, 3)
		Paratyphi B	-	(2)
		Paratyphi B var Java	-	(1), This study
		Reading	-	(1), This study
		Saintpaul	-	(1)
		Sandiego	-	(1)
		Schwarzengrund	-	(3)
		Stanley	-	(2)
		Stanleyville	+	This study
		Typhimurium	+	(1, 3)
	O:7 (C_1)	Braenderup	-	(2)
		Choleraesuis	+	(1, 3)
		Infantis	-	(2)
		Montevideo	-	(1-3)
		Ohio	-	(1)
		Oranienburg	-	(1)
		Thompson	-	(2)
		Virchow	-	(3)
	O:8 (C_2-C_3)	Hadar	-	(1, 3)

	Havana	-	(1)
	Muenchen	-	(2)
	Newport	-	(1, 3)
O:9 (D)	Dublin	+	(1, 3), This study
	Enteritidis	+	(1-3)
	Gallinarum	-	(1)
	Miami	-	(2)
	Naestved	+	(1)
	Panama	-	This study
	Sendai	-	(2)
	Typhi	-	(2) (1)
	9,12:l,v:-	-	(3)
O:3,10 (E)	Muenster	-	(2)
	Give var 15+ (Newbrunswick)	-	(1)
O:13 (G)	Poona	-	(2)
O:21 (L)	Minnesota	-	(1)
salamae		-	(2)
arizonae		-	(2)
diarizonae		-	(2)
houtenae		-	(2)
indica		-	(2)

* Virulence was determined by infecting BALB/c mice i.p. with 10^5 CFU (1) or $10^7 - 10^9$ CFU (this study) or intragastrically with 10^9 CFU (2), or C3H/HeN mice i.p. with ~5 x 10^3 CFU (3).

REFERENCES

1. **Roudier C, Krause M, Fierer J, Guiney DG.** 1990. Correlation between the presence of sequences homologous to the *vir* region of *Salmonella dublin* plasmid pSDL2 and the virulence of twenty-two *Salmonella* serotypes in mice. Infect Immun **58:**1180-1185.

2. **Swearingen MC, Porwollik S, Desai PT, McClelland M, Ahmer BM.** 2012. Virulence of 32 *Salmonella* strains in mice. PLoS One **7:**e36043.

3. **Suez J, Porwollik S, Dagan A, Marzel A, Schorr YI, Desai PT, Agmon V, McClelland M, Rahav G, Gal-Mor O.** 2013. Virulence gene profiling and pathogenicity characterization of non-typhoidal *Salmonella* accounted for invasive disease in humans. PLoS One **8:**e58449.