

Supplementary Data

SUPPLEMENTARY TABLE S1. pH1N1-SPECIFIC Th1
AND CYTOTOXIC T CELL SUBSETS BEFORE
AND AFTER ADMINISTRATION OF pH1N1 VACCINE

| Variable (%) ^a | Median (IQR) at baseline ^b | Median (IQR) postdose 1 ^c |
|---|--|---|
| CD4 ⁺ IL-2 ⁺ | 0.35 (-1.64, 2.07) | 0.08 (-1.22, 1.60) |
| CD4 ⁺ MIP1β ⁺ | 0.10 (-1.46, 1.73) | -0.05 (-1.73, 1.01) |
| CD4 ⁺ MIP1β ⁺ IL2 ⁺ | -0.12 (-0.44, 0.38) | -0.01 (-0.53, 0.23) |
| CD4 ⁺ PERF ⁺ | -0.35 (-1.83, 1.73) | -0.49 (-2.22, 1.31) |
| CD4 ⁺ TNF-α ⁺ | -0.19 (-1.35, 2.37) | -0.21 (-0.97, 1.04) |
| CD4 ⁺ PERF ⁺ TNF-α ⁺ | -0.08 (-0.41, 0.94) | 0.04 (-0.56, 0.46) |
| CD8 ⁺ IL-2 ⁺ | 0.34 (-1.22, 1.77) | 0.45 (-1.01, 2.08) |
| CD8 ⁺ MIP1β ⁺ | 0.67 (-1.43, 2.46) | 0.83 (-0.97, 2.49) |
| CD8 ⁺ MIP1β ⁺ IL-2 ⁺ | 0.06 (-0.91, 0.83) | 0.22 (-0.25, 1.06) |
| CD8 ⁺ PERF ⁺ | 0.95 (-2.14, 4.33) | 0.11 (-1.17, 1.76) |
| CD8 ⁺ TNF-α ⁺ | 0.72 (-0.52, 3.63) | 0.35 (-1.57, 2.48) |
| CD8 ⁺ PERF ⁺ TNF-α ⁺ | 0.38 (-0.58, 1.68) | 0.10 (-0.64, 0.84) |

^aSubsets, measured by flow cytometry, are expressed as a percentage of the parent CD4⁺ or CD8⁺ B cell population.

^bN=49.

^cN=51.