

Table S3 –Association of Log-transformed (base 2) Continuous Urine Creatinine adjusted Early Post-operative Biomarkers with Any AKI and interaction with Baseline eGFR

Biomarker	eGFR ≤ 60 ml/min/1.73 m ² (n= 424)		eGFR > 60 ml/min/1.73 m ² (n=795)		eGFR x Biomarker p-value
	unadjusted RR	adjusted RR	unadjusted RR	adjusted RR	
IL-18	1.07 [1.02, 1.12]	1.04 [0.98, 1.1]	1.14 [1.11, 1.18]	1.12 [1.08, 1.17]	0.0052
NGAL	1.03 [0.99, 1.07]	1.02 [0.98, 1.07]	1.08 [1.04, 1.11]	1.03 [1, 1.07]	0.19
KIM-1	1.17 [1.07, 1.28]	1.13 [1.03, 1.24]	1.2[1.11, 1.3]	1.16 [1.07, 1.26]	0.62
LFABP	1.02 [0.98, 1.05]	1.01 [0.97, 1.05]	1.07 [1.03, 1.1]	1.04 [1.01, 1.08]	0.1
CysC	0.9[0.84, 0.96]	0.92 [0.86, 1]	0.99 [0.92, 1.07]	0.96 [0.89, 1.03]	0.13
Albumin	1.05 [0.98, 1.12]	1.05 [0.97, 1.12]	1.08 [1.02, 1.15]	1.05 [0.98, 1.12]	0.75