

Table S6 : The change in post-operative biomarker concentration in those with and without a eGFR > 60 ml/min/1.73m² for the detection of All AKI

Log transformed delta biomarkers	eGFR ≤60 ml/min//1.73m ²		eGFR > 60 ml/min/1.73m ²		eGFR x Biomarker p-value
	unadjusted RR	adjusted RR	unadjusted RR	adjusted RR	
Urine IL-18	1.07 [1, 1.13]	1.04 [0.97, 1.12]	1.12 [1.06, 1.18]	1.12 [1.06, 1.19]	0.1
Urine NGAL	1.05 [0.99, 1.11]	1.05 [0.98, 1.13]	1.06 [1.01, 1.11]	1.04 [0.98, 1.1]	0.94
Urine KIM-1	1.03 [0.87, 1.24]	0.94 [0.78, 1.14]	1.13 [1.01, 1.27]	1.03 [0.91, 1.17]	0.61
Urine L-FABP	1.08 [0.96, 1.21]	1.08 [0.97, 1.2]	1.14 [1, 1.29]	1.14 [0.99, 1.31]	0.14
Urine Cystatin C	1.03 [0.97, 1.09]	1.02 [0.96, 1.09]	1.1 [1.04, 1.15]	1.09 [1.03, 1.15]	0.36
Urine Albumin	1.08 [0.97, 1.21]	1.1 [0.97, 1.25]	1.21 [1.11, 1.33]	1.17 [1.07, 1.29]	0.21