

Genetic Deletion of Cyclooxygenase-2 Downstream Enzyme Prostacyclin Synthase Suppresses Inflammation but Facilitates Carcinogenesis, unlike Deletion of Microsomal Prostaglandin E Synthase-1

Yuka Sasaki, Shuhei Kamiyama, Azusa Kamiyama, Konomi Matsumoto, Moe Akatsu, Yoshihito Nakatani, Hiroshi Kuwata, Yukio Ishikawa, Toshiharu Ishii, Chieko Yokoyama, and Shuntaro Hara *

*haras@pharm.showa-u.ac.jp

Supplementary Table 1. Composition of peritoneal leukocytes.

	control	PGIS KO	mPGES-1 KO	DKO
Macrophages %	45.2 ± 4.26	34.2 ± 9.79	47.6 ± 8.81	28.6 ± 7.62
Lymphocytes %	50.2 ± 3.59	60.8 ± 11.9	41.4 ± 12.6	68.6 ± 7.25
Polynuclear leukocytes %	4.57 ± 1.59	4.92 ± 2.74	11.0 ± 3.81	2.72 ± 2.40

Cytocentrifuge preparations of the exudate leukocytes were Giemsa-stained, and cell subsets were identified and counted. Results are mean±SEM (n=3)