

Supporting Information

Inactivation of *Fam20B* in the dental epithelium of mice leads to supernumerary incisors

TIAN Y, MA P, LIU C, YANG X, CRAWFORD D, YAN W, BAI D, QIN C, WANG X

Texas A&M University Baylor College of Dentistry, Dallas, TX, USA; and
West China School of Stomatology, Sichuan University, Chengdu, Sichuan, China

Table S1: Primers for PCR screen, South blotting probes, genotyping and RT-PCR

Primer name	Primer sequence (5'-3')
3' screen-F	GCTGACCGCTTCCTCGTGCTTTA
3' screen-R	CTGAACCACAGCCTATGGAGGCA
5' probe-F	AGCCAGACTTTAGTCCCAAAGTGCA
5' probe-R	CCAAGAGCAATGCCTTGCCA
3' probe-F	GCCACTTCCCACATGAGCAGAAAC
3' probe-R	TGTGTGTCTGTGGGCGTGCTGTA
Neo probe-F	CCTGAATGAACTGCAGGACGAGG
Neo probe-R	AGCTCTTCAGCAATATCACGGGTAGC
Flox-F	GAGCTTCCCCAAGAACTTAAACTCC
Flox-R	GAATACTGCTTCTACCTCTGCAACCATC
Rec-R	GAGCCTGAGGCATAACACCAACC
Cre-F	CCCGCAGAACCTGAAGATG
Cre-R	GACCCGGCAAAACAGGTAG
RT-F	GTAGAAGGGGAGCCATACGC
RT-R	CAGCATCGCAGTAGCTCTCA