

**Diel vertical dynamics of gelatinous zooplankton (Cnidaria, Ctenophora and Thaliacea) in a
subtropical stratified ecosystem (South Brazilian Bight)**

Miodeli Nogueira Júnior*, Frederico P Brandini & Juan Carlos Ugaz Codina

*miodeli@gmail.com

S4 Table. Thaliaceans species list and summary of the catches. DS = developmental stage; B = blastozooids (aggregates); O = oozoids (solitary), P = phorozoids; G = gonozooids; N = nurses; A = all stages combined. Other legends as in S1 Table.

Taxa	DS	Average density (±SD)		FC	RA	Weighted mean depth (±SD)		t
		Day	Night			Day	Night	
THALIACEA								
Salpida								
<i>Cyclosalpa</i> spp. (<i>C. bakeri</i> and <i>C. polae</i> ?)	B	0.26 (±0.33)	0.09 (±0.1)	8.33	0.33	77.8 (±10.1)	20	
	O	0	0.04 (±0.09)	1.39	0.04	-	20	
(?) <i>Ritteriella amboinensis</i> (Apstein, 1904)	O	0	0.04 (±0.09)	1.39	0.04	-	55	
<i>Salpa fusiformis</i> Cuvier, 1804	B	3.44 (±3.6)	2.13 (±1.8)	41.67	5.38	56.8 (±16.8)	27.5 (±9.8)	2.93*
	O	1.48 (±1.8)	0.74 (±0.41)	34.72	2.13	51.7 (±21.7)	44.3 (±13.3)	0.58
<i>Thalia democratica</i> (Forskål, 1775)	B	15.15 (±17.9)	7 (±2.9)	66.67	21.23	50.9 (±10.4)	22.2 (±3.8)	5.17**
	O	1.48 (±1.6)	0.17 (±0.14)	26.39	1.58	54.06 (±17.7)	41.7 (±37.5)	0.59
Doliolida								
<i>Doliolletta gegenbauri</i> (Uljanin, 1884)	P	7.53 (±9.7)	2.6 (±1.28)	61.11	9.92	44.7 (±17.1)	36.1 (±0.9)	1
	G	1.22 (±1)	0.48 (±0.26)	33.33	1.67	44.5 (±18.7)	22.7 (±5.4)	2.2
	N	2.52 (1.18)	2.48 (±1.3)	64.28	4.9	43.5 (±5.9)	30.7 (±2.3)	4**
<i>Doliolum nationalis</i> Borgert, 1893	P	10.14 (±10.8)	3.22 (±3.05)	56.94	12.8	41 (±12.4)	32 (±5.2)	1.34
	G	23.46 (±24.1)	12 (±9.8)	61.11	33.99	29.9 (±3.6)	21.9 (±0.6)	4.38**
	N	3.39 (±2.06)	1.87 (±1.13)	55.55	5.04	45.1 (±9.9)	23.4 (±4.3)	4.03**
Destroyed, unidentified				5.55	0.54			
Pirossomatida								
(?) <i>Pyrossomella verticillata</i> (Neumann, 1909)		0.09 (±0.1)	0.09 (±0.1)	5.55	0.17	67.5 (±17.7)	20	