

Supplementary Table 1. Incidence of condyloma^a by HPV type detected in the lesion among men^b with the same HPV type detected at the genitals either as a prevalent or incident infection

HPV Type ^c	Prevalent Genital HPV Infection					Incident Genital HPV Infection				
	Incidence Rate ^d (95% CI)	Cumulative Incidence (%)			Median time to EGL ^e	Incidence Rate ^d (95% CI)	Cumulative Incidence (%)			Median time to EGL ^e
		6-Month (95% CI)	12-Month (95% CI)	24-Month (95% CI)			6-Month (95% CI)	12-Month (95% CI)	24-Month (95% CI)	
Any	1.3 (1.0-1.6)	0.9 (0.4- 1.3)	2.4 (1.6-3.1)	3.4 (2.5-4.3)	7.6	0.9 (0.6-1.2)	0.5 (0.2-0.8)	1.2 (0.7-1.7)	2.1 (1.4-2.8)	7.7
Vaccine ^f	5.3 (3.9-7.0)	3.9 (2.0-5.8)	9.7 (6.7-12.8)	13.5 (9.8-17.2)	7.1	4.0 (2.7-5.6)	2.0 (0.7-3.3)	5.1 (2.9-7.2)	8.9 (5.7-12.1)	8.2
High Risk	0.3 (0.1-0.6)	0.1 (0.0-0.3)	0.6 (0.1-1.1)	0.8 (0.2-1.3)	8.1	0.3 (0.1-0.5)	0.2 (0.0-0.5)	0.4 (0.0-0.8)	0.8 (0.1-1.4)	7.4
16	0.7 (0.1-2.0)	0.0 (0.0-0.0)	1.2 (0.3-4.8)	2.0 (0.6-6.1)	6.4	0.3 (0.0-1.5)	0.0 (0.0-0.0)	0.7 (0.1-4.5)	0.7 (0.1-4.5)	8.3
18	0.6 (0.0-3.3)	1.6 (0.2-10.9)	1.6 (0.2-10.9)	1.6 (0.2-10.9)	5.7	0.0 (0.0-2.5)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE
31	0.0 (0.0-2.9)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE	0.9 (0.0-4.8)	1.6 (0.2-10.9)	1.6 (0.2-10.9)	1.6 (0.2-10.9)	5.7
39	0.0 (0.0-1.6)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE	0.4 (0.0-2.2)	0.7 (0.1-5.2)	0.7 (0.1-5.2)	0.7 (0.1-5.1)	1.0
45	0.0 (0.0-3.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE	0.6 (0.0-3.4)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	3.2 (0.5-20.8)	23.9
51	0.2 (0.0-1.2)	0.0 (0.0-0.0)	0.6 (0.1-4.4)	0.6 (0.1-4.4)	8.4	0.3 (0.0-1.5)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	1.1 (0.2-7.5)	17.9
52	1.0 (0.2-3.0)	0.0 (0.0-0.0)	2.0 (0.5-7.7)	2.0 (0.5-7.7)	9.2	0.3 (0.0-1.6)	0.0 (0.0-0.0)	0.7 (0.1-4.8)	0.7 (0.1-4.8)	7.4
56	0.0 (0.0-2.5)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE	0.8 (0.0-4.5)	1.7 (0.2-11.2)	1.7 (0.2-11.2)	1.7 (0.2-11.2)	0.4
Low Risk	2.6 (1.9-3.4)	1.9 (1.0-2.9)	4.7 (3.2-6.2)	7.0 (5.0-9.0)	7.5	1.6 (1.1-2.3)	0.8 (0.3-1.4)	2.2 (1.3-3.2)	3.8 (2.4-5.2)	8.0
6	15.1 (10.7-26.1)	9.3 (5.3-16.2)	22.5 (15.8-31.5)	32.4 (24.2-42.5)	7.5	9.8 (6.1-15.1)	2.5 (0.8-7.6)	10.3 (5.8-17.9)	20.1 (12.9-30.6)	10.8
11	12.7 (5.1-26.1)	12.0 (4.0-32.7)	29.6 (15.3-52.4)	29.6 (15.3-52.4)	6.7	13.5 (6.5-24.8)	12.5 (5.8-25.7)	17.8 (9.2-32.7)	26.4 (14.4-45.4)	3.7
40	0.0 (0.0-3.2)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE	0.8 (0.0-4.6)	0.0 (0.0-0.0)	1.8 (0.2-11.8)	1.8 (0.2-11.8)	6.9
53	0.5 (0.1-1.7)	0.6 (0.1-4.5)	0.6 (0.1-4.5)	1.6 (0.4-6.4)	11.1	0.0 (0.0-1.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE
54	0.4 (0.0-2.3)	0.0 (0.0-0.0)	1.2 (0.2-8.3)	1.2 (0.2-8.3)	7.8	0.0 (0.0-1.5)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	NE
66	0.6 (0.1-2.0)	0.0 (0.0-0.0)	0.0 (0.0-0.0)	2.0 (0.5-7.8)	15.0	0.2 (0.0-1.3)	0.0 (0.0-0.0)	0.6 (0.1-4.0)	0.6 (0.1-4.0)	8.9

Abbreviation: CI-confidence interval. NE-not estimable.

^aDNA detected using Linear Array. ^bNewly acquired-pathologically confirmed condyloma/suggestive of condyloma. ^cHPV types 33/35/58/59/26/68/69/71/70/73/82 did not progress to a condyloma lesion and therefore incidence rates and cumulative incidence could not be calculated. ^dIncidence rate is cases per 1000 person-months. ^eFollow-up time in months. ^fVaccine HPV types 6/11/16/18

Supplementary Table 2. HPV types detected within 161 condylomas

	n	%
Single infections		
HPV 6	90	0.56
HPV 11	33	0.20
HPV 16	1	0.01
HPV45	1	0.01
Coinfections		
HPV 6+ other HPV	16	0.10
HPV 11+ other HPV	8	0.05
HPV 6 and 11+	4	0.02
HPV 6, 11 and other HPV	4	0.02
Other HPV without 6 or 11	4	0.02