

Online Supplementary Document

Azmatullah et al. Systematic review of the global epidemiology, clinical and laboratory profile of enteric fever
 J Glob Health 2015;5:010407

Supplementary tables and definitions of variables

- Table S1: Epidemiology of enteric fever by age

Demographics/History	0 to 5 years 14 studies			5 to 10 years 6 studies			10 to 17 years 4 studies			Adults (12 years & up) 69 studies		
	N	N	%	n	N	%	n	N	%	n	N	%
Males	314	529	59	421	714	59	135	237	57	2761	4551	61
Blood/bone marrow Isolates	615	655	94	695	695	100	282	282	100	4153	4465	93
Pre-treatment antibiotics received	2	16	13							510	1240	41
Duration of illness ≤ 1 week	13	22	59							287	506	57
Duration of illness within 1-2 weeks	7	18	39							45	195	23
Duration of illness > 2 weeks	2	18	11							115	691	17
Duration of illness > 1 week	9	18	50							293	971	30
Multi-drug resistant isolates	94	383	25	121	558	22	58	237	24	913	1726	53
Fluoroquinolone resistance	10	161	6							11	1662	1
Chloramphenicol resistance	47	99	47							1119	1877	60
Ampicillin/Amoxyccillin resistance	36	71	51							1081	1461	74
Co-trimoxazole resistance	28	71	39							936	1158	81
Laboratory features												
Severe anaemia (<7.5 g/dL)	45	91	49							17	437	4
Neutrophilia(>7.5 x 10 ³ /µL)	9	9	100							4	50	8
Thrombocytopenia (<150 x 10 ³ /µL)	0	12	0	5	28	18	4	31	13	210	395	53
Severe thrombocytopenia(<50 x 10 ³ /µL)										13	194	7
Hemolysis or hemolytic anaemia										1	100	1
Other signs and symptoms												
Chills or rigors	4	33	12	4	125	3				362	769	47
Bradycardia/relative bradycardia										532	1142	47
Dehydration							1	8	13	164	288	57
Anorexia/refusal to feed	36	55	65	82	125	66				355	765	46
Weakness/dizziness or malaise							5	8	63	223	472	47
Continuous fever	8	18	44							113	274	41
Intermittent fever	5	18	28							32	191	17
Remittent fever	5	18	28							86	199	43
Stepladder fever										63	255	25
Chelosis										2	82	2
URTI	5	34	15	14	65	22	5	45	11	34	334	10
Other complications												
Myocarditis	1	51	2	6	125	5				72	862	8
Meningitis or meningismus	7	139	5	3	153	2	0	31	0	29	1645	2
Hepatitis or raised LFTs	17	47	36	19	72	26				547	1955	28
Jaundice	6	46	13	0	28	0	1	39	3	43	985	4
DIC										22	573	4
Acute acalculouscholecystitis	7	71	10							7	97	7
Edema or pleural effusion or ascites	0	12	0	6	69	9	1	31	3	9	512	2
Convulsions	10	122	8	3	125	2	0	8	0			

Focal neurological involvement												
Epistaxis/bleeding (unspecified)	16	140	11	16	166	10				96	1102	9
Any abscess	1	71	1							15	315	5
Osteomyelitis or septic arthritis	11	116	9							16	435	4
Renal involvement	2	71	3							2	30	7
Arthritis or arthralgia or myalgia	0	18	0	1	28	4	4	39	10	140	1277	11
										194	1285	15

- **Table S2: Associations of clinical features in enteric fever by age (0-5 yrs old versus 5-10 yrs old)**

Clinical Features	Odds Ratio	95% Confidence Interval
Headache	0.1	0.05 - 0.2
Diarrhea	2.5	2.0 - 3.2
Constipation	0.4	0.3 - 0.6
Nausea or vomiting	0.6	0.4 - 0.8
Hepatomegaly	1.3	1.0 - 1.6
Abdominal pain or tenderness	0.1	0.07 - 0.2
Abdominal distention or ileus	0.2	0.1 - 0.4
Leucopenia (< 5 x 10 ³ /µL)	0.2	0.1 - 0.4
Leucocytosis(> 15 x 10 ³ /µL)	4.6	3.5 - 6.2
Altered mental status	0.4	0.2 - 0.6
Chills or rigors	4.1	1.0 - 17.7
Death	4.6	2.3 - 9.2

- **Table S3: Associations of clinical features in enteric fever by age (children versus adults*)**

Clinical Features	Odds Ratio	95% Confidence Interval
Multi-drug resistant isolates	0.3	0.2- 0.3
Pre-treatment antibiotics received	0.2	0.04 - 0.9
Coated tongue	3.5	2.5- 5.1
Cough	0.7	0.6- 0.9
Nausea or vomiting	0.7	0.5- 0.8
Diarrhea	0.6	0.5 - 0.6
Constipation	0.7	0.6 - 0.9
Hepatomegaly	2.1	1.8- 2.5
Splenomegaly	0.6	0.5- 0.7
Abdominal distention or ileus	2.7	2.0- 3.7
Anemia (Hb < 12 g/dl)	5.8	3.8- 9.0
Leucopenia (< 5 x 10 ³ /µL)	1.4	1.0- 1.8
Leucocytosis (> 15 x 10 ³ /µL)	2.7	1.8- 4.0
Thrombocytopenia (<150 x 10 ³ /µL)	0.1	0.05- 0.3
Altered mental status	0.7	0.6- 0.9
GI perforation	0.4	0.2- 0.8
Pneumonia or chest signs	1.7	1.3- 2.2
Hepatitis or raised LFTs	5.9	3.9 - 8.8
Jaundice	1.9	1.0 - 3.5
Chills or rigors	0.06	0.03- 0.1

URTI	2.1	1.1- 3.9
High grade fever	0.3	0.1 - 0.5
Fluoroquinolone resistance	9.4	3.9- 22.5
Chloramphenicol resistance	0.6	0.4 - 0.9
Ampicillin/Amoxycillin resistance	0.4	0.2 - 0.6
Co-trimoxazole resistance	0.2	0.1 - 0.3
Severe anaemia (<7.5)	24.2	12.8 - 45.6
Anorexia/refusal to feed	2.2	1.6 - 3.1
Edema or pleural effusion or ascites	4.5	1.5 - 12.9
Epistaxis/bleeding (unspecified)	2.3	1.2 - 4.4
Renal involvement	0.2	0.06 - 1.0
Arthritis or arthralgia or myalgia	0.1	0.02 - 1.0

* Children - 0 - 5 yrs old versus 5 - 10 yrs old
 Adults - Author defined or aged 12 years and above

References to Table 1 , Table S1, S2 and S3: Epidemiology of enteric fever by age

Ages 0- 5 years

1. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. ArchDisChild. 1996;75(3):214-7.
2. Chiu CH, Lin TY. Typhoid fever in children. Lancet. 1999;354(9194):2001-2.
3. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. Acta Paediatr Taiwan. 2000;41(1):28-32.
4. Duggan MB, Beyer L. Enteric fever in young Yoruba children. ArchDisChild. 1975;50(1):67-71.
5. Emary K, Moore CE, Champheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 Salmonella enterica serovar Typhi with intermediate susceptibility to ciprofloxacin. Trans R Soc Trop Med Hyg. 2012 Dec;106(12):718-24.
6. Ferreccio C, Levine MM, Manterola A, Rodriguez G, Rivara I, Prenzel I, et al. Benign bacteremia caused by *Salmonella typhi* and *paratyphi* in children younger than 2 years. J Pediatr. 1984;104(6):899-901.
7. Garg RA, Krashak R. Typhoid fever before two years of age. Indian Pediatr. 1993;30(6):805-8.
8. Green SD, Cheesbrough JS. *Salmonella* bacteraemia among young children at a rural hospital in western Zaire. AnnTropPaediatr. 1993;13(1):45-53.
9. Oh HM, Masayu Z, Chew SK. Typhoid fever in hospitalized children in Singapore. J Infect. 1997;34(3):237-42.
10. Owais A, Sultana S, Zaman U, Rizvi A, Zaidi AK. Incidence of typhoid bacteremia in infants and young children in southern coastal Pakistan. Pediatr Infect Dis J. 2010 Nov;29(11):1035-9.
11. Palacios Malmaceda PG, Vela Acosta JJ, Gutierrez AW. [Typhoid fever in children under 2 years of age]. BolMedHospInfant Mex. 1981;38(3):473-83.
12. Rajajee S, Anandi TB, Subha S, Vatsala BR. Patterns of resistant *Salmonella typhi* infection in infants. JTropPediatr. 1995;41(1):52-4.

13. Sinha A, Sazawal S, Kumar R, Sood S, Reddaiah VP, Singh B, et al. Typhoid fever in children aged less than 5 years. Lancet. 1999;354(9180):734-7.
14. Soelistyowati S, Soenarto Y, Soesilo H, Widiarto, Widiatmodjo, Ismangoen. Typhoid fever in children. PaediatrIndones. 1982;22(7-8):138-46.
15. Walia M, Gaind R, Paul P, Mehta R, Aggarwal P, Kalaivani M. Age-related clinical and microbiological characteristics of enteric fever in India. TransRSocTropMed Hyg. 2006;100(10):942-8.
16. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. EpidemiolInfect. 2005;133(6):1073-9.

Ages 5 -10 years

1. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. ArchDisChild. 1996;75(3):214-7.
2. Chiu CH, Lin TY. Typhoid fever in children. Lancet. 1999;354(9194):2001-2.
3. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. Acta PaediatrTaiwan. 2000;41(1):28-32.
4. Emary K, Moore CE, Chanpheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 Salmonella enterica serovar Typhi with intermediate susceptibility to ciprofloxacin. Trans R Soc Trop Med Hyg. 2012 Dec;106(12):718-24
5. Leung DT, Bogetz J, Itoh M, Ganapathi L, Pietroni MA, Ryan ET, et al. Factors associated with encephalopathy in patients with Salmonella enterica serotype Typhi bacteremia presenting to a diarrheal hospital in Dhaka, Bangladesh. Am J Trop Med Hyg. 2012 Apr;86(4):698-702.
6. Moussa K. Enterica of childhood. J EgyptPublic Health Assoc. 1970;45(1):48-55.
7. Oh HM, Masayu Z, Chew SK. Typhoid fever in hospitalized children in Singapore. J Infect. 1997;34(3):237-42.
8. Soelistyowati S, Soenarto Y, Soesilo H, Widiarto, Widiatmodjo, Ismangoen. Typhoid fever in children. PaediatrIndones. 1982;22(7-8):138-46.
9. Walia M, Gaind R, Paul P, Mehta R, Aggarwal P, Kalaivani M. Age-related clinical and microbiological characteristics of enteric fever in India. TransRSocTropMed Hyg. 2006;100(10):942-8.

Ages 10-17yrs

1. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. ArchDisChild. 1996;75(3):214-7.
2. Chiu CH, Lin TY. Typhoid fever in children. Lancet. 1999;354(9194):2001-2.
3. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. Acta PaediatrTaiwan. 2000;41(1):28-32.
4. Emary K, Moore CE, Chanpheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 Salmonella enterica serovar Typhi with intermediate susceptibility to ciprofloxacin. Trans R Soc Trop Med Hyg. 2012 Dec;106(12):718-24
5. Kanra G, Yurdakok M, Secmeer G, Arvas A, Dereli O, Sarikayalar F, et al. Clinical findings of pediatric patients during an outbreak of typhoid fever in Ankara. TurkJ Pediatr. 1985;27(1):11-5.
6. Moussa K. Enterica of childhood. J EgyptPublic Health Assoc. 1970;45(1):48-55.

Adults

1. Abraham G, Teklu B. Typhoid fever: clinical analysis of 50 Ethiopian patients. *Ethiop Med J.* 1981;19(2):41-6.
2. Aghanwa HS, Morakinyo O. Correlates of psychiatric morbidity in typhoid fever in a Nigerian general hospital setting. *Gen Hosp Psychiatry.* 2001;23(3):158-62.
3. Ali G, Rashid S, Kamli MA, Shah PA, Allaqqaband GQ. Spectrum of neuropsychiatric complications in 791 cases of typhoid fever. *Trop Med Int Health.* 1997;2(4):314-8.
4. Abdool Gaffar MS, Seedat YK, Coovadia YM, Khan Q. The white cell count in typhoid fever. *Trop Geogr Med.* 1992;44(1-2):23-7.
5. Alsoub H, Uwaydah AK, Matar I, Zebeib M, Elhag KM. A clinical comparison of typhoid fever caused by susceptible and multidrug-resistant strains of *Salmonella typhi*. *Br J Clin Pract.* 1997;51(1):8-10.
6. Alam MN, Haq SA, Das KK, Baral PK, Mazid MN, Siddique RU, et al. Efficacy of ciprofloxacin in enteric fever: comparison of treatment duration in sensitive and multidrug-resistant *Salmonella*. *Am J Trop Med Hyg.* 1995;53(3):306-11.
7. Barrett FC, Knudsen JD, Johansen IS. Cases of typhoid fever in Copenhagen region: a retrospective study of presentation and relapse. *BMC research notes.* 2013;6:315.
8. Bhatia RS. Changing trends in the presentation and treatment of enteric fever: our experience. *The Journal of the Association of Physicians of India.* 1992;40(4):283-4.
9. Butler T, Bell WR, Levin J, Linh NN, Arnold K. Typhoid fever. Studies of blood coagulation, bacteremia, and endotoxemia. *Arch Intern Med.* 1978 Mar;138(3):407-10.
10. Butler T, Linh NN, Arnold K, Pollack M. Chloramphenicol-resistant typhoid fever in Vietnam associated with R factor. *Lancet.* 1973;2(7836):983-5.
11. Carcelen A, Chirinos J, Yi A. Furazolidone and chloramphenicol for treatment of typhoid fever. *Scand J Gastroenterol Suppl.* 1989;169:19-23.
12. Catalano G, Del Vecchio-Blanco C, Varone GL. Changing pattern of typhoid fever: clinical bacteriological and immunological aspects. *Boll Ist Sieroter Milan.* 1972;51(2):157-64.
13. Caumes E, Ehya N, Nguyen J, Bricaire F. Typhoid and paratyphoid fever: a 10-year retrospective study of 41 cases in a Parisian hospital. *J Travel Med.* 2001;8(6):293-7.
14. Chakravorty B, Jain N, Gupta B, Rajvanshi P, Sen MK, Krishna A. Chloramphenicol resistant enteric fever. *J Indian Med Assoc.* 1993;91(1):10-3.
15. Clark TW, Daneshvar C, Pareek M, Perera N, Stephenson I. Enteric fever in a UK regional infectious diseases unit: a 10 year retrospective review. *J Infect.* 2010 Feb;60(2):91-8.
16. Chuttani HK, Jain K, Misra RC. Small bowel in typhoid fever. *Gut.* 1971 Sep;12(9):709-12.
17. Daga MK, Sarin K, Sarkar R. A study of culture positive multidrug resistant enteric fever--changing pattern and emerging resistance to ciprofloxacin. *J Assoc Physicians India.* 1994;42(8):599-600.
18. Daoud AS, Zaki M, Pugh RN, al-Mutairi G, Beseiso R, Nasrallah AY. Clinical presentation of enteric fever: its changing pattern in Kuwait. *J Trop Med Hyg.* 1991;94(5):341-7.
19. Downey D, Anzimlt JC, Ellis-Pegler RB. *Salmonella typhi* infection in adults is not limited to travellers returning from the tropics. *NZ Med J.* 1986;99(804):443-6.

20. Duff P, Engelsgjerd B. Typhoid fever on an obstetrics-gynecology service. *AmJ ObstetGynecol.* 1983;145(1):113-4.
21. Dutta TK, Beerasha, Ghotekar LH. Atypical manifestations of typhoid fever. *JPostgradMed.* 2001;47(4):248-51.
22. Goh KT, Teo SH, Tay L, Monteiro EH. Epidemiology and control of an outbreak of typhoid in a psychiatric institution. *EpidemiolInfect.* 1992;108(2):221-9.
23. Gupta S, Meena HS. Changing profile of enteric fever--in summer-91. *J AssocPhysicians India.* 1992;40(11):726-9.
24. Gupta SP, Gupta MS, Bhardwaj S, Chugh TD. Current clinical patterns of typhoid fever: a prospective study. *J TropMed Hyg.* 1985;88(6):377-81.
25. Gupta V, Kaur U, Singh G, Prakash C, Sharma M, Aggarwal KC. An outbreak of typhoid fever in Chandigarh, North India. *TropGeogrMed.* 1986;38(1):51-4.
26. Haq SA, Alam MN, Hossain SM, Ahmed T, Tahir M. Value of clinical features in the diagnosis of enteric fever. *Bangladesh Med ResCouncBull.* 1997;23(2):42-6.
27. Hoa NT, Diep TS, Wain J, Parry CM, Hien TT, Smith MD, et al. Community-acquired septicaemia in southern Viet Nam: the importance of multidrug-resistant *Salmonella typhi*. *TransRSocTropMedHyg.* 1998;92(5):503-8.
28. Hosoglu S, Geyik MF, Akalin S, Ayaz C, Kokoglu OF, Loeb M. A simple validated prediction rule to diagnose typhoid fever in Turkey. *TransRSocTropMed Hyg.* 2006;100(11):1068-74.
29. Hyams KC, Oldfield EC, Scott RM, Bourgeois AL, Gardiner H, Pazzaglia G, et al. Evaluation of febrile patients in Port Sudan, Sudan: isolation of dengue virus. *AmJ TropMed Hyg.* 1986;35(4):860-5.
30. Ispahani P, Slack RC. Enteric fever and other extraintestinal salmonellosis in University Hospital, Nottingham, UK, between 1980 and 1997. *EurJ ClinMicrobiolInfectDis.* 2000;19(9):679-87.
31. Kamat SA. Evaluation of therapeutic efficacy of trimethoprim-sulphamethoxazole and chloramphenicol in enteric fever. *BrMedJ.* 1970;3(5718):320-2.
32. Kamat SA, Herzog C. Typhoid: clinical picture and response to chloramphenicol. Prospective study in Bombay (1972). *Infection.* 1977;5(2):85-91.
33. Khan M, Coovadia Y, Connolly C, Sturm AW. Risk factors predicting complications in blood culture-proven typhoid fever in adults. *ScandJ InfectDis.* 2000;32(2):201-5.
34. Khan M, Coovadia YM, Connolly C, Sturm AW. Influence of sex on clinical features, laboratory findings, and complications of typhoid fever. *AmJ TropMed Hyg.* 1999;61(1):41-6.
35. Khan MA, Hayat Z, Sadick A. Ofloxacin in the treatment of typhoid fever resistant to chloramphenicol and amoxicillin. *ClinTher.* 1994;16(5):815-8.
36. Khosla SN. Typhoid hepatitis. *PostgradMed J.* 1990;66(781):923-5.
37. Khosla SN, Anand A, Singh U, Khosla A. Haematological profile in typhoid fever. *TropDoct.* 1995;25(4):156-8.
38. Khosla SN, Kalra IJ, Sharma A. Plasma cortisol levels in enteric fever. *J AssocPhysicians India.* 1989;37(10):643-5.
39. Khosla SN, Lochan R. Renal dysfunction in enteric fever. *J AssocPhysicians India.* 1991;39(5):382-4.
40. Khosla SN, Saman A, Khosla P, Sabharwal U, Khosla A. Drug resistant typhoid fever. *TropDoct.* 1998;28(4):235-7.

41. Khosla SN, Singh R, Singh GP, Trehan VK. The spectrum of hepatic injury in enteric fever. *Am J Gastroenterol*. 1988;83(4):413-6.
42. Kuvandik C, Karaoglan I, Namiduru M, Baydar I. Predictive value of clinical and laboratory findings in the diagnosis of the enteric fever. *New Microbiol*. 2009 Jan;32(1):25-30.
43. Leung DT, Bogetz J, Itoh M, Ganapathi L, Pietroni MA, Ryan ET, et al. Factors associated with encephalopathy in patients with *Salmonella enterica* serotype Typhi bacteremia presenting to a diarrheal hospital in Dhaka, Bangladesh. *Am J Trop Med Hyg*. 2012 Apr;86(4):698-702.
44. Liberti A, Loiacono L. Ciprofloxacin versus chloramphenicol in the treatment of salmonella infection. *Int J Antimicrob Agents*. 2000;16(3):347-8.
45. Lin FY, Becke JM, Groves C, Lim BP, Israel E, Becker EF, et al. Restaurant-associated outbreak of typhoid fever in Maryland: identification of carrier facilitated by measurement of serum Vi antibodies. *J Clin Microbiol*. 1988;26(6):1194-7.
46. Mathura KC, Chaudhary D, Simkhada R, Pradhan M, Shrestha P, Gurubacharya DL. Study of clinical profile and antibiotic sensitivity pattern in culture positive typhoid fever cases. *Kathmandu Univ Med J (KUMJ)*. 2005;3(4):376-9.
47. Mathura KC, Gurubacharya DL, Shrestha A, Pant S, Basnet P, Karki DB. Clinical profile of typhoid patients. *Kathmandu Univ Med J (KUMJ)*. 2003;1(2):135-7.
48. Misra RC, Kulpati DD, Bala S, Prakash K, Gupta SK, Chuttani HK. Niridazole-in treatment of typhoid fever. *Chemotherapy*. 1971;16(5):326-31.
49. Morgenstern R, Hayes PC. The liver in typhoid fever: always affected, not just a complication. *Am J Gastroenterol*. 1991;86(9):1235-9.
50. Mukhtar ED, Mekki MO. Trimethoprim-sulphamethoxazole in the treatment of enteric fever in the Sudan. *Trans R Soc Trop Med Hyg*. 1981;75(6):771-3.
51. Nand N, Sharma M, Bhutani M, Singh GP, Sharma VK. Cardiac status in typhoid fever. *Angiology*. 1996;47(11):1095-100.
52. Nazar HS, Rabbani A, Riaz A, Anwar J. Presentation of typhoid fever patients in Hazara Division and response to different treatment regimens. *J Ayub Med Coll Abbottabad*. 2005;17(1):67-9.
53. Ndububa DA, Erhabor GE, Akinola DO. Typhoid and paratyphoid fever: a retrospective study. *Trop Gastroenterol*. 1992;13(2):56-63.
54. Neopane A, Singh SB, Bhatta R, Dhital B, Karki DB. Changing spectrum of antibiotic sensitivity in enteric fever. *Kathmandu Univ Med J (KUMJ)*. 2008 Jan-Mar;6(1):12-5.
55. Okome-Nkoumou M, Ayo NE, Bekale J, Kombila M. [Typhoid and paratyphoid fever in adults in the Internal Medicine Department at Libreville (Gabon)]. *Sante*. 2000;10(3):205-9.
56. Olle-Goig JE, Ruiz L. Typhoid fever in rural Haiti. *Bull Pan Am Health Organ*. 1993;27(4):382-8.
57. Panhotra BR, Saxena AK, Al-Ghamdi AM. Typhoid fever due to multiresistant *Salmonella enterica* serovar typhi having reduced susceptibility to ciprofloxacin and nalidixic acid resistance. *Saudi Med J*. 2004;25(10):1509-11.
58. Park SC, Lee CH, Kim SY, Park CH, Kim TW, Seok SE, et al. Clinical trial of cefotaxime in patients with typhoid fever. *Clin Ther*. 1985;7(4):448-51.
59. Pohan HT. Clinical and laboratory manifestations of typhoid fever at Persahabatan Hospital, Jakarta. *Acta Med Indones*. 2004;36(2):78-83.

60. Rogerson SJ, Spooner VJ, Smith TA, Richens J. Hydrocortisone in chloramphenicol-treated severe typhoid fever in Papua New Guinea. *TransRSocTropMed Hyg.* 1991;85(1):113-6.
61. Sanford JP, Linh NN, Kutscher E, Arnold K, Gould K. Oxolinic acid in the treatment of typhoid fever due to chloramphenicol-resistant strains of *Salmonella typhi*. *AntimicrobAgents Chemother.* 1976;9(3):387-92.
62. Sardesai HV, Melinkere RD, Diwate AB. Trimethoprim-sulphamethoxazole in typhoid fever. *TransRSocTropMed Hyg.* 1971;65(2):189-94.
63. Shin BM, Paik IK, Cho HI. Bone marrow pathology of culture proven typhoid fever. *J Korean Med Sci.* 1994;9(1):57-63.
64. Shlim DR, Schwartz E, Eaton M. Clinical Importance of *Salmonella Paratyphi A* Infection to Enteric Fever in Nepal. *JTravelMed.* 1995;2(3):165-8.
65. Smith SI, Odunukwe NN, Niemogha MT, Ahmed AO, Efienemokwu CA, Otuonye MN, et al. Diagnostic methods for typhoid fever in Nigeria. *BrJBiomedSci.* 2004;61(4):179-81.
66. Spencer DC, Pienaar NL, Atkinson PM. Disturbances of blood coagulation associated with *Salmonella typhi* infections. *J Infect.* 1988;16(2):153-61.
67. Su CP, Chen YC, Chang SC. Changing characteristics of typhoid fever in Taiwan. *J MicrobiolImmunolInfect.* 2004;37(2):109-14.
68. Sulaiman K, Sarwari AR. Culture-confirmed typhoid fever and pregnancy. *IntJInfectDis.* 2007;11(4):337-41.
69. Torin DE, Betts SL, McClenahan JB, Phillips GM, Shikamura MT. A typhoid fever outbreak on a university campus. *ArchInternMed.* 1969;124(5):606-10.
70. Trevett AJ, Nwokolo N, Lightfoot D, Naraqi S, Kevau IH, Temu PI, et al. Ataxia in patients infected with *Salmonella typhi* phage type D2: clinical, biochemical and immunohistochemical studies. *TransRSocTropMed Hyg.* 1994;88(5):565-8.
71. Unal S, Hayran M, Tuncer S, Gur D, Uzun O, Akova M, et al. Treatment of enteric fever with pefloxacin for 7 days versus 5 days: a randomized clinical trial. *AntimicrobAgents Chemother.* 1996;40(12):2898-900.
72. Uwaydah AK, al SH, Matar I. Randomized prospective study comparing two dosage regimens of ciprofloxacin for the treatment of typhoid fever. *J AntimicrobChemother.* 1992;30(5):707-11.
73. Valenciano M, Baron S, Fisch A, Grimont F, Desenclos JC. Investigation of concurrent outbreaks of gastroenteritis and typhoid fever following a party on a floating restaurant, France, March 1998. *AmJEpidemiol.* 2000;152(10):934-9.
74. van den Bergh ET, Gasem MH, Keuter M, Dolmans MV. Outcome in three groups of patients with typhoid fever in Indonesia between 1948 and 1990. *TropMed Int Health.* 1999;4(3):211-5.
75. Walia M, Gaind R, Paul P, Mehta R, Aggarwal P, Kalaivani M. Age-related clinical and microbiological characteristics of enteric fever in India. *TransRSocTropMed Hyg.* 2006;100(10):942-8.
76. Walia M, Gaind R, Mehta R, Paul P, Aggarwal P, Kalaivani M. Current perspectives of enteric fever: a hospital-based study from India. *AnnTropPaediatr.* 2005;25(3):161-74.
77. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. *EpidemiolInfect.* 2005;133(6):1073-9.
78. Wallace MR, Yousif AA, Mahroos GA, Mapes T, Threlfall EJ, Rowe B, et al. Ciprofloxacin versus ceftriaxone in the treatment of multiresistant typhoid fever. *EurJ ClinMicrobiolInfectDis.* 1993;12(12):907-10.

79. Yap PK, Chua CT. The haemogram in the diagnosis of acute typhoid fever--with special reference to thrombocytopenia. Singapore Med J. 1983;24(3):161-2.

• **Table S4: Spectrum of enteric fever in children by economic regions and geographical regions**

Demographics/History	High Income countries 17 studies			Europe& Central Asia 4 studies			Sub-Saharan Africa 12 studies			Middle East & North Africa 3 studies			South Asia 37 studies			East Asia & Pacific 10 studies		
	N	N	%	n	N	%	n	N	%	N	N	%	n	N	%	n	N	%
Males	262	472	56	77	131	59	574	1077	53	162	281	58	2626	4512	58	371	661	56
Duration of illness 1-2 weeks	63	161	39				109	420	26	40	150	27	556	1712	32	63	158	40
Duration of illness > 2 weeks	74	161	46				35	320	11	18	150	12	432	1863	23	5	158	3
Duration of illness > 1 week	63	134	47				131	424	31				672	1380	49	122	325	38
Chloramphenicol resistance	10	257	4	2	83	2	0	120	0				712	1075	66	66	426	15
Ampicillin/Amoxycillin resistance	19	245	8	3	83	4	6	120	5				506	781	65	41	329	12
Co-trimoxazole resistance	17	245	7	2	83	2	10	181	6				502	708	71	29	337	9
Laboratory features																		
Severe anaemia (<7.5 g/dl)							52	152	34	3	62	5	3	50	6	7	163	4
Neutrophilia(>7.5 x 10 ³ /µL)				21	24	88							72	131	55	116	177	66
Thrombocytopenia (<150 x 10 ³ /µL)	39	264	15	0	15	0	92	266	35				53	1208	4	47	173	27
Severe thrombocytopenia (<50 x 10 ³ /µL)							4	91	4	1	71	1				5	17	29
Hemolysis or hemolytic anaemia	11	127	9	1	27	4	2	144	1							7	279	3
Other signs and symptoms																		
Chills or rigors	40	106	38							4	71	6	154	503	31	83	638	13
Anorexia/refusal to feed	140	279	50	37	72	51	73	224	33				770	3184	24	246	502	49
Weakness or malaise	58	61	95	23	51	45	6	217	3				48	161	30	9	30	30
Continuous fever							3	54	6	31	150	21	262	351	75	32	137	23
Intermittent fever	96	127	76				11	35	31	5	150	3	171	354	48	52	137	38
Remittent fever	28	111	25				64	231	28	114	150	76	29	245	12	51	137	37
Stepladder fever													0	93	0	0	54	0
Chelosis	2	24	8										33	447	7	4	137	3
URTI	31	111	28				11	47	23	37	150	25	436	2765	16	19	314	6
Other complications																		
Myocarditis							1	107	1				24	908	3	12	397	3
Meningitis or meningismus	18	326	6				170	957	18	12	221	5	54	1231	4	9	488	2
Hepatitis or raised LFTs	23	56	41	13	43	30	13	296	4				199	2427	8	70	173	40
Jaundice	4	181	2	3	51	6	33	569	6				109	3485	3	12	538	2
DIC													91	336	27	1	54	2
Acute acalculouscholecystitis	1	28	4										39	970	4	4	156	3
Edema/pleural effusion/ascites	8	83	10				4	117	3				14	674	2	6	193	3
Convulsions	2	58	3	3	27	11	7	324	2				22	590	4	4	30	13
Focal neurological involvement	4	153	3	3	51	6	11	385	3	5	71	7	65	2249	3	17	458	4
Epistaxis/bleeding(unspecified)	24	240	10				3	117	3				33	772	4	16	242	7
Any abscess	1	50	2				3	164	2				2	488	0			

Osteomyelitis or septic arthritis	1	206	0		10	184	5		2	625	0	2	102	2				
Renal involvement	4	16	25		59	974	6	20	71	28	8	823	1	46	116	40		
Arthritis or arthralgia or myalgia	24	152	16	40	96	42	17	271	6	2	71	3	60	1182	5	16	384	4

Table S5: Spectrum of enteric fever in children in Latin America & Caribbean

	Latin America & Caribbean 2 studies		
	n	N	%
Males	8	13	62
Blood/bone marrow Isolates	13	13	100
Duration of illness≤ 1 week	4	4	100
Fever	4	4	100
High grade fever	0	4	0
Anorexia/refusal to feed	2	4	50
Cough	4	4	100
Nausea or vomiting	2	4	50
Diarrhea	0	4	0
Constipation	0	4	0
Hepatomegaly	0	4	0
Splenomegaly	0	4	0
Anaemia (Hb < 12 g/dL)	9	9	100
Neutrophilia(>7.5 x 10 ³ /μL)	9	9	100
Meningitis or meningismus	1	9	11
Pneumonia or chest signs	1	9	11
Hepatitis or raised LFTs	2	9	22
Epistaxis/bleeding	4	9	44
Death	2	13	15

- Table S6: Associations of clinical features in enteric fever by geographical regions (Africa* versus South Asia)

Clinical Features	Odds Ratio	95% Confidence Interval
Multi-drug resistant isolate	0.6	0.5 - 0.8
Toxicity	2.2	1.8 - 2.7
Diarrhea	2.3	2.0 - 2.6
Constipation	3.3	2.4 - 4.5
Relative bradycardia	15.3	9.6 - 24.3
Leucopenia (< 5 x 10 ³ /μL)	2.8	2.1 - 3.6
Leucocytosis (> 15)	0.6	0.4 - 0.9
Thrombocytopenia	11.5	7.9 - 16.7
Abdominal pain/tenderness	2.4	2.1 - 2.8
Abdominal distention or ileus	7.4	6.2 - 8.9
Altered mental status	1.5	1.2 - 1.8
Meningitis or meningismus	4.1	3.0 - 5.5
Pneumonia	3.1	2.6 - 3.8
GI bleeding	0.5	0.3 - 0.8
Death	3.6	2.7 - 4.9
Severe anaemia (<7.5)	5.4	1.6 - 18.1
URTI	1.7	1.2 - 2.3

Duration of illness > 2 weeks	0.4	0.3 - 0.6
Duration of illness > 1 week	0.8	0.6 - 1.0
Chills or rigors	0.2	0.1 - 0.6
Anorexia/refusal to feed	1.5	1.1 - 2.0
Weakness or malaise	0.07	0.03 - 0.2
Continuous fever	0.07	0.04 - 0.1
Remittent fever	6.5	4.2- 10.1
Intermittent fever	0.1	0.06 - 0.2
Hepatitis or raised LFTs	0.5	0.3 - 0.9
Jaundice	1.9	1.3- 2.9
Osteomyelitis or septic arthritis	17.9	3.9 - 82.5
Renal involvement	7.8	4.0 - 15.1

* Sub-Saharan Africa (12 studies) and Middle East & North Africa (3 studies)

References to Table 2, Table S4, Table S5 and Table S6: Spectrum of enteric fever in children by economic regions

High Income Countries

1. Chiu CH, Lin TY. Typhoid fever in children. Lancet. 1999;354(9194):2001-2.
2. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. Acta Paediatr Taiwan. 2000;41(1):28-32.
3. Chow CB, Wang PS, Leung NK. Typhoid fever in Hong Kong children. AustPaediatrJ. 1989;25(3):147-50.
4. Collins RN, Marine WM, Nahmias AJ. The 1964 epidemic of typhoid fever in Atlanta. Clinical and epidemiologic observations. JAMA. 1966;197(3):179-84.
5. Colon AR, Gross DR, Tamer MA. Typhoid fever in children. Pediatrics. 1975;56(4):606-9.
6. Colon AR. Gastrointestinal phenomena in childhood typhoid fever. SouthMed J. 1976;69(7):914.
7. Daoud AS, Zaki M, Pugh RN, al-Mutairi G, Beseiso R, Nasrallah AY. Clinical presentation of enteric fever: its changing pattern in Kuwait. J TropMed Hyg. 1991;94(5):341-7.
8. Del Rosario AJ, Haynes RE, Cramblett HG. Typhoid fever in children. OhioState Med J. 1971;67(6):516-20.
9. Galloway H, Clark NS, Blackhall M. Paediatric aspects of the Aberdeen typhoid outbreak. ArchDisChild. 1966;41(215):63-8.
10. Hardy G, Padfield CJ, Chadwick P, Partington MW. Typhoid outbreak in Kingston, Ont: experience with high-dose oral ampicillin. CanMed AssocJ. 1977;116(7):761-4, 7.
11. Ispahani P, Slack RC. Enteric fever and other extraintestinal salmonellosis in University Hospital, Nottingham, UK, between 1980 and 1997. EurJ ClinMicrobiolInfectDis. 2000;19(9):679-87.
12. Mallouh AA, Sa'di AR. White blood cells and bone marrow in typhoid fever. Pediatr InfectDisJ. 1987;6(6):527-9.

13. Misra S, Diaz PS, Rowley AH. Characteristics of typhoid fever in children and adolescents in a major metropolitan area in the United States. *Clin Infect Dis*. 1997;24(5):998-1000.
14. Oh HM, Masayu Z, Chew SK. Typhoid fever in hospitalized children in Singapore. *J Infect*. 1997;34(3):237-42.
15. Papaevangelou V, Syriopoulou V, Charissiadou A, Pangalis A, Mostrou G, Theodoridou M. Salmonella bacteraemia in a tertiary children's hospital. *Scand J Infect Dis*. 2004;36(8):547-51.
16. Stormon MO, McIntyre PB, Morris J, Fasher B. Typhoid fever in children: diagnostic and therapeutic difficulties. *Pediatr Infect Dis J*. 1997;16(7):713-4.
17. Su CP, Chen YC, Chang SC. Changing characteristics of typhoid fever in Taiwan. *J Microbiol Immunol Infect*. 2004;37(2):109-14.
18. Tolaymat A, Fakhreddine F, David CB, Whitworth JM. Typhoid fever in children: a forgotten disease? *South Med J*. 1979;72(2):136-8.
19. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. *Epidemiol Infect*. 2005;133(6):1073-9.

Sub Saharan Africa:

1. Ali Omer MI. Trimethoprim-sulphamethoxazole in the treatment of enteric fever. *J Trop Med Hyg*. 1975;78(7):162-6.
2. Duggan MB, Beyer L. Enteric fever in young Yoruba children. *Arch Dis Child*. 1975;50(1):67-71.
3. Ellis ME, Moosa A, Hillier V. A review of typhoid fever in South African black children. *Postgrad Med J*. 1990;66(782):1032-6.
4. Green SD, Cheesbrough JS. Salmonella bacteraemia among young children at a rural hospital in western Zaire. *Ann Trop Paediatr*. 1993;13(1):45-53.
5. Johnson AO, Aderele WI. Enteric fever in childhood. *J Trop Med Hyg*. 1981;84(1):29-35.
6. Kagwa-Nyanzi JA. Typhoid fever in African children. *J Trop Pediatr Environ Child Health*. 1971;17(3):105-9.
7. Khan M, Coovadia YM, Connolly C, Sturm AW. Influence of sex on clinical features, laboratory findings, and complications of typhoid fever. *Am J Trop Med Hyg*. 1999;61(1):41-6.
8. Laditan AA, Alausa KO. Problems in the clinical diagnosis of typhoid fever in children in the tropics. *Ann Trop Paediatr*. 1981;1(3):191-5.
9. Lepage P, Bogaerts J, Van GC, Ntahorutaba M, Nsengumuremyi F, Hitimana DG, et al. Community-acquired bacteraemia in African children. *Lancet*. 1987;1(8548):1458-61.
10. Mtove G, Amos B, von Seidlein L, Hendriksen I, Mwambuli A, Kimera J, et al. Invasive salmonellosis among children admitted to a rural Tanzanian hospital and a comparison with previous studies. *PLoS One*. 2010;5(2):e9244.
11. Mulligan TO. Typhoid fever in young children. *Br Med J*. 1971;4(5788):665-7.
12. Nesbitt A, Mirza NB. Salmonella septicaemias in Kenyan children. *J Trop Pediatr*. 1989;35(1):35-9.
13. Scragg JN, Rubidge CJ. Trimethoprim and sulphamethoxazole in typhoid fever in children. *Br Med J*. 1971;3(5777):738-41.
14. Topley JM. Mild typhoid fever. *Arch Dis Child*. 1986;61(2):164-7.

East Asia and Pacific:

1. Brian MJ. Typhoid in children in Goroka. *PNGMed J.* 1990;33(1):68-9.
 2. Choo KE, Razif A, Ariffin WA, Sepiah M, Gururaj A. Typhoid fever in hospitalized children in Kelantan, Malaysia. *AnnTropPaediatr.* 1988;8(4):207-12.
 3. Emary K, Moore CE, Chanpheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 *Salmonella enterica* serovar Typhi with intermediate susceptibility to ciprofloxacin. *Trans R Soc Trop Med Hyg.* 2012 Dec;106(12):718-24.
 4. Hoa NT, Diep TS, Wain J, Parry CM, Hien TT, Smith MD, et al. Community-acquired septicaemia in southern Viet Nam: the importance of multidrug-resistant *Salmonella typhi*. *Med J Malaysia.* 2001;56(4):478-90.
 5. Malik AS, Malik RH. Typhoid fever in Malaysian children. *Med J Malaysia.* 2001;56(4):478-90.
 6. Malik AS. Complications of bacteriologically confirmed typhoid fever in children. *J TropPediatr.* 2002;48(2):102-8
 7. Soelistiyowati S, Soenarto Y, Soesilo H, Widiarto, Widiatmodjo, Ismangoen. Typhoid fever in children. *Paediatr Indones.* 1982;22(7-8):138-46.
 8. Shwe TN, Nyein MM, Yi W, Mon A. Blood culture isolates from children admitted to Medical Unit III, Yangon Children's Hospital, 1998. *Southeast Asian J TropMed*
 9. Thisyakorn U, Mansuwan P, Taylor DN. Typhoid and paratyphoid fever in 192 hospitalized children in Thailand. *AmJ DisChild.* 1987;141(8):862-5
 10. Thisyakorn U, Mansuwan P. Comparative efficacy of mecillinam, mecillinam/amoxicillin and trimethoprim-sulfamethoxazole for treatment of typhoid fever in children. *PediatrInfectDisJ.*
 11. Wongsawat J, Pancharoen C, Thisyakorn U. Typhoid fever in children: experience in King Chulalongkorn Memorial Hospital. *J Med AssocThai.* 2002;85
 12. Yap YF, Puthucheary SD. Typhoid fever in children--a retrospective study of 54 cases from Malaysia. *Singapore Med J.* 1998;39(6):260-2.
- Europe and Central Asia:
1. Caglar MK, Ozsoylu S, Kanra G. Relative granulocytosis in childhood typhoid fever. *J Pediatr.* 1983;102(4):603-4
 2. Ciftci E, Guriz H, Derya AA, Ince E, Erdem B, Dogru U. *Salmonella* bacteraemia in Turkish children: 37 cases seen in a university hospital between 1993 and 2002.
 3. Kanra G, Yurdakok M, Secmeer G, Arvas A, Dereli O, Sarikayalar F, et al. Clinical findings of pediatric patients during an outbreak of typhoid fever in Ankara. *TurkJ*
 4. Secmeer G, Kanra G, Cemeroglu AP, Ozen H, Ceyhan M, Ecevit Z. *Salmonella typhi* infections. A 10-year retrospective study. *TurkJ Pediatr.* 1995;37(4):339-41.
 5. Tatli MM, Aktas G, Kosecik M, Yilmaz A. Treatment of typhoid fever in children with a flexible-duration of ceftriaxone, compared with 14-day treatment with chloramphenicol.
- Latin America:
1. Ferreccio C, Levine MM, Manterola A, Rodriguez G, Rivara I, Prenzel I, et al. Benign bacteremia caused by *Salmonella typhi* and *paratyphi* in children younger than 2
 2. Palacios Malmaceda PG, Vela Acosta JJ, Gutierrez AW. [Typhoid fever in children under 2 years of age]. *BolMedHospInfant Mex.* 1981;38(3):473-83
- Middle East and North Africa:

1. Grgis NI, Tribble DR, Sultan Y, Farid Z. Short course chemotherapy with cefixime in children with multidrug-resistant *Salmonella typhi* Septicaemia. *J TropPediatr.* 1995;41(6):364-5
2. Khuri-Bulos N. Enteric fevers in children. The importance of age in the varying clinical picture. *ClinPediatr (Phila).* 1981;20(7):448-52.
3. Moussa K. Enterica of childhood. *J EgyptPublic Health Assoc.* 1970;45(1):48-55.

South Asia:

1. Agarwal KS, Singh SK, Kumar N, Srivastav R, Rajkumar A. A study of current trends in enteric fever. *J CommunDis.* 1998;30(3):171-4.
2. Ahmad KA, Khan LH, Roshan B, Bhutta ZA. Factors associated with typhoid relapse in the era of multiple drug resistant strains. *J Infect Dev Ctries.* 2011 Oct;5(10):727-31.
3. Arora RK, Gupta A, Joshi NM, Kataria VK, Lall P, Anand AC. Multidrug resistant typhoid fever: study of an outbreak in Calcutta. *Indian Pediatr.* 1992;29(1):61-6.
4. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. *ArchDisChild.* 1996;75(3):214-7.
5. Bhutta ZA. Fulminant hepatic failure with typhoid fever in childhood. *J Pak Med Assoc.* 1991;41(6):123-6.
6. Bhutta ZA, Mansoorali N, Hussain R. Plasma cytokines in paediatric typhoidal salmonellosis: correlation with clinical course and outcome. *J Infect.* 1997;35(3):253-6.
7. Bhutta ZA, Naqvi SH. Hepatic dysfunction in paediatric typhoidal salmonellosis. *AnnTropPaediatr.* 1991;11(3):203-5.
8. Bhutta ZA, Naqvi SH, Durrani S, Suria A. Chloramphenicol therapy of typhoid fever and its relationship to hepatic dysfunction. *JTropPediatr.* 1991;37(6):320-2.
9. Bhutta ZA, Naqvi SH, Razzaq RA, Farooqui BJ. Multidrug-resistant typhoid in children: presentation and clinical features. *RevInfectDis.* 1991;13(5):832-6.
10. Bhutta ZA, Niazi SK, Suria A. Chloramphenicol clearance in typhoid fever: implications for therapy. *Indian J Pediatr.* 1992;59(2):213-9
11. Bhutta ZA. Enteric Fever: Risk Factors for Treatment Failure in the Era of Multidrug Resistance. UNPUBLISHED.
12. Biswal N. Neurological manifestations of typhoid fever in children. *J TropPediatr.* 1994;40(3):190.
13. Biswal N, Mathai B, Bhatia BD, Srinivasan S, Nalini P. Enteric fever: a changing perspective. *Indian Pediatr.* 1994;31(7):813-9.
14. Buch NA, Hassan MU, Kakroo DK. Enteric fever--a changing sensitivity pattern, clinical profile and outcome. *Indian Pediatr.* 1994;31(8):981-5
15. Chandra R, Srinivasan S, Nalini P, Rao RS. Multidrug resistant enteric fever. *J TropMed Hyg.* 1992;95(4):284-7.
16. Chawla V, Bhujwala RA, Chandra RK, Ghai OP. Typhoid fever in children. Clinical and bacteriological studies. *Indian J Pediatr.* 1970;37(269):239-44.
17. Dheer G, Kundra S, Singh T. Clinical and laboratory profile of enteric fever in children in northern India. *Trop Doct.* 2012 Jul;42(3):154-6.
18. Dutta P, Lahiri M, Bhattacharya SK, Saha MR, Mitra U, Rasaily R, et al. Hepatitis-like presentation in typhoid fever. *TransRSocTropMed Hyg.* 1992;86(1):92.

19. Dutta P, Rasaily R, Saha MR, Mitra U, Manna B, Chakraborty S, et al. Randomized clinical trial of furazolidone for typhoid fever in children. *Scand J Gastroenterol*. 1993;28(2):168-72.
20. Garg K, Mangal N, Mathur HC. Clinical profile of multi drug resistant typhoid fever in Jaipur City. *Indian Pediatr*. 1994;31(2):191-3.
21. Garg RA, Krashak R. Typhoid fever before two years of age. *Indian Pediatr*. 1993;30(6):805-8
22. Gulati S, Marwaha RK, Singhi S, Ayyagari A, Kumar L. Third generation cephalosporins in multi-drug resistant typhoid fever. *Indian Pediatr*. 1992;29(4):513-6.
23. Hassan M. Multiply-resistant *Salmonella typhi* in children. *Drugs*. 1995;49 Suppl 2:457-9.
24. Jagadish K, Patwari AK, Sarin SK, Prakash C, Srivastava DK, Anand VK. Hepatic manifestations in typhoid fever. *Indian Pediatr*. 1994;31(7):807-11.
25. Jain S, Chitnis DS, Sham A, Rathi S, Inamdar S, Rindani GJ. Outbreak of chloramphenicol resistant typhoid fever. *Indian Pediatr*. 1987;24(3):193-7
26. Kabra SK, Madhulika, Talati A, Soni N, Patel S, Modi RR. Multidrug-resistant typhoid fever. *Trop Doct*. 2000;30(4):195-7.
27. Kapoor JP, Mohan M, Talwar V, Daral TS, Bhargava SK. Typhoid fever in young children. *Indian Pediatr*. 1985;22(11):811-3.
28. Khan MI, Soofi SB, Ochiai RL, Khan MJ, Sahito SM, Habib MA, et al. Epidemiology, clinical presentation, and patterns of drug resistance of *Salmonella Typhi* in Karachi, Pakistan. *J Infect Dev Ctries*. 2012 Oct;6(10):704-14.
29. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramchandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: clinical profile and therapy. *Indian Pediatr*. 1991;28(4):357-61.
30. Kulkarni ML, Rego SJ. Acute acalculous cholecystitis in typhoid fever. *Indian Pediatr*. 1995;32(7):829-30
31. Kumar R, Gupta N. Multidrug-resistant typhoid fever. *Indian J Pediatr*. 2007;74(1):39-42.
32. Leung DT, Bogetz J, Itoh M, Ganapathi L, Pietroni MA, Ryan ET, et al. Factors associated with encephalopathy in patients with *Salmonella enterica* serotype Typhi bacteremia presenting to a diarrheal hospital in Dhaka, Bangladesh. *Am J Trop Med Hyg*. 2012 Apr;86(4):698-702
33. Libov A. Enteric fever in children. *Indian J Pediatr*. 1960;27:125-8. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr*. 1991;28(10):1171-4.
34. Memon IA, Billoo AG, Memon HI. Cefixime: an oral option for the treatment of multidrug-resistant enteric fever in children. *South Med J*. 1997;90(12):1204-7.
35. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr*. 1991;28(10):1171-4.
36. Naeemullah S. A 3 year study of children with enteric fever. *Pakistan Pediatric Journal*. 1994;18(4):165-9.
37. Owais A, Sultana S, Zaman U, Rizvi A, Zaidi AK. Incidence of typhoid bacteremia in infants and young children in southern coastal Pakistan. *Pediatr Infect Dis J*. 2010 Nov;29(11):1035-9.
38. Pradhan R, Shrestha U, Gautam SC, Thorson S, Shrestha K, Yadav BK, et al. Bloodstream infection among children presenting to a general hospital outpatient clinic in urban Nepal. *PLoS One*. 2012;7(10):e47531

39. Rajajee S, Anandi TB, Subha S, Vatsala BR. Patterns of resistant *Salmonella typhi* infection in infants. *JTropPediatr.* 1995;41(1):52-4.
40. Raman TS, Krishnamurthy L, Menon PK, Singh D, Jayaprakash DG. Clinical profile and therapy in enteric fever. *Indian Pediatr.* 1994;31(2):196-9.
41. Ramanan A, Pandit N, Yeshwanth M. Unusual complications in a multidrug resistant *Salmonella typhi* outbreak. *Indian Pediatr.* 1992;29(1):118-20.
42. Rasaily R, Dutta P, Saha MR, Mitra U, Lahiri M, Pal SC. Multi-drug resistant typhoid fever in hospitalised children. Clinical, bacteriological and epidemiological profiles.
43. Rathore MH, Bux D, Hasan M. Multidrug-resistant *Salmonella typhi* in Pakistani children: clinical features and treatment. *SouthMed J.* 1996;89(2):235-7.
44. Sethuraman S, Mahamood M, Kareem S. Furazolidone in multi-resistant childhood typhoid fever. *AnnTropPaediatr.* 1994;14(4):321-4.
45. Sharma A, Gathwala G. Clinical profile and outcome in enteric fever. *Indian Pediatr.* 1993;30(1):47-50.
46. Shetty AK, Mital SR, Bahrainwala AH, Khubchandani RP, Kumta NB. Typhoid hepatitis in children. *J TropPediatr.* 1999;45(5):287-90.
47. Siddiqui FJ, Rabbani F, Hasan R, Nizami SQ, Bhutta ZA. Typhoid fever in children: some epidemiological considerations from Karachi, Pakistan. *Int J*
48. Singh H, Raizada N. Chloramphenicol resistant typhoid fever. *Indian Pediatr.* 1991;28(4):433.
49. Sinha A, Sazawal S, Kumar R, Sood S, Reddaiah VP, Singh B, et al. Typhoid fever in children aged less than 5 years. *Lancet.* 1999;354(9180):734-7.
50. Smith M. Typhoid admissions in childhood. *TropDoct.* 1986;16(4):150-2
51. Takkar VP, Kumar R, Takkar R, Khurana S. Resurgence of chloramphenicol sensitive *Salmonella typhi*. *Indian Pediatr.* 1995;32(5):586-7.
52. Thambidorai CR, Shyamala J, Sarala R, Vatsala RB, Tamizhisai S. Acute acalculous cholecystitis associated with enteric fever in children. *Pediatr Infect Dis J.* 1995 Sep;14(9):812-3.
53. Verma PS, Mullick P, Ghosh S. Neurological manifestations of enteric fever: clinical profile and correlation with ultimate outcome. *Indian Pediatr.* 1972;9(11):681-5.
54. Walia M, Gaind R, Mehta R, Paul P, Aggarwal P, Kalaivani M. Current perspectives of enteric fever: a hospital-based study from India. *AnnTropPaediatr.* 2005;25(3):161-74.
55. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr.* 1991;28(10):1171-4.

- Table S7: Enteric fever in children in hospital-based versus community-based setting

Demographics/History	Hospital-based 80 studies			Community-based or health center (out-patients) 5 studies		
	N	N	%	n	N	%
Males	4075	7142	57	2	4	50
Duration of illness within 1-2 weeks	831	2601	32			

Duration of illness > 2 weeks	564	2652	21			
Duration of illness > 1 week	988	2263	44			
Chloramphenicol resistance	780	1895	41	10	28	36
Ampicillin/Amoxycillin resistance	567	1520	37			
Co-trimoxazole resistance	547	1516	36			
Laboratory features						
Severe anaemia (<7.5 g/dl)	53	393	13			
Neutrophilia(>7.5 x 10 ³ /µL)	218	341	64			
Thrombocytopenia (<150 x 10 ³ /µL)	231	1926	12			
Severe thrombocytopenia (<50 x 10 ³ /µL)	10	179	6			
Hemolysis or hemolytic anaemia	21	577	4			
Other signs and symptoms						
Chills or rigors	281	1318	21			
Anorexia/refusal to feed	1264	4212	30	4	53	8
Weakness/dizziness or malaise	142	471	30	2	49	4
Continuous fever	328	692	47			
Intermittent fever	335	803	42			
Remittent fever	286	874	33			
Stepladder fever	0	147	0			
Chelosis	39	608	6			
URTI	532	3371	16	2	16	13
Other complications						
Myocarditis	37	1412	3			
Meningitis or meningismus	260	3160	8			
Hepatitis or raised LFTs	320	3004	11			
Jaundice	156	4790	3			
DIC	92	390	24			
Acute acalculouscholecystitis	44	1154	4			
Edema or pleural effusion or ascites	32	1067	3			
Convulsions	103	3329	3			
Focal neurological involvement	38	1029	4			
Epistaxis/bleeding (unspecified)	79	1342	6			
Any abscess	6	702	1			
Osteomyelitis or septic arthritis	11	1083	1			
Renal involvement	137	2000	7			
Arthritis or arthralgia or myalgia	159	2156	7			

- Table S8: Associations of clinical features in enteric fever in children by setting (hospital-based versus community/health-center)

Clinical Features	Odds Ratio	95% Confidence Interval
Pre-treatment antibiotics received	2.8	2.0 - 4.0
Multidrug resistant isolates	1.7	1.3 - 2.1
High - grade fever	4.7	3.5 - 6.4
Headache	0.3	0.2 - 0.4
Toxicity	0.3	0.1 - 0.6
Nausea/vomiting	2.0	1.6 - 2.6
Cough	0.7	0.5 - 0.9
Diarrhea	5.2	3.8 - 7.2

Constipation	4.2	1.9 - 9.6
Hepatomegaly	7.1	4.2 - 12.0
Splenomegaly	2.7	1.7 - 4.1
Abdominal pain/tenderness	0.5	0.4 - 0.6
Abdominal distention or ileus	7.4	2.7 - 20.0
GI bleeding	9.0	1.2 - 64.3
Duration of illness ≤ 1 week	1.8	1.3 - 2.4
Fluoroquinolone resistance	0.03	0.01 - 0.08

References for Table 3 , Table S7, and Table S8: Typhoid fever in children in hospital-based vs. community/health-center

Community Based:

1. Ferreccio C, Levine MM, Manterola A, Rodriguez G, Rivara I, Prenzel I, et al. Benign bacteremia caused by *Salmonella typhi* and *paratyphi* in children younger than 2 years. *J Pediatr.* 1984;104(6):899-901.
2. Khan MI, Soofi SB, Ochiai RL, Khan MJ, Sahito SM, Habib MA, et al. Epidemiology, clinical presentation, and patterns of drug resistance of *Salmonella Typhi* in Karachi, Pakistan. *J Infect Dev Ctries.* 2012 Oct;6(10):704-14.
3. Owais A, Sultana S, Zaman U, Rizvi A, Zaidi AK. Incidence of typhoid bacteremia in infants and young children in southern coastal Pakistan. *Pediatr Infect Dis J.* 2010
4. Pradhan R, Shrestha U, Gautam SC, Thorson S, Shrestha K, Yadav BK, et al. Bloodstream infection among children presenting to a general hospital outpatient clinic in urban Nepal. *PLoS One.* 2012;7(10):e47531
5. Sinha A, Sazawal S, Kumar R, Sood S, Reddaiah VP, Singh B, et al. Typhoid fever in children aged less than 5 years. *Lancet.* 1999;354(9180):734-7.
6. Siddiqui FJ, Rabbani F, Hasan R, Nizami SQ, Bhutta ZA. Typhoid fever in children: some epidemiological considerations from Karachi, Pakistan. *Int J Infect Dis.* 2006;10(3):215-22.

Hospital Based:

1. Ali Omer MI. Trimethoprim-sulphamethoxazole in the treatment of enteric fever. *J Trop Med Hyg.* 1975;78(7):162-6.
2. Ahmad KA, Khan LH, Roshan B, Bhutta ZA. Factors associated with typhoid relapse in the era of multiple drug resistant strains. *J Infect Dev Ctries.* 2011 Oct;5(10):727-31.
3. Arora RK, Gupta A, Joshi NM, Kataria VK, Lall P, Anand AC. Multidrug resistant typhoid fever: study of an outbreak in Calcutta. *Indian Pediatr.* 1992;29(1):61-6.
4. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. *Arch Dis Child.* 1996;75(3):214-7.
5. Bhutta ZA. Fulminant hepatic failure with typhoid fever in childhood. *J Pak Med Assoc.* 1991;41(6):123-6.
6. Bhutta ZA, Mansoorali N, Hussain R. Plasma cytokines in paediatric typhoidal salmonellosis: correlation with clinical course and outcome. *J Infect.* 1997;35(3):253-6.

7. Bhutta ZA, Naqvi SH. Hepatic dysfunction in paediatric typhoidal salmonellosis. *AnnTropPaediatr.* 1991;11(3):203-5.
8. Bhutta ZA, Naqvi SH, Durrani S, Suria A. Chloramphenicol therapy of typhoid fever and its relationship to hepatic dysfunction. *JTropPediatr.* 1991;37(6):320-2.
9. Bhutta ZA, Naqvi SH, Razzaq RA, Farooqui BJ. Multidrug-resistant typhoid in children: presentation and clinical features. *RevInfectDis.* 1991;13(5):832-6.
10. Bhutta ZA, Niazi SK, Suria A. Chloramphenicol clearance in typhoid fever: implications for therapy. *Indian J Pediatr.* 1992;59(2):213-9
11. Bhutta ZA. Enteric Fever: Risk Factors for Treatment Failure in the Era of Multidrug Resistance. UNPUBLISHED.
12. Biswal N. Neurological manifestations of typhoid fever in children. *J TropPediatr.* 1994;40(3):190.
13. Biswal N, Mathai B, Bhatia BD, Srinivasan S, Nalini P. Enteric fever: a changing perspective. *Indian Pediatr.* 1994;31(7):813-9.
14. Brian MJ. Typhoid in children in Goroka. *PNGMed J.* 1990;33(1):68-9.
15. Buch NA, Hassan MU, Kakroo DK. Enteric fever--a changing sensitivity pattern, clinical profile and outcome. *Indian Pediatr.* 1994;31(8):981-5
16. Caglar MK, Ozsoylu S, Kanra G. Relative granulocytosis in childhood typhoid fever. *J Pediatr.* 1983;102(4):603-4
17. Chandra R, Srinivasan S, Nalini P, Rao RS. Multidrug resistant enteric fever. *J TropMed Hyg.* 1992;95(4):284-7.
18. Chawla V, Bhujwala RA, Chandra RK, Ghai OP. Typhoid fever in children. Clinical and bacteriological studies. *Indian J Pediatr.* 1970;37(269):239-44.
19. Chiu CH, Lin TY. Typhoid fever in children. *Lancet.* 1999;354(9194):2001-2.
20. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. *Acta PaediatrTaiwan.* 2000;41(1):28-32.
21. Choo KE, Razif A, Ariffin WA, Sepiah M, Gururaj A. Typhoid fever in hospitalized children in Kelantan, Malaysia. *AnnTropPaediatr.* 1988;8(4):207-12.
22. Chow CB, Wang PS, Leung NK. Typhoid fever in Hong Kong children. *AustPaediatrJ.* 1989;25(3):147-50.
23. Ciftci E, Guriz H, Derya AA, Ince E, Erdem B, Dogru U. Salmonella bacteraemia in Turkish children: 37 cases seen in a university hospital between 1993 and 2002. *AnnTropPaediatr.* 2004;24(1):75-80..
24. Collins RN, Marine WM, Nahmias AJ. The 1964 epidemic of typhoid fever in Atlanta. Clinical and epidemiologic observations. *JAMA.* 1966;197(3):179-84.
25. Colon AR, Gross DR, Tamer MA. Typhoid fever in children. *Pediatrics.* 1975;56(4):606-9.
26. Colon AR. Gastrointestinal phenomena in childhood typhoid fever. *SouthMed J.* 1976;69(7):914.
27. Daoud AS, Zaki M, Pugh RN, al-Mutairi G, Beseiso R, Nasrallah AY. Clinical presentation of enteric fever: its changing pattern in Kuwait. *J TropMed Hyg.* 1991;94(5):341-7
28. Del Rosario AJ, Haynes RE, Cramblett HG. Typhoid fever in children. *OhioState Med J.* 1971;67(6):516-20.
29. Dheer G, Kundra S, Singh T. Clinical and laboratory profile of enteric fever in children in northern India. *Trop Doct.* 2012 Jul;42(3):154-6.
30. Duggan MB, Beyer L. Enteric fever in young Yoruba children. *ArchDisChild.* 1975;50(1):67-71.

31. Dutta P, Lahiri M, Bhattacharya SK, Saha MR, Mitra U, Rasaily R, et al. Hepatitis-like presentation in typhoid fever. *TransRSocTropMed Hyg.* 1992;86(1):92.
32. Dutta P, Rasaily R, Saha MR, Mitra U, Manna B, Chakraborty S, et al. Randomized clinical trial of furazolidone for typhoid fever in children. *ScandJGastroenterol.* 1993;28(2):168-72.
33. Ellis ME, Moosa A, Hillier V. A review of typhoid fever in South African black children. *PostgradMed J.* 1990;66(782):1032-6..
34. Emary K, Moore CE, Chanpheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 *Salmonella enterica* serovar Typhi with intermediate susceptibility to ciprofloxacin. *Trans R Soc Trop Med Hyg.* 2012 Dec;106(12):718-24.
35. Galloway H, Clark NS, Blackhall M. Paediatric aspects of the Aberdeen typhoid outbreak. *ArchDisChild.* 1966;41(215):63-8.
36. Garg K, Mangal N, Mathur HC. Clinical profile of multi drug resistant typhoid fever in Jaipur City. *Indian Pediatr.* 1994;31(2):191-3.
37. Garg RA, Krashak R. Typhoid fever before two years of age. *Indian Pediatr.* 1993;30(6):805-8
38. Girgis NI, Tribble DR, Sultan Y, Farid Z. Short course chemotherapy with cefixime in children with multidrug-resistant *Salmonella typhi* Septicaemia. *J TropPediatr.* 1995;41(6):364-5.
39. Gulati S, Marwaha RK, Singhi S, Ayyagari A, Kumar L. Third generation cephalosporins in multi-drug resistant typhoid fever. *Indian Pediatr.* 1992;29(4):513-6.
40. Hardy G, Padfield CJ, Chadwick P, Partington MW. Typhoid outbreak in Kingston, Ont: experience with high-dose oral ampicillin. *CanMed AssocJ.* 1977;116(7):761-4, 7.
41. Hassan M. Multiply-resistant *Salmonella typhi* in children. *Drugs.* 1995;49 Suppl 2:457-9.
42. Hoa NT, Diep TS, Wain J, Parry CM, Hien TT, Smith MD, et al. Community-acquired septicaemia in southern Viet Nam: the importance of multidrug-resistant *Salmonella typhi*. *TransRSocTropMedHyg.* 1998;92(5):503-8.
43. Ispahani P, Slack RC. Enteric fever and other extraintestinal salmonellosis in University Hospital, Nottingham, UK, between 1980 and 1997. *EurJ ClinMicrobiolInfectDis.* 2000;19(9):679-87.
44. Jagadish K, Patwari AK, Sarin SK, Prakash C, Srivastava DK, Anand VK. Hepatic manifestations in typhoid fever. *Indian Pediatr.* 1994;31(7):807-11.
45. Jain S, Chitnis DS, Sham A, Rathi S, Inamdar S, Rindani GJ. Outbreak of chloramphenicol resistant typhoid fever. *Indian Pediatr.* 1987;24(3):193-7
46. Johnson AO, Aderele WI. Enteric fever in childhood. *J TropMed Hyg.* 1981;84(1):29-35.
47. Kabra SK, Madhulika, Talati A, Soni N, Patel S, Modi RR. Multidrug-resistant typhoid fever. *TropDoct.* 2000;30(4):195-7.
48. Kanra G, Yurdakok M, Secmeer G, Arvas A, Dereli O, Sarikayalar F, et al. Clinical findings of pediatric patients during an outbreak of typhoid fever in Ankara. *TurkJ Pediatr.* 1985;27(1):11-5.
49. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramchandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: clinical profile and therapy. *Indian Pediatr.* 1991;28(4):357-61.
50. Kumar R, Gupta N. Multidrug-resistant typhoid fever. *Indian J Pediatr.* 2007;74(1):39-42.

51. Kapoor JP, Mohan M, Talwar V, Daral TS, Bhargava SK. Typhoid fever in young children. *Indian Pediatr.* 1985;22(11):811-3.
52. Kulkarni ML, Rego SJ. Acute acalculous cholecystitis in typhoid fever. *Indian Pediatr.* 1995;32(7):829-30
53. Khan M, Coovadia YM, Connolly C, Sturm AW. Influence of sex on clinical features, laboratory findings, and complications of typhoid fever. *Am J Trop Med Hyg.* 1999;61(1):41-6.
54. Khuri-Bulos N. Enteric fevers in children. The importance of age in the varying clinical picture. *Clin Pediatr (Phila).* 1981;20(7):448-52
55. Kagwa-Nyanzi JA. Typhoid fever in African children. *J Trop Pediatr Environ Child Health.* 1971;17(3):105-9.
56. Laditan AA, Alausa KO. Problems in the clinical diagnosis of typhoid fever in children in the tropics. *Ann Trop Paediatr.* 1981;1(3):191-5.
57. Lepage P, Bogaerts J, Van GC, Ntahorutaba M, Nsengumuremyi F, Hitimana DG, et al. Community-acquired bacteraemia in African children. *Lancet.* 1987;1(8548):1458-61.
58. Leung DT, Bogetz J, Itoh M, Ganapathi L, Pietroni MA, Ryan ET, et al. Factors associated with encephalopathy in patients with *Salmonella enterica* serotype Typhi bacteremia presenting to a diarrheal hospital in Dhaka, Bangladesh. *Am J Trop Med Hyg.* 2012 Apr;86(4):698-702.
59. Libov A. Enteric fever in children. *Indian J Pediatr.* 1960;27:125-8.
60. Malik AS. Complications of bacteriologically confirmed typhoid fever in children. *J Trop Pediatr.* 2002;48(2):102-8.
61. Malik AS, Malik RH. Typhoid fever in Malaysian children. *Med J Malaysia.* 2001;56(4):478-90.
62. Mallouh AA, Sa'di AR. White blood cells and bone marrow in typhoid fever. *Pediatr Infect Dis J.* 1987;6(6):527-9.
63. Memon IA, Billoo AG, Memon HI. Cefixime: an oral option for the treatment of multidrug-resistant enteric fever in children. *South Med J.* 1997;90(12):1204-7.
64. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr.* 1991;28(10):1171-4.
65. Misra S, Diaz PS, Rowley AH. Characteristics of typhoid fever in children and adolescents in a major metropolitan area in the United States. *Clin Infect Dis.* 1997;24(5):998-
66. Moussa K. Enterica of childhood. *J Egypt Public Health Assoc.* 1970;45(1):48-55.
67. Mtové G, Amos B, von Seidlein L, Hendriksen I, Mwambuli A, Kimera J, et al. Invasive salmonellosis among children admitted to a rural Tanzanian hospital and a comparison with previous studies. *PLoS One.* 2010;5(2):e9244.
68. Mulligan TO. Typhoid fever in young children. *Br Med J.* 1971;4(5788):665-7.
69. Naeemullah S. A 3 year study of children with enteric fever. *Pakistan Pediatric Journal.* 1994;18(4):165-9.
70. Nesbitt A, Mirza NB. *Salmonella* septicaemias in Kenyan children. *J Trop Pediatr.* 1989;35(1):35-9.
71. Oh HM, Masayu Z, Chew SK. Typhoid fever in hospitalized children in Singapore. *J Infect.* 1997;34(3):237-42.
72. Palacios Malmaceda PG, Vela Acosta JJ, Gutierrez AW. [Typhoid fever in children under 2 years of age]. *Bol Med Hosp Infant Mex.* 1981;38(3):473-83.

73. Papaevangelou V, Syriopoulou V, Charissiadou A, Pangalis A, Mostrou G, Theodoridou M. Salmonella bacteraemia in a tertiary children's hospital. *ScandJ InfectDis.* 2004;36(8):547-51.
74. Rajajee S, Anandi TB, Subha S, Vatsala BR. Patterns of resistant *Salmonella typhi* infection in infants. *JTropPediatr.* 1995;41(1):52-4.
75. Raman TS, Krishnamurthy L, Menon PK, Singh D, Jayaprakash DG. Clinical profile and therapy in enteric fever. *Indian Pediatr.* 1994;31(2):196-9.
76. Ramanan A, Pandit N, Yeshwanth M. Unusual complications in a multidrug resistant *Salmonella typhi* outbreak. *Indian Pediatr.* 1992;29(1):118-20.
77. Rasaily R, Dutta P, Saha MR, Mitra U, Lahiri M, Pal SC. Multi-drug resistant typhoid fever in hospitalised children. Clinical, bacteriological and epidemiological profiles. *EurJ Epidemiol.* 1994;10(1):41-6.
78. Rathore MH, Bux D, Hasan M. Multidrug-resistant *Salmonella typhi* in Pakistani children: clinical features and treatment. *SouthMed J.* 1996;89(2):235-7.
79. Scragg JN, Rubidge CJ. Trimethoprim and sulphamethoxazole in typhoid fever in children. *BrMed J.* 1971;3(5777):738-41.
80. Secmeer G, Kanra G, Cemeroglu AP, Ozen H, Ceyhan M, Ecevit Z. *Salmonella typhi* infections. A 10-year retrospective study. *TurkJ Pediatr.* 1995;37(4):339-41.
81. Sethuraman S, Mahamood M, Kareem S. Furazolidone in multi-resistant childhood typhoid fever. *AnnTropPaediatr.* 1994;14(4):321-4.
82. Sharma A, Gathwala G. Clinical profile and outcome in enteric fever. *Indian Pediatr.* 1993;30(1):47-50.
83. Shetty AK, Mital SR, Bahrainwala AH, Khubchandani RP, Kumta NB. Typhoid hepatitis in children. *J TropPediatr.* 1999;45(5):287-90.
84. Shwe TN, Nyein MM, Yi W, Mon A. Blood culture isolates from children admitted to Medical Unit III, Yangon Children's Hospital, 1998. *Southeast Asian J TropMed Public Health.*
85. Singh H, Raizada N. Chloramphenicol resistant typhoid fever. *Indian Pediatr.* 1991;28(4):433.
86. Smith M. Typhoid admissions in childhood. *TropDoct.* 1986;16(4):150-2
87. Soelistowati S, Soenarto Y, Soesilo H, Widiarto, Widiatmodjo, Ismangoen. Typhoid fever in children. *PaediatrIndones.* 1982;22(7-8):138-46.
88. Stormon MO, McIntyre PB, Morris J, Fasher B. Typhoid fever in children: diagnostic and therapeutic difficulties. *Pediatr InfectDisJ.* 1997;16(7):713-4.
89. Su CP, Chen YC, Chang SC. Changing characteristics of typhoid fever in Taiwan. *J MicrobiolImmunolInfect.* 2004;37(2):109-14.
90. Takkar VP, Kumar R, Takkar R, Khurana S. Resurgence of chloramphenicol sensitive *Salmonella typhi*. *Indian Pediatr.* 1995;32(5):586-7.
91. Thambidorai CR, Shyamala J, Sarala R, Vatsala RB, Tamizhisai S. Acute acalculous cholecystitis associated with enteric fever in children. *Pediatr InfectDisJ.* 1995;14(9):812-3.
92. Tatli MM, Aktas G, Kosecik M, Yilmaz A. Treatment of typhoid fever in children with a flexible-duration of ceftriaxone, compared with 14-day treatment with chloramphenicol.
93. Thisakorn U, Mansuwan P. Comparative efficacy of mecillinam, mecillinam/amoxicillin and trimethoprim-sulfamethoxazole for treatment of typhoid fever in children. *PediatrInfectDisJ.* 1992;11(11):979-80.
94. Thisakorn U, Mansuwan P, Taylor DN. Typhoid and paratyphoid fever in 192 hospitalized children in Thailand. *AmJ DisChild.* 1987;141(8):862-5.

95. Tolaymat A, Fakhreddine F, David CB, Whitworth JM. Typhoid fever in children: a forgotten disease? *SouthMed J.* 1979;72(2):136-8.
96. Topley JM. Mild typhoid fever. *ArchDisChild.* 1986;61(2):164-7.
97. Verma PS, Mullick P, Ghosh S. Neurological manifestations of enteric fever: clinical profile and correlation with ultimate outcome. *Indian Pediatr.* 1972;9(11):681-5.
98. Walia M, Gaind R, Mehta R, Paul P, Aggarwal P, Kalaivani M. Current perspectives of enteric fever: a hospital-based study from India. *AnnTropPaediatr.* 2005;25(3):161-74.
99. Walia M, Gaind R, Paul P, Mehta R, Aggarwal P, Kalaivani M. Age-related clinical and microbiological characteristics of enteric fever in India. *TransRSocTropMed Hyg.* 2006;100(10):942-8.
100. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. *EpidemiolInfect.* 2005;133(6):1073-9.
101. Wongsawat J, Pancharoen C, Thisyakorn U. Typhoid fever in children: experience in King Chulalongkorn Memorial Hospital. *J Med AssocThai.* 2002;85(12):1247-50.
102. Yap YF, Puthucheary SD. Typhoid fever in children--a retrospective study of 54 cases from Malaysia. *Singapore Med J.* 1998;39(6):260-2.

- Table S9: Typhoid fever in children in with multi-drug resistant versus sensitive strains of salmonella**

Demographics/History	Multi-drug resistant 13 studies			Sensitive 22 studies		
	n	N	%	n	N	%
Males	358	601	60	864	1500	58
Duration of illness 1 - 2 weeks	80	272	29	208	897	23
Duration of illness >2 weeks	83	272	31	164	897	18
Duration of illness > 1 week	208	334	62	407	1034	39
Ampicillin/Amoxyillin resistance	125	125	100	9	186	5
Co-trimoxazole resistance	125	125	100	14	186	8
Laboratory features						
Severe anaemia (<7.5 g/dl)	2	39	5	13	75	17
Thrombocytopenia (<150 x 10 ³ /µL)	3	62	5	39	119	33
Other signs and symptoms						
Chills or rigors	33	116	28	22	192	11
Anorexia/refusal to feed	27	77	35	85	253	34
Continuous fever	81	81	100	66	171	39
Intermittent fever	44	62	71	52	137	38
Remittent fever	11	11	100	51	137	37
Chelosis	5	112	4	6	161	4
Other complications						
Myocarditis	10	157	6	5	313	2
Meningitis or meningismus	3	54	6	6	210	3
Hepatitis or raised LFTs	59	413	14	22	210	10
Jaundice	43	416	10	12	317	4
Acute acalculouscholecystitis	4	80	5	3	122	2
Edema or pleural effusion or ascites	5	181	3	0	11	0
Convulsions	2	54	4	8	233	3
Epistaxis/bleeding (unspecified)	2	39	5	3	62	5
Any abscess	0	62	0	1	47	2

Renal involvement	3	95	3	46	116	40
Arthritis or arthralgia or myalgia	1	77	1	45	233	19

- Table S10: Associations of clinical features in enteric fever in children with multi-drug resistant versus sensitive strains of *S. Typhi* and *S. Paratyphi*

Clinical Features	Odds Ratio	95% Confidence Interval
Duration of illness ≤ 1 week	0.4	0.3 - 0.5
Headache	2.4	1.7 - 3.3
Toxicity	2.1	1.6 - 2.6
Hepatomegaly	2.3	1.9 - 2.7
Splenomegaly	2.4	2.0- 2.9
Abdominal pain/tenderness	2.4	2.0 – 3.0
Abdominal distention or ileus	2.6	1.7 - 4.1
Anaemia	1.6	1.3 - 2.0
Leucopenia	0.5	0.3 - 0.8
Shock	2.9	1.2 - 7.3
Pneumonia	2.2	1.3 - 3.7
GI bleeding	2.3	1.1 - 4.5
Relapse	0.3	0.1 - 0.7
High grade fever	0.6	0.5 - 0.8
Bradycardia or relative bradycardia	9.1	2.3 - 36.4
Dehydration	10.9	3.3 - 36.0
Duration of illness > 1 week	2.7	2.1 - 3.4
Jaundice	2.9	1.5 - 5.7
Renal involvement	0.05	0.01 - 0.2
Duration of illness >2 weeks	2.0	1.4 - 2.7
Thrombocytopenia (<150)	0.1	0.03 - 0.3
Chills or rigors	3.1	1.7 - 5.6
Myocarditis	4.2	1.4 - 12.5
Arthritis or arthralgia or myalgia	0.05	0.01 - 0.4

References to Table 4, Table S9 and Table S10: Typhoid fever in children with multi-drug resistant versus sensitive strains of salmonella

Sensitive

1. Agarwal KS, Singh SK, Kumar N, Srivastav R, Rajkumar A. A study of current trends in enteric fever. J CommunDis. 1998;30(3):171-4.
2. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. ArchDisChild. 1996;75(3):214-7.
3. Bhutta ZA. Enteric Fever: Risk Factors for Treatment Failure in the Era of Multidrug Resistance. UNPUBLISHED.

4. Buch NA, Hassan MU, Kakroo DK. Enteric fever--a changing sensitivity pattern, clinical profile and outcome. *Indian Pediatr.* 1994;31(8):981-5.
5. Chandra R, Srinivasan S, Nalini P, Rao RS. Multidrug resistant enteric fever. *J TropMed Hyg.* 1992;95(4):284-7.
6. Choo KE, Razif A, Ariffin WA, Sepiah M, Gururaj A. Typhoid fever in hospitalized children in Kelantan, Malaysia. *AnnTropPaediatr.* 1988;8(4):207-12.
7. Hardy G, Padfield CJ, Chadwick P, Partington MW. Typhoid outbreak in Kingston, Ont: experience with high-dose oral ampicillin. *CanMed AssocJ.* 1977;116(7):761-4, 7.
8. Jain S, Chitnis DS, Sham A, Rathi S, Inamdar S, Rindani GJ. Outbreak of chloramphenicol resistant typhoid fever. *Indian Pediatr.* 1987;24(3):193-7.
9. Kabra SK, Madhulika, Talati A, Soni N, Patel S, Modi RR. Multidrug-resistant typhoid fever. *TropDoct.* 2000;30(4):195-7.
10. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramchandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: clinical profile and therapy. *Indian Pediatr.* 1991;28(4):357-61.
11. Lepage P, Bogaerts J, Van GC, Ntahorutaba M, Nsengumuremyi F, Hitimana DG, et al. Community-acquired bacteraemia in African children. *Lancet.* 1987;1(8548):1458-61.
12. Malik AS. Complications of bacteriologically confirmed typhoid fever in children. *J TropPediatr.* 2002;48(2):102-8.
13. Malik AS, Malik RH. Typhoid fever in Malaysian children. *Med J Malaysia.* 2001;56(4):478-90.
14. ishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr.* 1991;28(10):1171-4.
15. Nesbitt A, Mirza NB. *Salmonella* septicaemias in Kenyan children. *J TropPediatr.* 1989;35(1):35-9.
16. Papaevangelou V, Syriopoulou V, Charissiadou A, Pangalis A, Mostrou G, Theodoridou M. *Salmonella* bacteraemia in a tertiary children's hospital. *ScandJ InfectDis.* 2004;36(8):547-51.
17. Raman TS, Krishnamurthy L, Menon PK, Singh D, Jayaprakash DG. Clinical profile and therapy in enteric fever. *Indian Pediatr.* 1994;31(2):196-9.
18. Sharma A, Gathwala G. Clinical profile and outcome in enteric fever. *Indian Pediatr.* 1993;30(1):47-50.
19. Su CP, Chen YC, Chang SC. Changing characteristics of typhoid fever in Taiwan. *J MicrobiolImmunolInfect.* 2004;37(2):109-14.
20. Takkar VP, Kumar R, Takkar R, Khurana S. Resurgence of chloramphenicol sensitive *Salmonella typhi*. *Indian Pediatr.* 1995;32(5):586-7.
21. Tatli MM, Aktas G, Kosecik M, Yilmaz A. Treatment of typhoid fever in children with a flexible-duration of ceftriaxone, compared with 14-day treatment with chloramphenicol. *IntJAntimicrobAgents.* 2003;21(4):350-3.
22. Thisyakorn U, Mansuwan P. Comparative efficacy of mecillinam, mecillinam/amoxicillin and trimethoprim-sulfamethoxazole for treatment of typhoid fever in children. *PediatrInfectDisJ.* 1992;11(11):979-80.
23. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. *EpidemiolInfect.* 2005;133(6):1073-9.
24. Wongsawat J, Pancharoen C, Thisyakorn U. Typhoid fever in children: experience in King Chulalongkorn Memorial Hospital. *J Med AssocThai.* 2002;85(12):1247-50.

MDR

1. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. *ArchDisChild.* 1996;75(3):214-7.
2. Bhutta ZA. Enteric Fever: Risk Factors for Treatment Failure in the Era of Multidrug Resistance. UNPUBLISHED.
3. Buch NA, Hassan MU, Kakroo DK. Enteric fever--a changing sensitivity pattern, clinical profile and outcome. *Indian Pediatr.* 1994;31(8):981-5.
4. Chandra R, Srinivasan S, Nalini P, Rao RS. Multidrug resistant enteric fever. *J TropMed Hyg.* 1992;95(4):284-7.
5. Dutta P, Lahiri M, Bhattacharya SK, Saha MR, Mitra U, Rasaily R, et al. Hepatitis-like presentation in typhoid fever. *TransRSocTropMed Hyg.* 1992;86(1):92.
6. Grgis NI, Tribble DR, Sultan Y, Farid Z. Short course chemotherapy with cefixime in children with multidrug-resistant *Salmonella typhi* Septicaemia. *J TropPediatr.* 1995;41(6):364-5.
7. Gulati S, Marwaha RK, Singhi S, Ayyagari A, Kumar L. Third generation cephalosporins in multi-drug resistant typhoid fever. *Indian Pediatr.* 1992;29(4):513-6.
8. Jagadish K, Patwari AK, Sarin SK, Prakash C, Srivastava DK, Anand VK. Hepatic manifestations in typhoid fever. *Indian Pediatr.* 1994;31(7):807-11.
9. Kabra SK, Madhulika, Talati A, Soni N, Patel S, Modi RR. Multidrug-resistant typhoid fever. *TropDoct.* 2000;30(4):195-7.
10. Khan MI, Soofi SB, Ochiai RL, Khan MJ, Sahito SM, Habib MA, et al. Epidemiology, clinical presentation, and patterns of drug resistance of *Salmonella Typhi* in Karachi, Pakistan. *J Infect Dev Ctries.* 2012 Oct;6(10):704-14.
11. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramchandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: clinical profile and therapy. *Indian Pediatr.* 1991;28(4):357-61.
12. Kumar R, Gupta N. Multidrug-resistant typhoid fever. *Indian J Pediatr.* 2007;74(1):39-42.
13. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr.* 1991;28(10):1171-4.
14. Raman TS, Krishnamurthy L, Menon PK, Singh D, Jayaprakash DG. Clinical profile and therapy in enteric fever. *Indian Pediatr.* 1994;31(2):196-9.
15. Rasaily R, Dutta P, Saha MR, Mitra U, Lahiri M, Pal SC. Multi-drug resistant typhoid fever in hospitalised children. Clinical, bacteriological and epidemiological profiles. *EurJ Epidemiol.* 1994;10(1):41-6.
16. Sharma A, Gathwala G. Clinical profile and outcome in enteric fever. *Indian Pediatr.* 1993;30(1):47-50.

Table S11 :Time-line of included studies on enteric fever in children

	1950-1970s 13 studies			1980-1989 21 studies			1990 – 1999 30 studies			2000-2013 12 studies		
Demographics/History	N	N	%	n	N	%	n	N	%	n	N	%
Blood/bone marrow												
Isolates	882	1113	79	1283	1629	79	4562	4941	92	918	918	100
Pre-treatment antibiotics received	23	34	68	137	295	46	1875	3283	57	45	227	20
Duration of illness≤ 1 week	204	404	50	516	841	61	711	1570	45	82	251	33
Multi-drug resistant isolates	0	127	0	1	351	0	1609	3772	43	281	598	47
Fluoroquinolone resistance							12	1148	1	8	167	5
Chloramphenicol resistance	0	127	0	49	497	10	591	1042	57	0	72	0
Signs & Symptoms: Systemic												
Fever	662	711	93	1327	1399	95	3694	4014	92	731	732	100
High grade fever	8	70	11	199	385	52	1847	2706	68	247	465	53
Chills or rigors	22	137	16	39	488	8	185	566	33	48	190	25
Bradycardia or relative bradycardia	30	390	8	355	771	46	24	149	16	9	270	3
Anorexia/refusal to feed	203	463	44	269	523	51	706	3046	23	92	319	29
Headache	147	538	27	456	1235	37	295	3256	9	314	766	41
Toxicity	171	295	58	79	288	27	978	3310	30	48	76	63
Coated tongue	81	150	54	157	196	80	146	294	50	65	198	33
Respiratory and abdominal												
Cough	244	641	38	290	878	33	464	2377	20	178	666	27
Nausea or vomiting	290	655	44	354	935	38	1361	3757	36	282	783	36
Diarrhea	280	701	40	718	1561	46	1312	3964	33	182	802	23
Constipation	87	348	25	243	811	30	251	3445	7	35	583	6
Hepatomegaly	139	471	30	372	747	50	1922	4123	47	359	603	60
Splenomegaly	343	869	39	266	905	29	1280	4358	29	222	603	37
Abdominal pain or tenderness	273	559	49	574	1476	39	764	3194	24	383	763	50
Abdominal distention or ileus	235	611	38	292	1155	25	171	3657	5	43	515	8
Laboratory features												
Anaemia (Hb < 12 g/dl)	233	467	50	323	967	33	2124	2959	72	189	337	56
Leucopenia (< 5 x 10 ³ /µL)	92	379	24	272	906	30	109	1589	7	26	338	8
Leucocytosis (> 15 x 10 ³ /µL)	16	161	10	25	302	8	330	1465	23	54	275	20
Thrombocytopenia (<150 x 10 ³ /µL)	41	161	25	104	415	25	79	1188	7	39	198	20
Complications & outcome												
Shock or hypotension	8	188	4	3	81	4	89	2344	4	24	408	6
Altered mental status	146	650	22	276	1495	18	290	4176	7	75	517	15
Pneumonia or chest signs	117	610	19	276	1204	23	148	1580	9	37	394	9
GI bleeding	40	784	5	23	1351	2	48	716	7	39	586	7
GI perforation	58	660	9	12	1047	1	7	775	1	3	130	2
Hepatitis or raised LFTs	18	277	6	66	145	46	172	2072	8	43	266	16

Jaundice	21	484	4	22	534	4	81	3381	2	38	408	9
Relapse	59	658	9	29	1053	3	173	3235	5	5	248	2
Death	73	986	7	40	1518	3	62	4288	1	5	722	1

References to Table S6 : Time-line of included studies on enteric fever in children

1950 - 1970

1. Collins RN, Marine WM, Nahmias AJ. The 1964 epidemic of typhoid fever in Atlanta. Clinical and epidemiologic observations. *JAMA*. 1966;197(3):179-84.
2. Colon AR. Gastrointestinal phenomena in childhood typhoid fever. *SouthMed J*. 1976;69(7):914.
3. Colon AR, Gross DR, Tamer MA. Typhoid fever in children. *Pediatrics*. 1975;56(4):606-9.
4. Del Rosario AJ, Haynes RE, Cramblett HG. Typhoid fever in children. *OhioState Med J*. 1971;67(6):516-20.
5. Duggan MB, Beyer L. Enteric fever in young Yoruba children. *ArchDisChild*. 1975;50(1):67-71.
6. Galloway H, Clark NS, Blackhall M. Paediatric aspects of the Aberdeen typhoid outbreak. *ArchDisChild*. 1966;41(215):63-8.
7. Hardy G, Padfield CJ, Chadwick P, Partington MW. Typhoid outbreak in Kingston, Ont: experience with high-dose oral ampicillin. *CanMed AssocJ*. 1977;116(7):761-4, 7.
8. Johnson AO, Aderele WI. Enteric fever in childhood. *J TropMed Hyg*. 1981;84(1):29-35.
9. Laditan AA, Alausa KO. Problems in the clinical diagnosis of typhoid fever in children in the tropics. *AnnTropPaediatr*. 1981;1(3):191-5.
10. Khuri-Bulos N. Enteric fevers in children. The importance of age in the varying clinical picture. *ClinPediatr (Phila)*. 1981;20(7):448-52.
11. Kagwa-Nyanzi JA. Typhoid fever in African children. *J TropPediatr EnvironChild Health*. 1971;17(3):105-9.
12. Moussa K. Enterica of childhood. *J EgyptPublic Health Assoc*. 1970;45(1):48-55.
13. Mulligan TO. Typhoid fever in young children. *BrMed J*. 1971;4(5788):665-7.
14. Scragg JN, Rubidge CJ. Trimethoprim and sulphamethoxazole in typhoid fever in children. *BrMed J*. 1971;3(5777):738-41.
15. Tolaymat A, Fakhreddine F, David CB, Whitworth JM. Typhoid fever in children: a forgotten disease? *SouthMed J*. 1979;72(2):136-8.
16. Verma PS, Mullick P, Ghosh S. Neurological manifestations of enteric fever: clinical profile and correlation with ultimate outcome. *Indian Pediatr*. 1972;9(11):681-5.

1980 - 1989

1. Brian MJ. Typhoid in children in Goroka. *PNGMed J*. 1990;33(1):68-9.
2. Caglar MK, Ozsoylu S, Kanra G. Relative granulocytosis in childhood typhoid fever. *J Pediatr*. 1983;102(4):603-4.
3. Chiu CH, Lin TY. Typhoid fever in children. *Lancet*. 1999;354(9194):2001-2.

4. Chiu CH, Tsai JR, Ou JT, Lin TY. Typhoid fever in children: a fourteen-year experience. *Acta Paediatr Taiwan*. 2000;41(1):28-32.
5. Choo KE, Razif A, Ariffin WA, Sepiah M, Gururaj A. Typhoid fever in hospitalized children in Kelantan, Malaysia. *AnnTropPaediatr*. 1988;8(4):207-12.
6. Chow CB, Wang PS, Leung NK. Typhoid fever in Hong Kong children. *AustPaediatrJ*. 1989;25(3):147-50.
7. Daoud AS, Zaki M, Pugh RN, al-Mutairi G, Beseiso R, Nasrallah AY. Clinical presentation of enteric fever: its changing pattern in Kuwait. *J TropMed Hyg*. 1991;94(5):341-7.
8. Ellis ME, Moosa A, Hillier V. A review of typhoid fever in South African black children. *PostgradMed J*. 1990;66(782):1032-6.
9. Ferreccio C, Levine MM, Manterola A, Rodriguez G, Rivara I, Prenzel I, et al. Benign bacteremia caused by *Salmonella typhi* and *paratyphi* in children younger than 2 years. *J Pediatr*. 1984;104(6):899-901.
10. Green SD, Cheesbrough JS. *Salmonella* bacteraemia among young children at a rural hospital in western Zaire. *AnnTropPaediatr*. 1993;13(1):45-53.
11. Ispahani P, Slack RC. Enteric fever and other extraintestinal salmonellosis in University Hospital, Nottingham, UK, between 1980 and 1997. *EurJ ClinMicrobiolInfectDis*. 2000;19(9):679-87.
12. Jain S, Chitnis DS, Sham A, Rathi S, Inamdar S, Rindani GJ. Outbreak of chloramphenicol resistant typhoid fever. *Indian Pediatr*. 1987;24(3):193-7.
13. Kanra G, Yurdakok M, Secmeer G, Arvas A, Dereli O, Sarikayalar F, et al. Clinical findings of pediatric patients during an outbreak of typhoid fever in Ankara. *TurkJ Pediatr*. 1985;27(1):11-5.
14. Kapoor JP, Mohan M, Talwar V, Daral TS, Bhargava SK. Typhoid fever in young children. *Indian Pediatr*. 1985;22(11):811-3.
15. Lepage P, Bogaerts J, Van GC, Ntahorutaba M, Nsengumuremyi F, Hitimana DG, et al. Community-acquired bacteraemia in African children. *Lancet*. 1987;1(8548):1458-61.
16. Mallouh AA, Sa'di AR. White blood cells and bone marrow in typhoid fever. *Pediatr InfectDisJ*. 1987;6(6):527-9
17. Nesbitt A, Mirza NB. *Salmonella* septicaemias in Kenyan children. *J TropPediatr*. 1989;35(1):35-9.
18. Secmeer G, Kanra G, Cemeroglu AP, Ozen H, Ceyhan M, Ecevit Z. *Salmonella typhi* infections. A 10-year retrospective study. *TurkJ Pediatr*. 1995;37(4):339-41.
19. Soelistiyowati S, Soenarto Y, Soesilo H, Widiarto, Widiatmodjo, Ismangoen. Typhoid fever in children. *PaediatrIndones*. 1982;22(7-8):138-46.
20. Stormon MO, McIntyre PB, Morris J, Fasher B. Typhoid fever in children: diagnostic and therapeutic difficulties. *Pediatr InfectDisJ*. 1997;16(7):713-4.
21. Thisyakorn U, Mansuwan P, Taylor DN. Typhoid and paratyphoid fever in 192 hospitalized children in Thailand. *AmJ DisChild*. 1987;141(8):862-5.
22. Yap YF, Puthucheary SD. Typhoid fever in children--a retrospective study of 54 cases from Malaysia. *Singapore Med J*. 1998;39(6):260-2.

1990 - 1994

1. Ahmad KA, Khan LH, Roshan B, Bhutta ZA. Factors associated with typhoid relapse in the era of multiple drug resistant strains. *J Infect Dev Ctries*. 2011 Oct;5(10):727-31.

2. Arora RK, Gupta A, Joshi NM, Kataria VK, Lall P, Anand AC. Multidrug resistant typhoid fever: study of an outbreak in Calcutta. *Indian Pediatr.* 1992;29(1):61-6.
3. Bhutta ZA. Impact of age and drug resistance on mortality in typhoid fever. *ArchDisChild.* 1996;75(3):214-7.
4. Bhutta ZA. Fulminant hepatic failure with typhoid fever in childhood. *J Pak Med Assoc.* 1991;41(6):123-6.
5. Bhutta ZA, Mansoorali N, Hussain R. Plasma cytokines in paediatric typhoidal salmonellosis: correlation with clinical course and outcome. *J Infect.* 1997;35(3):253-6.
6. Bhutta ZA, Naqvi SH. Hepatic dysfunction in paediatric typhoidal salmonellosis. *AnnTropPaediatr.* 1991;11(3):203-5.
7. Bhutta ZA, Naqvi SH, Durrani S, Suria A. Chloramphenicol therapy of typhoid fever and its relationship to hepatic dysfunction. *JTropPediatr.* 1991;37(6):320-2.
8. Bhutta ZA, Naqvi SH, Razzaq RA, Farooqui BJ. Multidrug-resistant typhoid in children: presentation and clinical features. *RevInfectDis.* 1991;13(5):832-6.
9. Bhutta ZA, Niazi SK, Suria A. Chloramphenicol clearance in typhoid fever: implications for therapy. *Indian J Pediatr.* 1992;59(2):213-9
10. Bhutta ZA. Enteric Fever: Risk Factors for Treatment Failure in the Era of Multidrug Resistance. UNPUBLISHED.
11. Biswal N. Neurological manifestations of typhoid fever in children. *J TropPediatr.* 1994;40(3):190.
12. Biswal N, Mathai B, Bhatia BD, Srinivasan S, Nalini P. Enteric fever: a changing perspective. *Indian Pediatr.* 1994;31(7):813-9.
13. Buch NA, Hassan MU, Kakroo DK. Enteric fever--a changing sensitivity pattern, clinical profile and outcome. *Indian Pediatr.* 1994;31(8):981-5.
14. Chandra R, Srinivasan S, Nalini P, Rao RS. Multidrug resistant enteric fever. *J TropMed Hyg.* 1992;95(4):284-7.
15. Dutta P, Lahiri M, Bhattacharya SK, Saha MR, Mitra U, Rasaily R, et al. Hepatitis-like presentation in typhoid fever. *TransRSocTropMed Hyg.* 1992;86(1):92.
16. Dutta P, Rasaily R, Saha MR, Mitra U, Manna B, Chakraborty S, et al. Randomized clinical trial of furazolidone for typhoid fever in children. *ScandJGastroenterol.* 1993;28(2):
17. Garg RA, Krashak R. Typhoid fever before two years of age. *Indian Pediatr.* 1993;30(6):805-8.
18. Girgis NI, Tribble DR, Sultan Y, Farid Z. Short course chemotherapy with cefixime in children with multidrug-resistant *Salmonella typhi* Septicaemia. *J TropPediatr.* 1995;41(6):364-5.
19. Hassan M. Multiply-resistant *Salmonella typhi* in children. *Drugs.* 1995;49 Suppl 2:457-9.
20. Hoa NT, Diep TS, Wain J, Parry CM, Hien TT, Smith MD, et al. Community-acquired septicaemia in southern Viet Nam: the importance of multidrug-resistant *Salmonella typhi*. *TransRSocTropMedHyg.* 1998;92(5):503-8.
21. Khan M, Coovadia YM, Connolly C, Sturm AW. Influence of sex on clinical features, laboratory findings, and complications of typhoid fever. *AmJ TropMed Hyg.* 1999;61(1):41-6.
22. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramchandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: clinical profile and therapy. *Indian Pediatr.* 1991;28(4):357-61.

23. Memon IA, Billoo AG, Memon HI. Cefixime: an oral option for the treatment of multidrug-resistant enteric fever in children. *SouthMed J.* 1997;90(12):1204-7.
24. Misra S, Diaz PS, Rowley AH. Characteristics of typhoid fever in children and adolescents in a major metropolitan area in the United States. *ClinInfectDis.* 1997;24(5):998-1000.
25. Mishra S, Patwari AK, Anand VK, Pillai PK, Aneja S, Chandra J, et al. A clinical profile of multidrug resistant typhoid fever. *Indian Pediatr.* 1991;28(10):1171-4.
26. Oh HM, Masayu Z, Chew SK. Typhoid fever in hospitalized children in Singapore. *J Infect.* 1997;34(3):237-42.
27. Rajajee S, Anandi TB, Subha S, Vatsala BR. Patterns of resistant *Salmonella typhi* infection in infants. *JTropPediatr.* 1995;41(1):52-4.
28. Ramanan A, Pandit N, Yeshwanth M. Unusual complications in a multidrug resistant *Salmonella typhi* outbreak. *Indian Pediatr.* 1992;29(1):118-20.
29. Rasaily R, Dutta P, Saha MR, Mitra U, Lahiri M, Pal SC. Multi-drug resistant typhoid fever in hospitalised children. Clinical, bacteriological and epidemiological profiles. *EurJ Epidemiol.* 1994;10(1):41-6
30. Rathore MH, Bux D, Hasan M. Multidrug-resistant *Salmonella typhi* in Pakistani children: clinical features and treatment. *SouthMed J.* 1996;89(2):235-7.
31. Shetty AK, Mital SR, Bahrainwala AH, Khubchandani RP, Kumta NB. Typhoid hepatitis in children. *J TropPediatr.* 1999;45(5):287-90
32. Sethuraman S, Mahamood M, Kareem S. Furazolidone in multi-resistant childhood typhoid fever. *AnnTropPaediatr.* 1994;14(4):321-4.
33. Thambidorai CR, Shyamala J, Sarala R, Vatsala RB, Tamizhisai S. Acute acalculous cholecystitis associated with enteric fever in children. *Pediatr InfectDisJ.* 1995;14(9):812-3.

1995 - 1999

1. Ciftci E, Guriz H, Derya AA, Ince E, Erdem B, Dogru U. *Salmonella* bacteraemia in Turkish children: 37 cases seen in a university hospital between 1993 and 2002. *AnnTropPaediatr.* 2004;24(1):75-80.
2. Malik AS. Complications of bacteriologically confirmed typhoid fever in children. *J TropPediatr.* 2002;48(2):102-8.
3. Malik AS, Malik RH. Typhoid fever in Malaysian children. *Med J Malaysia.* 2001;56(4):478-90.
4. Papaevangelou V, Syriopoulou V, Charissiadou A, Pangalis A, Mostrou G, Theodoridou M. *Salmonella* bacteraemia in a tertiary children's hospital. *ScandJ InfectDis.* 2004;36(8):547-51.
5. Shwe TN, Nyein MM, Yi W, Mon A. Blood culture isolates from children admitted to Medical Unit III, Yangon Children's Hospital, 1998. *Southeast Asian J TropMed Public Health.* 2002;33(4):764-71.
6. Sinha A, Sazawal S, Kumar R, Sood S, Reddaiah VP, Singh B, et al. Typhoid fever in children aged less than 5 years. *Lancet.* 1999;354(9180):734-7.
7. Su CP, Chen YC, Chang SC. Changing characteristics of typhoid fever in Taiwan. *J MicrobiolImmunolInfect.* 2004;37(2):109-14.
8. Wongsawat J, Pancharoen C, Thisyakorn U. Typhoid fever in children: experience in King Chulalongkorn Memorial Hospital. *J Med AssocThai.* 2002;85(12):1247-50.

2000 onwards

1. Dheer G, Kundra S, Singh T. Clinical and laboratory profile of enteric fever in children in northern India. *Trop Doct.* 2012 Jul;42(3):154-6.
2. Emary K, Moore CE, Chanpheaktra N, An KP, Chheng K, Sona S, et al. Enteric fever in Cambodian children is dominated by multidrug-resistant H58 *Salmonella enterica* serovar Typhi with intermediate susceptibility to ciprofloxacin. *Trans R Soc Trop Med Hyg.* 2012 Dec;106(12):718-24.
3. Khan MI, Soofi SB, Ochiai RL, Khan MJ, Sahito SM, Habib MA, et al. Epidemiology, clinical presentation, and patterns of drug resistance of *Salmonella* Typhi in Karachi, Pakistan. *J Infect Dev Ctries.* 2012 Oct;6(10):704-14.
4. Kumar R, Gupta N. Multidrug-resistant typhoid fever. *Indian J Pediatr.* 2007;74(1):39-42.
5. Leung DT, Bogetz J, Itoh M, Ganapathi L, Pietroni MA, Ryan ET, et al. Factors associated with encephalopathy in patients with *Salmonella enterica* serotype Typhi bacteremia presenting to a diarrheal hospital in Dhaka, Bangladesh. *Am J Trop Med Hyg.* 2012 Apr;86(4):698-702.
6. Mtove G, Amos B, von Seidlein L, Hendriksen I, Mwambuli A, Kimera J, et al. Invasive salmonellosis among children admitted to a rural Tanzanian hospital and a comparison with previous studies. *PLoS One.* 2010;5(2):e9244.
7. Owais A, Sultana S, Zaman U, Rizvi A, Zaidi AK. Incidence of typhoid bacteremia in infants and young children in southern coastal Pakistan. *Pediatr Infect Dis J.* 2010 Nov;29(11):1035-9.
8. Pradhan R, Shrestha U, Gautam SC, Thorson S, Shrestha K, Yadav BK, et al. Bloodstream infection among children presenting to a general hospital outpatient clinic in urban Nepal. *PLoS One.* 2012;7(10):e47531
9. Siddiqui FJ, Rabbani F, Hasan R, Nizami SQ, Bhutta ZA. Typhoid fever in children: some epidemiological considerations from Karachi, Pakistan. *Int J Infect Dis.* 2006;10(3):215-22.
10. Tatli MM, Aktas G, Kosecik M, Yilmaz A. Treatment of typhoid fever in children with a flexible-duration of ceftriaxone, compared with 14-day treatment with chloramphenicol. *Int J Antimicrob Agents.* 2003;21(4):350-3.
11. Walia M, Gaind R, Mehta R, Paul P, Aggarwal P, Kalaivani M. Current perspectives of enteric fever: a hospital-based study from India. *Ann Trop Paediatr.* 2005;25(3):161-74.
12. Walia M, Gaind R, Paul P, Mehta R, Aggarwal P, Kalaivani M. Age-related clinical and microbiological characteristics of enteric fever in India. *Trans R Soc Trop Med Hyg.* 2006;100(10):942-8.
13. Wang JL, Kao JH, Tseng SP, Teng LJ, Ho SW, Hsueh PR. Typhoid fever and typhoid hepatitis in Taiwan. *Epidemiol Infect.* 2005;133(6):1073-9.

Chart 1 : Definitions of variables used

Variables	As defined by author, or :
Multi-drug resistance	Resistance to all 3 of chloramphenicol, ampicillin/amoxycillin, cotrimoxazole
Weakness or malaise	fatigue, weakness, dizziness, malaise, prostration
High grade fever	> 104 F or > 40 C, hyperpyrexia, extreme hyperpyrexia
Toxicity	toxemia or ‘enteric state’
Chelosis	stomatitis, mouth ulcers, aphous ulcers, parotitis, stomatitis, cracked lips
Pneumonia, bronchopneumonia or chest signs	abnormal CXR, congestion, consolidation, rales, rhonchi, crepts, dyspnea, respiratory/chest/lung/bronchitic signs, pulmonary infiltration, bronchitis, respiratory distress, bronchial breathing, broncho-pulmonary involvement
Complications	
Altered mental status	mental change, encephalopathy, encephalitis, drowsy, lethargy, confusion, coma, semi-coma, delirium, psychosis, irritability, disoriented, stupor, obtundation, disordered thought, CNS disturbances, neuro-psychiatric features/manifestations, altered sensorium, clouded/altered consciousness, acute schizophrenia, global cerebral dysfunction, somnolence, apathy, aphasia, ataxia, deafness, motor paresis, hemiplegia, tremor, difficulty speaking, nystagmus, dysarthria, slurred speech, chorea/choreoform movements, cerebellar ataxia, cerebellitis, cerebellar disease, Parkinsonism, facial palsy, unsteady gait, unsteady/falls, myelitis, transverse myelitis, CN palsy
Focal neurological signs or symptoms	elevated SGPT/ALT, hepatic dysfunction, raised LFTs or transaminases (if not disaggregated, both ALT and/or AST elevated extracted), hepato-biliary involvement (taken ALT raised as > 35 if not specified by author), or ALT and/or bilirubin elevation, liver failure
Hepatitis or raised LFTs	tympanities, tympanic abdomen, meteorism, tumid abdomen, decreased or absent bowel sounds, signs of intestinal obstruction
Abdominal distention or ileus	nephritis, nephrotic, proteinuria, albuminuria, raised BUN or Cr, abnormal urine d/r, hematuria, pyuria, renal failure, oliguria, nephropathy
Renal involvement	limb/extremity pain, leg pain, backache, bone pain, joint pain, joint swelling, reactive arthritis, bodyache, muscle pain
Outcome	
Relapse	Clinical, or clinical and culture-confirmed
Laboratory features	
Anaemia(<12g/dl)	undefined, moderate anaemia, Hb 5-10, 8.5-10Hct< 30%/32%, Hb<9, < 10, <=10.5, <= 11, < 12, <=12
Severe anaemia (<7.5 g/dl)	undefined, <5, < 6, <6.5, < 6.7, < 7.4
Leucopenia (<5 x 10 ³ /µL)	<4, <4.5, <3, or undefined
Leucocytosis (>15 x 10 ³ /µL)	or undefined
Thrombocytopenia (<150 x 10 ³ /µL)	<100, or undefined
Severe thrombocytopenia (<50 x 10 ³ /µL)	or undefined
Neutrophilia (>7.5 x 10 ³ /µL)	or undefined