

**Supplemental Table 1.**

	<b>Included (n=253)</b>	<b>Excluded (n=518)</b>	<b>p-value</b>
<b>Age, yrs</b>	56.4(14.0)	57.5(14.4)	0.32
<b>Male sex, n(%)</b>	140(55)	315(61)	0.15
<b>African American, n(%)<sup>±</sup></b>	203(81)	269(52)	<0.001
<b>Diabetes mellitus, n(%)</b>	108(43)	249(48)	0.16
<b>Peripheral Vascular Disease, n(%)</b>	22(9)	52(10)	0.55
<b>ESRD vintage (yrs)</b>	3.4(1.7,7.9)	3.2(1.4,6.6)	0.005
<b>Albumin (g/dl)</b>	4.0(0.36)	4.0(0.36)	0.41
<b>Hemoglobin (g/dl)</b>	11.9(3.7)	11.7(1.9)	0.33
<b>Arterio-venous Fistula, n(%)<sup>χ</sup></b>	134(53)	342(67)	<0.001
<b>ICW<sup>†*</sup> (L)</b>	21.8(5.1)	21.9(5.7)	0.71
<b>ECW<sup>°*</sup> (L)</b>	19.2(4.4)	19.2(4.9)	0.99
<b>Fat mass* (kg)</b>	24.8(13.6)	24.9(12.8)	0.94

<sup>±</sup>Includes 1 missing value

Values are mean (sd) for normally distributed variables and median (25th, 75th percentile) for non-normally distributed continuous variables.

<sup>χ</sup>Includes 6 missing values

\*Data for 82% of study population with bioimpedance spectroscopy performed

<sup>†</sup>Intracellular water

<sup>°</sup>Extracellular water