

S1 File. Supplementary tables

Table A. Three models on gait speed in later life (fully adjusted).

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	0.716 ± 0.264 [†]	0.803 ± 0.264 [†]	0.745 ± 0.264 [†]
Sex, Female	-0.048 ± 0.008 [†]	-0.049 ± 0.008 [†]	-0.050 ± 0.008 [†]
Age	0.006 ± 0.007	0.006 ± 0.007	0.006 ± 0.007
Age ²	0.000 ± 0.000*	0.000 ± 0.000*	0.000 ± 0.000*
Household size	0.007 ± 0.006	0.007 ± 0.006	0.007 ± 0.006
Routine manual	-0.041 ± 0.009 [†]	-0.039 ± 0.009 [†]	-0.038 ± 0.009 [†]
Wealth mid-tertile	0.043 ± 0.01 [†]	0.041 ± 0.010 [†]	0.041 ± 0.010 [†]
Wealth top-tertile	0.088 ± 0.011 [†]	0.086 ± 0.011 [†]	0.086 ± 0.011 [†]
Age left education	0.006 ± 0.002 [†]	0.005 ± 0.002 [†]	0.005 ± 0.002 [†]
Married-cohab.	0.038 ± 0.020	0.038 ± 0.020	0.038 ± 0.020
Separated-div.	0.038 ± 0.023	0.037 ± 0.023	0.038 ± 0.023
Widow	0.034 ± 0.021	0.034 ± 0.021	0.034 ± 0.021
Social support	-0.018 ± 0.037	-0.018 ± 0.037	-0.019 ± 0.037
Social connection	0.005 ± 0.001 [†]	0.005 ± 0.001 [†]	0.005 ± 0.001 [†]
CVD	-0.024 ± 0.010*	-0.024 ± 0.010*	-0.024 ± 0.010*
COPD	-0.039 ± 0.011 [†]	-0.039 ± 0.011 [†]	-0.039 ± 0.011 [†]
Cancer	-0.006 ± 0.015	-0.007 ± 0.015	-0.007 ± 0.015
Smoking	-0.027 ± 0.013*	-0.026 ± 0.013*	-0.026 ± 0.013*
Drinking	0.020 ± 0.009*	0.020 ± 0.009*	0.019 ± 0.009*
Exercise, vigorous	0.046 ± 0.011 [†]	0.046 ± 0.011 [†]	0.046 ± 0.011 [†]
Exercise, moderate	0.107 ± 0.008 [†]	0.107 ± 0.008 [†]	0.107 ± 0.008 [†]
Childhood lack facility	-0.005 ± 0.003	-0.031 ± 0.003 [†]	-0.035 ± 0.011 [†]
<i>RMSE</i>	0.0038	0.0025	0.0037

Source: The English Longitudinal Study of Ageing.

* $p < 0.05$.

[†] $p < 0.01$.

Table B. Three models on episodic memory in later life (fully adjusted).

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	11.722 ± 2.118 [†]	12.140 ± 2.118 [†]	11.894 ± 2.118 [†]
Sex, Female	0.772 ± 0.088 [†]	0.768 ± 0.088 [†]	0.765 ± 0.088 [†]
Age	-0.047 ± 0.058	-0.048 ± 0.058	-0.050 ± 0.058
Age ²	-0.001 ± 0.000	-0.001 ± 0.000	-0.001 ± 0.000
Household size	0.037 ± 0.052	0.036 ± 0.052	0.036 ± 0.052
Routine manual	-0.632 ± 0.095 [†]	-0.622 ± 0.095 [†]	-0.618 ± 0.095 [†]
Wealth mid-tertile	0.308 ± 0.109 [†]	0.303 ± 0.109 [†]	0.299 ± 0.109 [†]
Wealth top-tertile	0.596 ± 0.116 [†]	0.585 ± 0.116 [†]	0.579 ± 0.116 [†]
Age left education	0.185 ± 0.019 [†]	0.180 ± 0.019 [†]	0.177 ± 0.019 [†]
Married-cohab.	0.105 ± 0.203	0.106 ± 0.203	0.105 ± 0.203
Separated-div.	0.137 ± 0.230	0.135 ± 0.229	0.130 ± 0.229
Widow	0.156 ± 0.219	0.156 ± 0.219	0.156 ± 0.219
Social support	0.173 ± 0.399	0.176 ± 0.399	0.175 ± 0.399
Social connection	0.064 ± 0.012 [†]	0.063 ± 0.012 [†]	0.063 ± 0.012 [†]
CVD	-0.083 ± 0.108	-0.084 ± 0.108	-0.085 ± 0.108
COPD	0.170 ± 0.116	0.172 ± 0.116	0.174 ± 0.116
Cancer	0.333 ± 0.177	0.328 ± 0.177	0.323 ± 0.177
Smoking	-0.351 ± 0.129 [†]	-0.346 ± 0.129 [†]	-0.342 ± 0.129 [†]
Drinking	0.477 ± 0.091 [†]	0.474 ± 0.091 [†]	0.474 ± 0.091 [†]
Exercise, vigorous	0.085 ± 0.109	0.086 ± 0.109	0.087 ± 0.109
Exercise, moderate	0.384 ± 0.091 [†]	0.384 ± 0.091 [†]	0.383 ± 0.091 [†]
CESD	-0.116 ± 0.024 [†]	-0.116 ± 0.024 [†]	-0.116 ± 0.024 [†]
Childhood lack facility	-0.054 ± 0.029	-0.191 ± 0.029 [†]	-0.273 ± 0.118*
<i>RMSE</i>	0.0406	0.0092	0.0405

Source: The English Longitudinal Study of Ageing.

* $p < 0.05$.

† $p < 0.01$.

Table C. Three models on mental health in later life (fully adjusted).

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	2.185 ± 1.229	2.450 ± 1.229*	2.183 ± 1.230
Sex, Female	-0.318 ± 0.051 [†]	-0.320 ± 0.051 [†]	-0.317 ± 0.051 [†]
Age	0.063 ± 0.034	0.062 ± 0.034	0.063 ± 0.034
Age ²	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Household size	0.060 ± 0.030*	0.060 ± 0.030	0.060 ± 0.030*
Routine manual	-0.097 ± 0.055	-0.091 ± 0.055	-0.097 ± 0.055
Wealth mid-tertile	0.287 ± 0.063 [†]	0.283 ± 0.063 [†]	0.287 ± 0.063 [†]
Wealth top-tertile	0.517 ± 0.067 [†]	0.510 ± 0.067 [†]	0.517 ± 0.067 [†]
Age left education	0.009 ± 0.011	0.006 ± 0.011	0.009 ± 0.011
Married-cohab.	0.115 ± 0.118	0.116 ± 0.118	0.115 ± 0.118
Separated-div.	-0.211 ± 0.133	-0.213 ± 0.133	-0.212 ± 0.133
Widow	-0.502 ± 0.127 [†]	-0.502 ± 0.127 [†]	-0.502 ± 0.127 [†]
Social support	1.107 ± 0.232 [†]	1.109 ± 0.232 [†]	1.108 ± 0.232 [†]
Social connection	0.030 ± 0.007 [†]	0.030 ± 0.007 [†]	0.030 ± 0.007 [†]
CVD	-0.104 ± 0.063	-0.105 ± 0.063	-0.104 ± 0.063
COPD	-0.474 ± 0.067 [†]	-0.473 ± 0.067 [†]	-0.474 ± 0.067 [†]
Cancer	-0.115 ± 0.103	-0.118 ± 0.103	-0.115 ± 0.103
Smoking	-0.255 ± 0.075 [†]	-0.252 ± 0.075 [†]	-0.255 ± 0.075 [†]
Drinking	0.132 ± 0.053*	0.130 ± 0.053*	0.132 ± 0.053*
Exercise, vigorous	0.134 ± 0.064*	0.134 ± 0.064*	0.134 ± 0.064*
Exercise, moderate	0.504 ± 0.053 [†]	0.504 ± 0.053 [†]	0.504 ± 0.053 [†]
Childhood lack facility	-0.005 ± 0.017	-0.091 ± 0.017 [†]	-0.003 ± 0.017
<i>RMSE</i>	0.0236	0.0206	0.0236

Source: The English Longitudinal Study of Ageing.

* $p < 0.05$.

[†] $p < 0.01$.

Table D. Three models on gait speed in later life, with adult health (fully adjusted).

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	0.667 ± 0.262*	0.746 ± 0.262 [†]	0.697 ± 0.262 [†]
Sex, Female	-0.049 ± 0.008 [†]	-0.050 ± 0.008 [†]	-0.050 ± 0.008 [†]
Age	0.009 ± 0.007	0.009 ± 0.007	0.008 ± 0.007
Age ²	0.000 ± 0.000 [†]	0.000 ± 0.000 [†]	0.000 ± 0.000 [†]
Household size	0.004 ± 0.006	0.004 ± 0.006	0.004 ± 0.006
Routine manual	-0.039 ± 0.009 [†]	-0.037 ± 0.009 [†]	-0.037 ± 0.009 [†]
Wealth mid-tertile	0.039 ± 0.010 [†]	0.037 ± 0.010 [†]	0.037 ± 0.010 [†]
Wealth top-tertile	0.081 ± 0.011 [†]	0.080 ± 0.011 [†]	0.079 ± 0.011 [†]
Age left edu	0.006 ± 0.002 [†]	0.005 ± 0.002 [†]	0.005 ± 0.002 [†]
Married-cohab.	0.040 ± 0.020*	0.040 ± 0.020*	0.040 ± 0.020*
Separated-div.	0.040 ± 0.023	0.040 ± 0.023	0.040 ± 0.023
Widow	0.033 ± 0.021	0.033 ± 0.021	0.033 ± 0.021
Social support	-0.023 ± 0.036	-0.023 ± 0.036	-0.023 ± 0.036
Social connection	0.005 ± 0.001 [†]	0.005 ± 0.001 [†]	0.005 ± 0.001 [†]
CVD	-0.021 ± 0.010*	-0.021 ± 0.010*	-0.021 ± 0.010*
COPD	-0.028 ± 0.011 [†]	-0.027 ± 0.011*	-0.027 ± 0.011*
Cancer	0.004 ± 0.015	0.003 ± 0.015	0.003 ± 0.015
Smoking	-0.027 ± 0.013*	-0.027 ± 0.013*	-0.026 ± 0.013*
Drinking	0.018 ± 0.009*	0.017 ± 0.009*	0.017 ± 0.009*
Exercise, vigorous	0.041 ± 0.011 [†]	0.041 ± 0.011 [†]	0.041 ± 0.011 [†]
Exercise, moderate	0.101 ± 0.008 [†]	0.101 ± 0.008 [†]	0.101 ± 0.008 [†]
Childhood lack facility	-0.005 ± 0.003	-0.029 ± 0.003 [†]	-0.035 ± 0.011 [†]
Adult illness episodes	-0.032 ± 0.004 [†]	-0.032 ± 0.004 [†]	-0.032 ± 0.004 [†]
<i>RMSE</i>	0.0038	0.0006	0.0037

Source: The English Longitudinal Study of Ageing 2006.

* $p < 0.05$.

[†] $p < 0.01$.

Table E. Three models on episodic memory in later life, with adult health (fully adjusted).

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	11.750 ± 2.120 [†]	12.174 ± 2.120 [†]	11.927 ± 2.120 [†]
Sex, Female	0.772 ± 0.088 [†]	0.769 ± 0.088 [†]	0.766 ± 0.088 [†]
Age	-0.048 ± 0.058	-0.050 ± 0.058	-0.052 ± 0.058
Age ²	-0.001 ± 0.000	-0.001 ± 0.000	-0.001 ± 0.000
Household size	0.037 ± 0.052	0.037 ± 0.052	0.037 ± 0.052
Routine manual	-0.633 ± 0.095 [†]	-0.623 ± 0.095 [†]	-0.618 ± 0.095 [†]
Wealth mid-tertile	0.310 ± 0.109 [†]	0.305 ± 0.109 [†]	0.301 ± 0.109 [†]
Wealth top-tertile	0.598 ± 0.116 [†]	0.587 ± 0.116 [†]	0.582 ± 0.116 [†]
Age left edu	0.184 ± 0.019 [†]	0.18 ± 0.0190 [†]	0.177 ± 0.019 [†]
Married-cohab.	0.105 ± 0.203	0.106 ± 0.203	0.105 ± 0.203
Separated-div.	0.137 ± 0.229	0.135 ± 0.229	0.130 ± 0.229
Widow	0.158 ± 0.219	0.158 ± 0.219	0.158 ± 0.219
Social support	0.175 ± 0.399	0.177 ± 0.399	0.177 ± 0.399
Social connection	0.064 ± 0.012 [†]	0.063 ± 0.012 [†]	0.063 ± 0.012 [†]
CVD	-0.084 ± 0.108	-0.086 ± 0.108	-0.086 ± 0.108
COPD	0.166 ± 0.117	0.168 ± 0.117	0.169 ± 0.116
Cancer	0.329 ± 0.178	0.323 ± 0.178	0.319 ± 0.178
Smoking	-0.351 ± 0.129 [†]	-0.346 ± 0.129 [†]	-0.341 ± 0.129 [†]
Drinking	0.478 ± 0.091 [†]	0.475 ± 0.091 [†]	0.475 ± 0.091 [†]
Exercise, vigorous	0.087 ± 0.110	0.088 ± 0.110	0.089 ± 0.110
Exercise, moderate	0.386 ± 0.091 [†]	0.386 ± 0.091 [†]	0.385 ± 0.091 [†]
Childhood lack facility	-0.118 ± 0.024 [†]	-0.117 ± 0.024 [†]	-0.117 ± 0.024 [†]
Adult illness episodes	-0.054 ± 0.029	-0.192 ± 0.029 [†]	-0.273 ± 0.118*
<i>RMSE</i>	0.0406	0.0189	0.0405

Source: The English Longitudinal Study of Ageing.

* $p < 0.05$.

[†] $p < 0.01$.

Table F. Three models on mental health in later life, with adult health (fully adjusted)

Predictors	Uncorrected	Bias-corrected	Error-corrected
Constant	1.676 ± 1.209	1.871 ± 1.209	1.664 ± 1.210
Sex, Female	-0.317 ± 0.050 [†]	-0.318 ± 0.050 [†]	-0.316 ± 0.050 [†]
Age	0.089 ± 0.033 [†]	0.088 ± 0.033 [†]	0.089 ± 0.033 [†]
Age ²	-0.001 ± 0.000*	-0.001 ± 0.000*	-0.001 ± 0.000*
Household size	0.042 ± 0.030	0.042 ± 0.030	0.042 ± 0.030
Routine manual	-0.084 ± 0.054	-0.079 ± 0.054	-0.084 ± 0.054
Wealth mid-tertile	0.236 ± 0.062 [†]	0.233 ± 0.062 [†]	0.236 ± 0.062 [†]
Wealth top-tertile	0.442 ± 0.066 [†]	0.437 ± 0.066 [†]	0.442 ± 0.066 [†]
Age left edu	0.010 ± 0.011	0.008 ± 0.011	0.010 ± 0.011
Married-cohab.	0.109 ± 0.116	0.109 ± 0.116	0.108 ± 0.116
Separated-div.	-0.205 ± 0.131	-0.206 ± 0.131	-0.205 ± 0.131
Widow	-0.528 ± 0.125 [†]	-0.528 ± 0.125 [†]	-0.529 ± 0.125 [†]
Social support	1.037 ± 0.228 [†]	1.038 ± 0.228 [†]	1.037 ± 0.228 [†]
Social connection	0.030 ± 0.007 [†]	0.030 ± 0.007 [†]	0.030 ± 0.007 [†]
CVD	-0.071 ± 0.062	-0.072 ± 0.062	-0.071 ± 0.062
COPD	-0.359 ± 0.066 [†]	-0.358 ± 0.066 [†]	-0.359 ± 0.066 [†]
Cancer	-0.005 ± 0.102	-0.008 ± 0.102	-0.005 ± 0.102
Smoking	-0.262 ± 0.074 [†]	-0.260 ± 0.074 [†]	-0.262 ± 0.074 [†]
Drinking	0.099 ± 0.052	0.098 ± 0.052	0.100 ± 0.052
Exercise, vigorous	0.079 ± 0.063	0.080 ± 0.063	0.079 ± 0.063
Exercise, moderate	0.443 ± 0.052 [†]	0.443 ± 0.052 [†]	0.443 ± 0.052 [†]
Childhood lack facility	-0.011 ± 0.016	-0.074 ± 0.016 [†]	-0.008 ± 0.017
Adult illness episodes	-0.318 ± 0.024 [†]	-0.317 ± 0.024 [†]	-0.318 ± 0.024 [†]
<i>RMSE</i>	0.0232	0.0107	0.0232

Source: The English Longitudinal Study of Ageing.

* $p < 0.05$.

† $p < 0.01$.