

Table S2. Discrepant recommendations of cancer genomic tests between organizations, by order of statement issuance

| Test | Purpose | First recommendation statement | | | Second recommendation statement | | | Third recommendation statement | | | Fourth recommendation statement | | |
|--------------------|--|--------------------------------|--------|-----------------------|---------------------------------|--------|-----------------------|--------------------------------|--------|-----------------------|---------------------------------|--------|-----------------------|
| | | Year | Source | Category ^a | Year | Source | Category ^a | Year | Source | Category ^a | Year | Source | Category ^a |
| BRAF | Pharmacogenomic for anti-EGFR therapy in colon cancer | 2011 | NCCN | 2 | 2013 | NCCN | 2 | | | | | | |
| | | | | | | EGAPP | 3 | | | | | | |
| CYP2D6 | Pharmacogenomic for tamoxifen use in breast cancer | 2005 | ASCO | 3 | 2010 | NCAB | 2 | 2011 | BCBS | 3 | | | |
| | | | | | | ASCO | 3 | | AHS | 3 | | | |
| ERRC1 | Prognostic / Pharmacogenomic use in NSCLC | 2009 | ASCO | - / 3 | 2011 | NCCN | - / 3 | 2013 | NCCN | 2 / 2 | | | |
| Oncotype DX | Prognostic / Pharmacogenomic use in lymph node-negative, ER-positive breast cancer | 2007 | ASCO | 2 / 2 | 2009 | NACB | 1 / 2 | 2011 | NCCN | 2 / 2 | 2013 | NICE | 1 / 1 |
| | | | | | | EGAPP | - / 3 | | | | | | |
| RRM1 | Prognostic / Pharmacogenomic use in NSCLC | 2009 | ASCO | - / 3 | 2010 | NCCN | 2 / 2 | 2011 | NCCN | 2 / 2 | | | |
| UGT1A1 | Pharmacogenomic use for irinotecan dosing in colon cancer | 2009 | EGAPP | 3 | 2010 | NCAB | 2 | | | | | | |

^a The - symbol indicates that the corresponding organization did not comment on the tests' prognostic purpose.