

S7 Table.

	Gene	qRT-PCR fold change (SF370SmR/CEM1Δ4)	RNA-seq fold change (SF370SmR/CEM1Δ4)*
37 °C Early Log	<i>nga</i>	2.0	2.6
	<i>slo</i>	4.5	3
	<i>norA</i>	-14.3	-18.5
	<i>emm</i>	11.9	11.1
	<i>speB</i>	**	**
	<i>hasB</i>	2.1	6.2
37 °C Late Log	<i>nga</i>	-1.9	1.5
	<i>slo</i>	-2.7	1.2
	<i>norA</i>	3.5	3.1
	<i>emm</i>	36.3	11.4
	<i>speB</i>	1.0	1
	<i>hasB</i>	-1.6	1
39 °C Early Log	<i>nga</i>	3.1	2.3
	<i>slo</i>	7.9	3.6
	<i>norA</i>	-3.8	-6.8
	<i>emm</i>	27	12.2
	<i>speB</i>	**	**
	<i>hasB</i>	1.7	4.9
39 °C Late Log	<i>nga</i>	1.9	1.4
	<i>slo</i>	1.0	1.3
	<i>norA</i>	-1.6	-2.8
	<i>emm</i>	79	5
	<i>speB</i>	251	143.9
	<i>hasB</i>	-4.4	-2.8

* Values calculated by Gene Sifter as described in Methods. The identical cDNA preparations were used for qRT-PCR analysis and RNA-seq analysis

** Not expressed in EL