

## Pathogen distribution and drug resistance in a burn ward

**Supplementary Table 1.** Annual distribution of Gram-positive and negative bacteria and fungi detected in clinical samples

Year	Strains identified (n)	Gram-positive bacteria [n (%)]	Gram-negative bacteria [n (%)]	Fungi [n (%)]
Wound Secretion Culture				
2011	468	168 (35.9)	259 (55.3)	41 (8.8)
2012	518	188 (36.3)	253 (48.8)	77 (14.9)
2013	480	187 (39)	250 (52.1)	43 (8.9)
Sputum Culture				
2011	56	9 (16.1)	46 (82.1)	1 (1.8)
2012	41	10 (24.4)	30 (73.2)	1 (2.4)
2013	56	7 (12.5)	48 (85.7)	1 (1.8)
Urine Culture				
2011	100	13 (13)	62 (2)	25 (25)
2012	116	22 (19)	38 (32.8)	56 (48.2)
2013	119	27 (22.7)	59 (49.6)	33 (27.7)
Stool Culture				
2011	2	0 (0)	1 (50)	1 (50)
2012	3	0 (0)	0 (0)	3 (100)
2013	0	0 (0)	0 (0)	0 (0)
Blood Culture				
2011	49	26 (53)	22 (44.9)	1 (1.1)
2012	35	20 (57.1)	12 (34.3)	3 (8.6)
2013	44	23 (52.3)	20 (45.5)	1 (2.2)
Deep vein catheter culture				
2011	48	19 (39.6)	27 (56.3)	2 (4.1)
2012	37	17 (46)	16 (43)	4 (11)
2013	40	14 (35)	23 (57.5)	3 (7.5)