

Supplemental Data

Inhibition of SET Domain-Containing Lysine Methyltransferase 7/9 Ameliorates Renal Fibrosis

Kensuke Sasaki,¹ Shigehiro Doi,^{1*} Ayumu Nakashima,¹ Taisuke Irifuku,¹ Kyoko Yamada,¹ Keiko Kokoroishi,¹ Toshinori Ueno,¹ Toshiki Doi,¹ Eisuke Hida,² Koji Arihiro,³ Nobuoki Kohno,⁴ and Takao Masaki^{1*}

¹Department of Nephrology, Hiroshima University Hospital, Hiroshima, Japan; ²Center for Integrated Medical Research, Hiroshima University Hospital, Hiroshima, Japan;

³Department of Pathology, Hiroshima University Hospital, Hiroshima, Japan; and

⁴Department of Molecular and Internal Medicine, Graduate School of Biomedical & Health Sciences, Hiroshima University, Hiroshima, Japan

***CORRESPONDENCE TO:**

Shigehiro Doi, M.D., Ph.D. Assistant Professor

Department of Nephrology, Hiroshima University Hospital

1-2-3 Kasumi, Minami-Ku, Hiroshima 734-8551, Japan

E-mail: sdoi@hiroshima-u.ac.jp

TEL: +81-82-257-5960

FAX: +81-82-256-5560

Takao Masaki, M.D., Ph.D. Professor

Department of Nephrology, Hiroshima University Hospital

1-2-3 Kasumi, Minami-Ku, Hiroshima 734-8551, Japan

E-mail: masakit@hiroshima-u.ac.jp

TEL: +81-82-257-1506

FAX: +81-82-257-1508

This Supplemental Data includes:

Supplemental Table 1 - 2

Table 1. Clinical characteristics of IgAN patients

eGFR, mL/min/1.73m ²	83.0 - 118.0 (n = 8)	64.9 - 80.3 (n = 6)	40.2 - 64.0 (n = 12)
Age, years	28 ± 10	34 ± 9	47 ± 10
Sex, M/F	5/3	4/2	10/2
BMI, kg/m ²	21 ± 4.2	21 ± 1.5	24 ± 2.5
Ccr, mL/min	114.6 ± 18.43	82.27 ± 10.74	76.55 ± 19.77
eGFR, mL/min/1.73m ²	99 ± 12	74 ± 5	54 ± 7
Urinary protein, g/24 h	0.19 ± 0.12	0.51 ± 0.42	0.99 ± 0.81
Renal fibrosis, %	4.4 ± 2.7	9.9 ± 1.1	15.2 ± 4.4

Values indicate means ± SD

BMI, body mass index; Ccr, creatinine clearance; eGFR, estimated glomerular filtration rate

Table 2. Clinical characteristics of MN patients

eGFR, mL/min/1.73m ²	75.0 - 84.5 (n = 4)	64.6 - 74.4 (n = 4)	48.6 - 63.4 (n = 4)
Age, years	66 ± 9	66 ± 7	73 ± 4
Sex, M/F	2/2	3/1	3/1
BMI, kg/m ²	22 ± 4.3	23 ± 1.1	24 ± 1.2
Ccr, mL/min	103.9 ± 20.85	89.1 ± 13.06	74.1 ± 9.70
eGFR, mL/min/1.73m ²	79 ± 3.5	71 ± 3.8	56 ± 5.4
Urinary protein, g/24 h	4.37 ± 3.15	2.86 ± 1.07	3.35 ± 1.31
Renal fibrosis, %	6.7 ± 1.5	8.2 ± 1.2	8.8 ± 0.5

Values indicate means ± SD

BMI, body mass index; Ccr, creatinine clearance; eGFR, estimated glomerular filtration rate