

S2 Table. The induction of resistance of *Enterococcus faecalis* and *Streptococcus agalactiae* to compounds DR5047 and DR5026. Minimum inhibitory concentrations of the tested LPPOs after each induction step. The induction of resistance was carried out by repeated passages of *E. faecalis* and *S. agalactiae* strains with subinhibitory concentrations of the relevant LPPOs. A total of 14 passages were performed. The results clearly showed that the MICs remained relatively unchanged. It meant that the resistance was not induced.

Step	MIC (mg/L)			
	<i>Streptococcus agalactiae</i>		<i>Enterococcus faecalis</i> CCM 4224	
	DR5026	DR5047	DR5026	DR5047
1	3.1	6.2	6.2	12.5
2	3.1	6.2	6.2	12.5
3	3.1	6.2	6.2	12.5
4	3.1	6.2	6.2	12.5
5	3.1	6.2	6.2	25
6	3.1	6.2	12.5	25
7	6.2	3.1	12.5	25
8	6.2	6.2	12.5	25
9	3.1	6.2	12.5	25
10	3.1	6.2	12.5	50
11	3.1	6.2	12.5	25
12	6.2	6.2	12.5	25
13	6.2	6.2	12.5	25
14	6.2	6.2	12.5	25