Cognitive ageing in Great Britain in the new century: Supplementary Tables

Gindo Tampubolon[1]

1 University of Manchester

Table 1. Sensitivity analysis using decadal cohorts (ELSA 2002-2013)

Predictors	Episodic memory	
	Coeff.	Std. error
Constant	-6.9057	0.8725
Age	0.4097	$0.0270^{\dagger}$
Age^2	-0.0033	$0.0002^{\dagger}$
Cohort 60s	0.8093	$0.1110^{\dagger}$
Cohort 70s	1.6935	$0.1235^{\dagger}$
Cohort 80s	2.9422	$0.1337^{\dagger}$
Female	0.7232	$0.0467^{\dagger}$
Intermediate	0.9496	$0.0578^{\dagger}$
Managerial	1.1044	$0.0547^{\dagger}$
Married/cohab.	-0.0345	0.0435
College	0.9514	$0.0542^{\dagger}$
Middle tertile	0.1710	$0.0370^{\dagger}$
Top tertile	0.3484	$0.0424^{\dagger}$
Ethnic minority	-1.4015	$0.1377^{\dagger}$
Social connect.	0.0710	$0.0184^{\dagger}$
Physical problem	-0.0385	$0.0062^{\dagger}$
Hypertensives	-0.0754	0.0381
Diabetes	-0.1747	$0.0624^{\dagger}$
Cancer	0.0793	0.0637
CVD	0.0290	0.0473
Stroke	-0.6062	$0.0889^{\dagger}$
Arthritis	0.2173	$0.0404^{\dagger}$
Smoker	-0.0403	0.0435
Drink regularly	0.4743	$0.0508^{\dagger}$
Exercise mod.	0.2554	$0.0348^{\dagger}$
Exercise vig.	0.1043	$0.0325^{\dagger}$
CESD	-0.0770	$0.0083^{\dagger}$
Practice	0.6392	$0.0722^{\dagger}$
Within-person σ	0.8043	$0.1348^{\dagger}$
R^2	0.531	

† $p < 0.01$.