

Table S4. Univariate and multivariate predictors of functional outcome assuming the worst scenario for patients lost to follow up

Predictors	48h			
	univariate		multivariate	
	OR(CI 95%)	<i>p</i>	OR(CI 95%)	<i>p</i>
Age	1.01(0.97- 1.05)	0.58	-	-
Gender	0.82(0.34-1.99)	0.66	-	-
Hypertension	0.32(0.12-0.84)	0.02	0.31(0.11- 0.87)	0.03
Diabetes	1.18(0.47-2.94)	0.73	-	-
Dyslipidemia	0.53(0.22-1.29)	0.16	-	-
Cardiovascular Diseases	1.05(0.17-6.67)	0.96	-	-
Atrial Fibrillation	0.75(0.23-2.43)	0.63	-	-
Smoking History	1.06(0.35-3.05)	0.90	-	-
Toast Classification	1.01(0.85-1.43)	0.47	-	-
NIHSS	1.14(1.06-1.24)	0.001	1.14(1.06-1.24)	0.001
Ficolin-1	1.55(0.13-18.01)	0.73	-	-
Ficolin-3	0.95(0.88-1.03)	0.21	-	-

Exact *p* value for univariate and multivariate logistic regression analysis is reported. 95% Confidence Interval (CI 95%); Odds Ratio (OR). NIHSS: National Institutes of Health Stroke Scale; 3- month mRS: 3-month modified Rankin