Checklist S1. PRISMA 2009 Checklist

| Section/topic | pic # Checklist item | | | | |
|------------------------------------|--|---|-----|--|--|
| Title | | | | | |
| Title | 1 | Identify the report as a systematic review, meta-analysis, or both. | 1 | | |
| Abstract: | | | | | |
| Structured summary | 2 | Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number. | 3 | | |
| Introduction | | | | | |
| Rationale | 3 | Describe the rationale for the review in the context of what is already known. | 5 | | |
| Objectives | 4 | Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS). | 5-6 | | |
| Methods | | | | | |
| Protocol and registration | 5 | Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number. | 6 | | |
| Eligibility criteria | 6 | Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale. | 7 | | |
| Information sources | 7 | Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched. | 6 | | |
| Search | 8 | Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated. | 6 | | |
| Study selection | 9 | State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis). | 6 | | |
| Data collection process | 10 | Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators. | 7 | | |
| Data items | 11 | List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made. | 7 | | |
| Risk of bias in individual studies | 12 | Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis. | 8 | | |
| Summary measures | 13 | State the principal summary measures (e.g., risk ratio, difference in means). | 7 | | |
| Synthesis of results | 14 | Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I²) for each meta-analysis. | 7 | | |
| Risk of bias across studies | 15 | Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies). | 7 | | |
| Additional analyses | analyses 16 Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified. | | | | |

| Results | | | | | | | |
|-------------------------------|----|--|-------|--|--|--|--|
| Study selection | 17 | Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram. | | | | | |
| Study characteristics | 18 | For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations. | 9 | | | | |
| Risk of bias within studies | 19 | Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12). | 9-10 | | | | |
| Results of individual studies | 20 | For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot. | 10 | | | | |
| Synthesis of results | 21 | Present results of each meta-analysis done, including confidence intervals and measures of consistency. | 10 | | | | |
| Risk of bias across studies | 22 | Present results of any assessment of risk of bias across studies (see Item 15). | 10-11 | | | | |
| Additional analysis | 23 | Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]). | 10-11 | | | | |
| Discussion | | | | | | | |
| Summary of evidence | 24 | Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers). | 12-15 | | | | |
| Limitations | 25 | Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias). | 15 | | | | |
| Conclusions | 26 | Provide a general interpretation of the results in the context of other evidence, and implications for future research. | 15 | | | | |
| Funding | | | | | | | |
| Funding | 27 | Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review. | 1 | | | | |
| | | | | | | | |

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097. For more information, visit: www.prisma-statement.org.


Supplementary Table 1. Risk of bias of included trials

| Study Representative exposed coho | | Selection of non- exposed cohort | Exposure | Outcome of interest not present at start | Comparability of LH vs. OH | Assessment of outcome | Follow up | Adequacy of follow- up/missing data | Score | |
|-----------------------------------|----------------------|-------------------------------------|------------------|--|--|-----------------------|-----------|--|-------|--|
| Belli et al. | Truly representative | Same | Surgical records | Yes | Restrictions in exophytic or subcapsular tumors, not matched | Record linkage | 3 yr | Unclear | 7 | |
| Cheung et al. | Truly representative | Same | Surgical records | Yes | No restrictions, matched | Record linkage | 5 yr | None | 8 | |
| Kanazawa et al. | Truly representative | Same | Surgical records | Yes | No restrictions, not matched | Record linkage | 5 yr | Unclear | 7 | |
| Memeo et al. | Truly representative | Same | Surgical records | Yes | No restrictions, matched | Record linkage | 10 yr | Unclear | 7 | |
| Siniscalchi et al. | Truly representative | Same | Surgical records | Yes | Restrictions in tumors diameters and should located in the anterior or lateral segments II-VI, not matched | Record linkage | 7 d | Clear | 7 | |
| Truant et al. | Truly representative | Same | Surgical records | Yes | Restrictions in subcapsular tumors located in the anterior or lateral seg- ments II-VI, matched | Record linkage | 5 yr | None | 7 | |
| Yamashita et al. | Truly representative | Same | Surgical records | Yes | No restrictions, not matched | Record linkage | 5 yr | Unclear | 7 | |


A Tumor margin

| | Laparoscop | Open Hepatectomy | | | Mean Difference | | | Mean Difference | | | | | |
|--|------------|------------------|-------|------|-----------------|-------|--------|--------------------|-------------|---------|--------------|------|-----|
| Study or Subgroup | Mean | SD | Total | Mean | SD | Total | Weight | IV. Fixed, 95% C | | IV. Fix | ced. 95° | % CI | |
| Cheung 2013 | 0.95 | 0.75 | 32 | 0.8 | 0.875 | 64 | 6.1% | 0.15 [-0.19, 0.49] | | | + | | |
| Kanazawa 2013 | 0.5 | 0.45 | 28 | 0.3 | 0.375 | 28 | 14.7% | 0.20 [-0.02, 0.42] | | | † · | | |
| Memeo 2014 | 1 | 1.25 | 45 | 0.6 | 1.45 | 45 | 2.2% | 0.40 [-0.16, 0.96] | | | 1 | | |
| Truant 2011 | 0.95 | 0.28 | 36 | 0.86 | 0.17 | 53 | 66.2% | 0.09 [-0.01, 0.19] | | | | | |
| Yamashita 2014 | 0.74 | 0.87 | 63 | 0.58 | 0.69 | 99 | 10.7% | 0.16 [-0.09, 0.41] | | | † | | |
| Total (95% CI) | | | 204 | | | 289 | 100.0% | 0.12 [0.04, 0.21] | | | | | |
| Heterogeneity: Chi ² = 1.93, df = 4 (P = 0.75); I ² = 0% | | | | | | | | | | -50 | _ | 50 | 100 |
| Test for overall effect: Z = 2.92 (P = 0.003) | | | | | | | | -100 | Favours [Ol | H) Favo | | 100 | |


B Operative time



C Blood loss




D Blood transfusion




Supplementary Figure 1. Meta-analysis of data on opeartion outcomes in LH and OH.


A Postoperative mobility




B Postoperative mortality



C Curative resection



D Length of hospital stay



Supplementary Figure 2. Meta-analysis of data on short-term outcomes in LH and OH.