


SUPPLEMENTAL MATERIALS


Title: The role of erythrocyte CD47 in intracerebral hematoma clearance

Supplemental Figures




Supplemental Figure I: Double labeling of CD86 and CD206 in the peri-hematoma area at day-3 after 30 μ l blood injected into right basal ganglia, scale bar = 50 μ m.

The primary antibodies were: polyclonal rabbit anti-CD206 IgG (Abcam, USA, 1:200 dilution) and monoclonal rat anti-CD86 IgG (Thermo Scientific, USA; 1:100 dilution). The secondary antibodies were Alexa Fluor 594-conjugated donkey anti-rabbit mAb and Alexa Fluor 488-conjugated donkey anti-rat mAb (Invitrogen, Grand Island, NY, USA; both at a 1:500 dilution).



Supplemental Figure II: Immunoreactivity of Iba-1 in the ipsilateral basal ganglia at day-3 after injection of 30 μ l saline with or without 5 μ l clodronate liposome into the right basal ganglia, scale bar = 50 μ m.



Supplemental Figure III: Immunoreactivity of Iba-1 in the ipsilateral basal ganglia at day-3 after injection of $30\mu\text{l}$ saline with or without $5\mu\text{l}$ clodronate liposome into the right basal ganglia, scale bar = $20\mu\text{m}$.