

# A cross modal performance-based measure of sensory stimuli intricacy

Kobi Snitz<sup>1\*,</sup>, Anat Arzi<sup>1,</sup>, Merav Jacobson<sup>1,</sup>, Lavi Secundo<sup>1,</sup>, Kineret Weissler<sup>1,</sup>,  
Adi Yablonka<sup>1,</sup>

**1 Dept of Neurobiology, Weizmann Institute of Science, Rehovot, Israel**

**These authors contributed equally to this work.**

\* [kobi.snitz@weizmann.ac.il](mailto:kobi.snitz@weizmann.ac.il)

## 0.1 S1 Table

**Data set A odorants.**

Name	CAS number	mean variance
Isoamyl acetate	123-92-2	0.8119
Nonane	111-84-2	0.5398
Ethyl valerate	539-82-2	0.9723
5-methyl-2-hexanone	110-12-3	1.3859
Isopropylbenzene (Cumene)	98-82-8	0.7061
1-pentanol	71-41-0	0.6954
1,7-octadiene	3710-30-3	0.8330
2-heptanone	110-43-0	0.6895
4-methyl-3-penten-2-one (mesityl oxide)	141-79-7	0.8428
3-methyl-2-buten-1-ol	556-82-1	0.5233