

Authors:

Kalani L. Raphael, MD, MS¹, Rachel A. Murphy, PhD², Michael G. Shlipak, MD, MPH³, Suzanne Satterfield, MD, DrPh⁴, Hunter K. Huston, MD¹, Anthony Sebastian, MD⁵, Deborah E. Sellmeyer, MD⁶, Kushang V. Patel, PhD, MPH⁷, Anne B. Newman, MD, MPH⁸, Mark J. Sarnak, MD, MS⁹, Joachim H. Ix, MD, MAS¹⁰, and Linda F. Fried, MD, MPH¹¹ for the Health ABC Study

Affiliations:

¹Salt Lake City Veterans Affairs Healthcare System and Department of Internal Medicine, University of Utah, Salt Lake City, Utah;

²Laboratory of Epidemiology and Population Sciences, Intramural Research Program, National Institute on Aging, Bethesda, Maryland;

³San Francisco Veterans Affairs Medical Center and Department of Medicine, University of California San Francisco, San Francisco, California;

⁴Department of Preventive Medicine, University of Tennessee Health Science Center, Memphis, Tennessee;

⁵Department of Medicine, Division of Nephrology, University of California San Francisco, San Francisco, California;

⁶Division of Endocrinology, Department of Medicine, Johns Hopkins Hospital, Johns Hopkins University, Baltimore, Maryland;

⁷Department of Anesthesiology and Pain Medicine, University of Washington, Seattle, Washington;

⁸Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania;

⁹Division of Nephrology, Department of Medicine, Tufts Medical Center, Boston, Massachusetts;

¹⁰Division of Nephrology-Hypertension, Department of Medicine, University of California San Diego and Veterans Affairs San Diego Healthcare System, San Diego, California;
and

¹¹VA Pittsburgh Healthcare System and Department of Medicine and Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania.