

## SUPPLEMENTARY TABLE

**Supplementary Table S1: Blood examination of tumor-bearing nude mice after treatment with varying formulations at day 22 after inoculation**

Assay	Physiological saline	Vincristine liposomes	Vincristine plus dasatinib liposomes	Functional vincristine plus dasatinib liposomes	Free vincristine
WBC <sup>a)</sup> (10 <sup>9</sup> /L)	5.57 ± 1.51	6.20 ± 1.35	7.80 ± 2.02	5.57 ± 1.16	3.53 ± 0.46*
RBC <sup>b)</sup> (10 <sup>12</sup> /L)	7.75 ± 0.46	7.13 ± 0.77	7.81 ± 1.07	8.30 ± 1.02	8.59 ± 2.96
MCHC <sup>c)</sup> (g/L)	304.67 ± 1.53	299.00 ± 20.66	298.33 ± 2.31	291.00 ± 12.53	299.67 ± 26.50
RDW <sup>d)</sup> (%)	14.10 ± 0.92	13.30 ± 0.50	13.33 ± 0.45	13.23 ± 0.12	13.67 ± 1.98
LYM <sup>e)</sup> (10 <sup>9</sup> /L)	0.93 ± 0.84	2.77 ± 2.25	1.13 ± 0.45	1.77 ± 0.15	0.50 ± 0.20
MID <sup>f)</sup> (10 <sup>9</sup> /L)	1.30 ± 0.78	1.33 ± 0.76	1.03 ± 0.42	0.90 ± 0.10	0.50 ± 0.30

<sup>a)</sup>White blood cells; <sup>b)</sup>Red blood cells; <sup>c)</sup>Mean corpuscular hemoglobin concentration; <sup>d)</sup>Red cell distribution width;

<sup>e)</sup>Lymphocytes; <sup>f)</sup>Intermediate cell number. Data are presented as the mean ± SD (*n* = 3). \**p* < 0.05, vs. physiological saline.