

Table S1. Representative Examples of Functional Effects of mtDNA Haplogroups

CONTINENT	POPULATION	ENDURANCE	SPRINT/POWER	REFERENCE
Africa	Kenyans	↑L0		(Scott et al., 2009)
	African-Americans		↑non-L	(Deason et al., 2012)
Europe	Finnish	↑I	↑Uk, ↑J	(Niemi and Majamaa, 2005)
	Polish	↑H, ↑16080C, ↓16362C		(Maruszak et al., 2014)
Asia	Spanish	↑V		(Nogales-Gadea et al., 2011)
	Japanese	↑G1	↑F	(Mikami et al., 2011)
	Korean	↑M*, ↑N9, ↓B		(Kim et al., 2012)

Table S2. Representative Examples of Haplogroup Associations which Predisposition to Clinical Phenotypes

Clinical Phenotype	Haplogroup	Effect	References
LHON	J	↑ risk	(Brown et al., 1997; Brown et al., 1995; Johns et al., 1992; Torroni et al., 1997)
AMD, drusen levels, retinal pigment	J + U	↑ risk	(Udar et al., 2009), (Jones et al., 2007)
AMD	H	↓ risk	(Jones et al., 2007)
Longevity, Europeans	J	Increased	(De Benedictis et al., 1999; Ivanova et al., 1998; Niemi et al., 2003; Rose et al., 2001)
Longevity, Europeans	J2 & U3, CR 150C>T	Increased	(Chen et al., 2012).
Longevity, Asians	D	Increased	(Tanaka et al., 1998)
AD & PD	J + Uk	↓ risk	(Ghezzi et al., 2005; van der Walt et al., 2004; van der Walt et al., 2003)
PD	H	↑ risk	(Khusnutdinova et al., 2008)
AD & PD	H5a, tRNA ^{Gln} 4336A>G	↑ risk	(Shoffner et al., 1993)

Diabetes, Europeans	J	↑ risk	(Crispim et al., 2006)
Diabetes, Asians	N9a	↓ risk	(Fuku et al., 2007)
Myocardial Infarction, Asians	N9a	↑ risk	(Nishigaki et al., 2007)
Sepsis	H	↓ risk	(Baudouin et al., 2005)
Serum IgE levels	U	increased	(Raby et al., 2007)
AIDS Progression	Uk	impedes	(Hendrickson et al., 2008)
Osteoarthritis	J + T	↓ risk	(Soto-Hermida et al., 2014)
Prostate & Renal Cancer	U	↑ risk	(Booker et al., 2006)
Breast Cancer	N, 10398G>A (A>T)	↑ risk	(Darvishi et al., 2007)
Breast Cancer	M, D5	↑ risk	(Fang et al., 2010)
Thyroid Cancer	D4a	↑ risk	(Fang et al., 2010)

REFERENCES

- Baudouin, S.V., Saunders, D., Tiangyou, W., Elson, J.L., Poynter, J., Pyle, A., Keers, S., Turnbull, D.M., Howell, N., and Chinnery, P.F. (2005). Mitochondrial DNA and survival after sepsis: a prospective study. *Lancet* 366, 2118-2121.
- Booker, L.M., Habermacher, G.M., Jessie, B.C., Sun, Q.C., Baumann, A.K., Amin, M., Lim, S.D., Fernandez-Golarz, C., Lyles, R.H., Brown, M.D., *et al.* (2006). North American white mitochondrial haplogroups in prostate and renal cancer. *J Urol* 175, 468-472; discussion 472-473.
- Brown, M.D., Sun, F., and Wallace, D.C. (1997). Clustering of Caucasian Leber hereditary optic neuropathy patients containing the 11778 or 14484 mutations on an mtDNA lineage. *Am J Hum Genet* 60, 381-387.
- Brown, M.D., Torroni, A., Reckord, C.L., and Wallace, D.C. (1995). Phylogenetic analysis of Leber's hereditary optic neuropathy mitochondrial DNA's indicates multiple independent occurrences of the common mutations. *Hum Mutat* 6, 311-325.
- Chen, A., Raule, N., Chomyn, A., and Attardi, G. (2012). Decreased reactive oxygen species production in cells with mitochondrial haplogroups associated with longevity. *PLoS One* 7, e46473.
- Crispim, D., Canani, L.H., Gross, J.L., Tschiedel, B., Souto, K.E., and Roisenberg, I. (2006). The European-specific mitochondrial cluster J/T could confer an increased risk of insulin-resistance and type 2 diabetes: an analysis of the m.4216T > C and m.4917A > G variants. *Ann Hum Genet* 70, 488-495.
- Darvishi, K., Sharma, S., Bhat, A.K., Rai, E., and Bamezai, R.N. (2007). Mitochondrial DNA G10398A polymorphism imparts maternal Haplogroup N a risk for breast and esophageal cancer. *Cancer Lett* 249, 249-255.
- De Benedictis, G., Rose, G., Carrieri, G., De Luca, M., Falcone, E., Passarino, G., Bonafe, M., Monti, D., Baggio, G., Bertolini, S., *et al.* (1999). Mitochondrial DNA inherited variants are associated with successful aging and longevity in humans. *FASEB J* 13, 1532-1536.
- Deason, M., Scott, R., Irwin, L., Macaulay, V., Fuku, N., Tanaka, M., Irving, R., Charlton, V., Morrison, E., Austin, K., *et al.* (2012). Importance of mitochondrial haplotypes and maternal lineage in sprint performance among individuals of West African ancestry. *Scand J Med Sci Sports* 22, 217-223.
- Fang, H., Shen, L., Chen, T., He, J., Ding, Z., Wei, J., Qu, J., Chen, G., Lu, J., and Bai, Y. (2010). Cancer type-specific modulation of mitochondrial haplogroups in breast, colorectal and thyroid cancer. *BMC cancer* 10, 421.
- Fuku, N., Park, K.S., Yamada, Y., Nishigaki, Y., Cho, Y.M., Matsuo, H., Segawa, T., Watanabe, S., Kato, K., Yokoi, K., *et al.* (2007). Mitochondrial haplogroup N9a confers resistance against type 2 diabetes in Asians. *Am J Hum Genet* 80, 407-415.

- Ghezzi, D., Marelli, C., Achilli, A., Goldwurm, S., Pezzoli, G., Barone, P., Pellecchia, M.T., Stanzone, P., Brusa, L., Bentivoglio, A.R., *et al.* (2005). Mitochondrial DNA haplogroup K is associated with a lower risk of Parkinson's disease in Italians. *Eur J Hum Genet* 13, 748-752.
- Hendrickson, S.L., Hutcheson, H.B., Ruiz-Pesini, E., Poole, J.C., Lautenberger, J., Sezgin, E., Kingsley, L., Goedert, J.J., Vlahov, D., Donfield, S., *et al.* (2008). Mitochondrial DNA haplogroups influence AIDS progression. *AIDS* 22, 2429-2439.
- Ivanova, R., Lepage, V., Charron, D., and Schachter, F. (1998). Mitochondrial genotype associated with French Caucasian centenarians. *Gerontology* 44, 349.
- Johns, D.R., Neufeld, M.J., and Park, R.D. (1992). An ND-6 mitochondrial DNA mutation associated with Leber hereditary optic neuropathy. *Biochem Biophys Res Commun* 187, 1551-1557.
- Jones, M.M., Manwaring, N., Wang, J.J., Rochtchina, E., Mitchell, P., and Sue, C.M. (2007). Mitochondrial DNA haplogroups and age-related maculopathy. *Arch Ophthalmol* 125, 1235-1240.
- Khusnutdinova, E., Gilyazova, I., Ruiz-Pesini, E., Derbeneva, O., Khusainova, R., Khidiyatova, I., Magzhanov, R., and Wallace, D.C. (2008). A mitochondrial etiology of neurodegenerative diseases: evidence from Parkinson's disease. *Ann N Y Acad Sci* 1147, 1-20.
- Kim, K.C., Cho, H.I., and Kim, W. (2012). MtDNA haplogroups and elite Korean athlete status. *Int J Sports Med* 33, 76-80.
- Maruszak, A., Adamczyk, J.G., Siewierski, M., Sozanski, H., Gajewski, A., and Zekanowski, C. (2014). Mitochondrial DNA variation is associated with elite athletic status in the Polish population. *Scand J Med Sci Sports* 24, 311-318.
- Mikami, E., Fuku, N., Takahashi, H., Ohiwa, N., Scott, R.A., Pitsiladis, Y.P., Higuchi, M., Kawahara, T., and Tanaka, M. (2011). Mitochondrial haplogroups associated with elite Japanese athlete status. *Br J Sports Med* 45, 1179-1183.
- Niemi, A.K., Hervonen, A., Hurme, M., Karhunen, P.J., Jylha, M., and Majamaa, K. (2003). Mitochondrial DNA polymorphisms associated with longevity in a Finnish population. *Hum Genet* 112, 29-33.
- Niemi, A.K., and Majamaa, K. (2005). Mitochondrial DNA and ACTN3 genotypes in Finnish elite endurance and sprint athletes. *Eur J Hum Genet* 13, 965-969.
- Nishigaki, Y., Yamada, Y., Fuku, N., Matsuo, H., Segawa, T., Watanabe, S., Kato, K., Yokoi, K., Yamaguchi, S., Nozawa, Y., *et al.* (2007). Mitochondrial haplogroup N9b is protective against myocardial infarction in Japanese males. *Hum Genet* 120, 827-836.
- Nogales-Gadea, G., Pinos, T., Ruiz, J.R., Marzo, P.F., Fiúza-Luces, C., Lopez-Gallardo, E., Ruiz-Pesini, E., Martin, M.A., Arenas, J., Moran, M., *et al.* (2011). Are mitochondrial haplogroups associated with elite athletic status? A study on a Spanish cohort. *Mitochondrion* 11, 905-908.

Raby, B.A., Klanderman, B., Murphy, A., Mazza, S., Camargo, C.A., Jr., Silverman, E.K., and Weiss, S.T. (2007). A common mitochondrial haplogroup is associated with elevated total serum IgE levels. *J Allergy Clin Immunol* 120, 351-358.

Rose, G., Passarino, G., Carrieri, G., Altomare, K., Greco, V., Bertolini, S., Bonafe, M., Franceschi, C., and De Benedictis, G. (2001). Paradoxes in longevity: sequence analysis of mtDNA haplogroup J in centenarians. *Eur J Hum Genet* 9, 701-707.

Scott, R.A., Fuku, N., Onywera, V.O., Boit, M., Wilson, R.H., Tanaka, M., Goodwin, W.H., and Pitsiladis, Y.P. (2009). Mitochondrial haplogroups associated with elite Kenyan athlete status. *Med Sci Sports Exerc* 41, 123-128.

Shoffner, J.M., Brown, M.D., Torroni, A., Lott, M.T., Cabell, M.R., Mirra, S.S., Beal, M.F., Yang, C., Gearing, M., Salvo, R., et al. (1993). Mitochondrial DNA variants observed in Alzheimer disease and Parkinson disease patients. *Genomics* 17, 171-184.

Soto-Hermida, A., Fernandez-Moreno, M., Oreiro, N., Fernandez-Lopez, C., Rego-Perez, I., and Blanco, F.J. (2014). mtDNA haplogroups and osteoarthritis in different geographic populations. *Mitochondrion* 15, 18-23.

Tanaka, M., Gong, J.S., Zhang, J., Yoneda, M., and Yagi, K. (1998). Mitochondrial genotype associated with longevity. *Lancet* 351, 185-186.

Torroni, A., Petrozzi, M., D'Urbano, L., Sellitto, D., Zeviani, M., Carrara, F., Carducci, C., Leuzzi, V., Carelli, V., Barboni, P., et al. (1997). Haplotype and phylogenetic analyses suggest that one European-specific mtDNA background plays a role in the expression of Leber hereditary optic neuropathy by increasing the penetrance of the primary mutations 11778 and 14484. *Am J Hum Genet* 60, 1107-1121.

Udar, N., Atilano, S.R., Memarzadeh, M., Boyer, D.S., Chwa, M., Lu, S., Maguen, B., Langberg, J., Coskun, P., Wallace, D.C., et al. (2009). Mitochondrial DNA haplogroups associated with age-related macular degeneration. *Invest Ophthalmol Vis Sci* 50, 2966-2974.

van der Walt, J.M., Dementieva, Y.A., Martin, E.R., Scott, W.K., Nicodemus, K.K., Kroner, C.C., Welsh-Bohmer, K.A., Saunders, A.M., Roses, A.D., Small, G.W., et al. (2004). Analysis of European mitochondrial haplogroups with Alzheimer disease risk. *Neurosci Lett* 365, 28-32.

van der Walt, J.M., Nicodemus, K.K., Martin, E.R., Scott, W.K., Nance, M.A., Watts, R.L., Hubble, J.P., Haines, J.L., Koller, W.C., Lyons, K., et al. (2003). Mitochondrial polymorphisms significantly reduce the risk of Parkinson disease. *Am J Hum Genet* 72, 804-811.