

Online Supplemental Table. Theory of Planned Behavior mean scores^a and standard deviations (SD) for adolescents and parents overall, adolescents by last grade completed (i.e. middle school or high school), and parents by BMI category from a Mixed Methods Analysis of Beverage Choices in Adolescents and Parents using the Theory of Planned Behavior.

	Adolescents (N=100)		Parents (N=66)	
Attitude	Overall	3.44 (0.57)	Overall	4.85 (1.29)
	Middle school	3.30 (0.50) ^b	Underweight	5.29 (1.02)
	High school	3.63 (0.61) ^b	Normal weight	5.02 (1.53)
			Overweight	4.46 (0.96)
			Obese	4.87 (1.16)
Subjective norm	Overall	3.59 (0.74)	Overall	4.56 (1.18)
	Middle school	3.59 (0.74)	Underweight	4.92 (1.20)
	High school	3.60 (0.74)	Normal weight	4.59 (1.29)
			Overweight	4.56 (1.04)
			Obese	4.43 (1.23)
Perceived behavioral control	Overall	4.48 (0.61)	Overall	6.52 (0.65)
	Middle school	4.34 (0.60) ^b	Underweight	6.17 (1.00)
	High school	4.66 (0.58) ^b	Normal weight	6.78 (0.47) ^c
			Overweight	6.54 (0.64)
			Obese	6.07 (0.64) ^c

Intention	Overall	3.45 (0.85)	Overall	5.69 (1.73)
	Middle school	3.34 (0.87)	Underweight	5.69 (2.63)
	High school	3.60 (0.80)	Normal weight	6.16 (1.37)
			Overweight	5.43 (1.83)
			Obese	5.04 (1.90)

^aResponses to the Theory of Planned Behavior Questionnaire were scored using a seven-point semantic differential scale in parents (i.e. "1" to "7") and a five-point semantic differential scale in adolescents (i.e. "1" to "5").

^bSignificantly different responses between middle school and high school participants, $p < 0.01$

^cSignificantly different responses between normal weight and obese parents, $p < 0.01$