

ONLINE SUPPLEMENT

PATTERNS AND CORRELATES OF BASELINE THIAZIDE-TYPE DIURETIC PRESCRIPTION IN THE SYSTOLIC BLOOD PRESSURE INTERVENTION TRIAL (SPRINT)

Authors & Affiliations:

Tara I. Chang, Stanford University School of Medicine
Gregory Evans, Wake Forest School of Medicine
Alfred K. Cheung, University of Utah
William C. Cushman, Memphis Veterans Affairs Medical Center
Matthew J Diamond, The Medical College of Georgia at Georgia Regents University
Jamie P. Dwyer, Vanderbilt University
Yonghong Huan, University of Pennsylvania
Dalane Kitzman, Wake Forest School of Medicine
John B. Kostis, Robert Wood Johnson Medical School
Suzanne Oparil, University of Alabama at Birmingham
Anjay Rastogi, University of California, Los Angeles
Christianne Roumie, Tennessee Valley Health Care Center & Vanderbilt University
Rukmani Sahay, Stanford University School of Medicine
Randall S. Stafford, Stanford University School of Medicine
Addison A. Taylor, Michael E. DeBakey VA Medical Center and Baylor College of Medicine
Jackson T. Wright Jr., Case Western Reserve University
Glenn M. Chertow, Stanford University School of Medicine
For the SPRINT Study Research Group

Corresponding Author:

Tara I. Chang, MD, MS
Assistant Professor of Medicine
Stanford Division of Nephrology
777 Welch Road Suite DE
Palo Alto, CA 94304
650-724-1297 (phone)
650-721-1443 (fax)
tichang@stanford.edu

Table S1. Baseline antihypertensive medication use in 7582 SPRINT participants

prescribed at least 1 antihypertensive medication.

| Variable | % |
|---|----------|
| Number of antihypertensive agents | |
| 1 | 32.1 |
| 2 | 39.1 |
| 3+ | 28.7 |
| Thiazide-type diuretic | 43.2 |
| Angiotensin converting enzyme inhibitor | 41.0 |
| Angiotensin II receptor blocker | 23.6 |
| Beta-blocker | 39.5 |
| Dihydropyridine calcium channel blocker | 33.0 |
| Non-dihydropyridine calcium channel blocker | 6.0 |
| Loop diuretic | 5.0 |
| Mineralocorticoid receptor antagonists | 2.1 |
| Direct renin inhibitors | 0.4 |
| Alpha-blocker | 4.4 |
| Centrally acting agent | 2.4 |
| Direct vasodilator | 1.7 |




Figure S1: Prevalence of baseline thiazide prescription in 7582 SPRINT participants prescribed at least 1 antihypertensive medication at baseline, among specified subgroups based on demographics.




Figure S2: Prevalence of baseline thiazide prescription in 7582 SPRINT participants prescribed at least 1 antihypertensive medication at baseline, among specified subgroups based on comorbid conditions

Abbreviation: eGFR = estimated glomerular filtration rate, in mL/min per 1.73m²