

Purity matters: A workflow for the valid high-resolution lipid profiling of mitochondria from cell culture samples

Lisa Kappler^{1,+}, Jia Li^{2,+}, Hans-Ulrich Häring^{1,3,4}, Cora Weigert^{1,3,4}, Rainer Lehmann^{1,3,4}, Guowang Xu^{2,*}, Miriam Hoene^{1,*}

¹Division of Clinical Chemistry and Pathobiochemistry, Department of Diagnostic Laboratory Medicine, University Hospital Tuebingen, Tuebingen, Germany

²Key Laboratory of Separation Science for Analytical Chemistry, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, China


³Department of Molecular Diabetology, Institute for Diabetes Research and Metabolic Diseases of the Helmholtz Centre Munich at the University of Tuebingen, Tuebingen, Germany

⁴German Center for Diabetes Research (DZD), Tuebingen, Germany

⁺Lisa Kappler and Jia Li contributed equally to this study.

^{*}Corresponding Authors.

SUPPLEMENT



Supplementary Figure 1: Correlation of the sum of cardiolipins (CL) with the content of Mitochondrial ATP synthase 5 (ATP5) protein as determined by western blot analysis ($r^2 = 0.79$, $p < 0.0001$).