

THE LANCET

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Jagger C, Matthews FE, Wohland P, et al, on behalf of the Medical Research Council Cognitive Function and Ageing Collaboration. A comparison of health expectancies over two decades in England: results of the Cognitive Function and Ageing Study I and II. *Lancet* 2015; published online Dec 8. [http://dx.doi.org/10.1016/S0140-6736\(15\)00947-2](http://dx.doi.org/10.1016/S0140-6736(15)00947-2).

Supplementary material

This material formed part of the original submission and has been peer reviewed.

Supplement to: Jagger C, Matthews FE, Wohland P, et al, on behalf of the Medical Research Council Cognitive Function and Ageing Collaboration. A comparison of health expectancies from the Cognitive Function and Ageing Studies over two decades in England. *Lancet* 2015.

Supplementary Table 1: Prevalence of cognitive impairment by severity (no impairment MMSE 26-30, mild impairment MMSE 18-25, severe impairment MMSE 0-17), CFAS I and CFAS II, by age group and sex

| | No cognitive impairment | | Mild impairment | | Severe impairment | |
|--------------------|-------------------------|------------------------|------------------------|------------------------|------------------------|-----------------------|
| | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) |
| Men | | | | | | |
| 65-69 years | 83.1 (80.6 to 85.4) | 87.9 (86.2 to 89.5) | 16.0 (13.6 to 18.4) | 11.5 (9.9 to 13.1) | 1.0 (0.4 to 1.6) | 0.6 (0.2 to 1.0) |
| 70-74 years | 78.6 (75.7 to 81.4) | 84.3 (82.4 to 86.2) | 20.0 (17.3 to 22.8) | 14.7 (12.8 to 16.5) | 1.4 (0.6 to 2.2) | 1.0 (0.5 to 1.6) |
| 75-79 years | 66.2 (62.8 to 69.7) | 76.5 (74.2 to 78.9) | 30.3 (26.9 to 33.6) | 20.9 (18.6 to 23.1) | 3.5 (2.2 to 4.8) | 2.6 (1.7 to 3.5) |
| 80-84 years | 55.2 (50.7 to 59.6) | 71.6 (68.7 to 74.5) | 39.8 (35.4 to 44.2) | 23.3 (20.6 to 26.1) | 5.0 (3.1 to 7.0) | 5.0 (3.6 to 6.5) |
| 85+ years | 36.6 (30.3 to 43.0) | 56.5 (52.7 to 60.3) | 43.2 (36.7 to 49.7) | 37.2 (33.5 to 40.9) | 20.2 (14.9 to 25.4) | 6.3 (4.4 to 8.1) |
| Women | | | | | | |
| 65-69 years | 75.6 (73.0 to 78.1) | 86.9 (85.3 to 88.6) | 23.6 (21.1 to 26.1) | 12.7 (11.1 to 14.4) | 0.9 (0.3 to 1.4) | 0.4 (0.1 to 0.6) |
| 70-74 years | 70.5 (67.7 to 73.3) | 80.2 (78.2 to 82.1) | 27.8 (25.1 to 30.5) | 18.7 (16.8 to 20.7) | 1.7 (0.9 to 2.5) | 1.1 (0.6 to 1.6) |
| 75-79 years | 60.4 (57.5 to 63.4) | 72.6 (70.3 to 74.9) | 36.1 (33.2 to 38.9) | 25.0 (22.8 to 27.3) | 3.5 (2.4 to 4.6) | 2.3 (1.6 to 3.1) |
| 80-84 years | 41.8 (38.6 to 45.0) | 61.5 (58.8 to 64.1) | 46.1 (42.9 to 49.3) | 32.6 (30.1 to 35.2) | 12.1 (10.0 to 14.2) | 5.9 (4.6 to 7.2) |
| 85+ years | 24.5 (21.3 to 27.7) | 46.3 (43.5 to 49.1) | 50.8 (47.2 to 54.5) | 42.0 (39.2 to 44.7) | 24.7 (21.5 to 27.8) | 11.7 (9.9 to 13.6) |

Supplementary Table 2: Prevalence of self-perceived health (SPH), CFAS I and CFAS II, by age group and sex

| | Excellent or good SPH | | Fair SPH | | Poor SPH | |
|--------------------|------------------------|------------------------|------------------------|------------------------|----------------------|-----------------------|
| | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) |
| Men | | | | | | |
| 65-69 years | 71.2 (68.3 to 74.1) | 74.3 (72.2 to 76.5) | 22.0 (19.3 to 24.6) | 19.1 (17.1 to 21.0) | 6.8 (5.2 to 8.4) | 6.6 (5.4 to 7.9) |
| 70-74 years | 68.0 (64.8 to 71.2) | 74.4 (72.1 to 76.7) | 26.1 (23.0 to 29.1) | 19.3 (17.2 to 21.3) | 5.9 (4.3 to 7.6) | 6.3 (5.0 to 7.6) |
| 75-79 years | 65.9 (62.4 to 69.4) | 71.1 (68.5 to 73.6) | 26.2 (23.0 to 29.5) | 23.1 (20.7 to 25.4) | 7.9 (5.9 to 9.8) | 5.9 (4.6 to 7.2) |
| 80-84 years | 63.5 (59.1 to 67.8) | 71.4 (68.5 to 74.3) | 29.2 (25.1 to 33.3) | 23.1 (20.3 to 25.8) | 7.3 (5.0 to 9.7) | 5.5 (4.0 to 7.0) |
| 85+ years | 66.1 (59.6 to 72.5) | 67.6 (64.0 to 71.1) | 28.0 (21.9 to 34.1) | 26.5 (23.1 to 29.8) | 5.9 (2.7 to 9.1) | 5.9 (4.2 to 7.7) |
| Women | | | | | | |
| 65-69 years | 70.2 (67.5 to 72.9) | 78.1 (76.0 to 80.1) | 23.2 (20.7 to 25.7) | 17.0 (15.1 to 18.8) | 6.7 (5.2 to 8.2) | 4.9 (3.9 to 6.0) |
| 70-74 years | 69.1 (66.3 to 71.9) | 71.2 (69.0 to 73.5) | 25.0 (22.3 to 27.6) | 23.5 (21.4 to 25.6) | 6.0 (4.5 to 7.4) | 5.3 (4.2 to 6.4) |
| 75-79 years | 65.1 (62.2 to 67.9) | 66.3 (63.9 to 68.7) | 28.0 (23.0 to 29.5) | 28.1 (25.8 to 30.5) | 7.0 (5.4 to 8.5) | 5.5 (4.4 to 6.7) |
| 80-84 years | 62.2 (59.0 to 65.4) | 63.0 (60.4 to 65.7) | 31.1 (28.1 to 34.2) | 29.6 (27.1 to 32.1) | 6.7 (5.1 to 8.3) | 7.3 (5.9 to 8.7) |
| 85+ years | 66.2 (62.5 to 69.8) | 69.1 (66.5 to 71.7) | 27.6 (24.2 to 31.1) | 26.1 (23.6 to 28.6) | 6.2 (4.4 to 8.1) | 4.8 (3.6 to 6.0) |

Supplementary Table 3: Prevalence of disability, CFAS I and CFAS II, by age group and sex

| | No disability | | Mild disability | | Moderate/severe disability | |
|------------------------|------------------------|------------------------|------------------------|------------------------|----------------------------|------------------------|
| | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) | CFAS I % (95% CI) | CFAS II % (95% CI) |
| Men | | | | | | |
| 65-69 years | 88.7 (86.6 to 90.7) | 85.3 (83.5 to 87.1) | 5.1 (3.7 to 6.6) | 9.2 (7.7 to 10.6) | 6.2 (4.6 to 7.7) | 5.5 (4.4 to 6.7) |
| 70-74 years | 86.1 (83.7 to 88.5) | 81.8 (79.8 to 83.8) | 6.2 (4.5 to 7.9) | 11.1 (9.5 to 12.8) | 7.7 (5.9 to 9.6) | 7.0 (5.7 to 8.4) |
| 75-79 years | 77.5 (74.4 to 80.5) | 77.3 (75.0 to 79.7) | 9.8 (7.6 to 12.0) | 12.4 (10.5 to 14.2) | 12.7 (10.3 to 15.2) | 10.3 (8.6 to 12.0) |
| 80-84 years | 63.7 (59.4 to 68.0) | 66.4 (63.3 to 69.4) | 15.1 (11.9 to 18.3) | 17.3 (14.8 to 19.7) | 21.2 (17.6 to 24.8) | 16.4 (14 to 18.7) |
| 85+ years | 37.4 (31.1 to 43.8) | 40.5 (36.7 to 44.3) | 19.1 (14.0 to 24.3) | 26.7 (23.2 to 30.1) | 43.5 (37.0 to 50.0) | 32.8 (29.2 to 36.5) |
| Women | | | | | | |
| 65-69 years | 81.6 (79.3 to 83.9) | 81.4 (79.5 to 83.3) | 11.9 (10.0 to 13.8) | 13.6 (11.9 to 15.3) | 6.5 (5.0 to 7.9) | 5.0 (3.9 to 6.1) |
| 70-74 years | 79.1 (76.6 to 81.5) | 71.3 (69.0 to 73.5) | 14.3 (12.2 to 16.4) | 20.4 (18.4 to 22.4) | 6.6 (5.1 to 8.2) | 8.3 (6.9 to 9.7) |
| 75-79 years | 66.5 (63.7 to 69.3) | 58.1 (55.6 to 60.7) | 17.7 (15.5 to 20.0) | 30.0 (27.6 to 32.3) | 15.7 (13.6 to 17.9) | 11.9 (10.3 to 13.6) |
| 80-84 years | 47.0 (43.8 to 50.2) | 41.0 (38.3 to 43.7) | 22.3 (19.6 to 25.0) | 36.4 (33.8 to 39.0) | 30.7 (27.7 to 33.6) | 22.5 (20.3 to 24.8) |
| 85+ years | 22.7 (19.7 to 25.8) | 19.0 (16.8 to 21.2) | 22.9 (19.8 to 25.9) | 35.1 (32.5 to 37.8) | 54.4 (50.8 to 58.1) | 45.8 (43.1 to 48.6) |

Supplementary Table 4: Life expectancy, cognitive impairment-free life expectancy (CIFLE), and proportion of life free of cognitive impairment at age 85 in 1991 and 2011 and change between 1991 and 2011, by sex

| | 1991 | | 2011 | | Difference (2011-1991) | |
|--|------------------------|------------------------|------------------------|------------------------|-------------------------|--------------------------|
| | Men | Women | Men | Women | Men | Women |
| Life expectancy (years) | 4.2 | 5.3 | 5.2 | 6.6 | 0.9 | 1.4 |
| CIFLE (MMSE 26-30) (95% CI) | 1.5 (1.0 to 2.0) | 1.3 (0.9 to 1.6) | 2.9 (2.5 to 3.3) | 3.0 (2.6 to 3.5) | 1.4 (1.2 to 1.6) | 1.8 (1.6 to 1.9) |
| Proportion of life free of cognitive impairment (95% CI) | 36.2 (24.7 to 47.7) | 24.2 (17.2 to 31.1) | 56.3 (49.0 to 63.5) | 46.0 (39.8 to 52.2) | 20.1 (6.5 to 33.7) | 21.8 (12.5 to 31.1) |
| CILE (MMSE 0-25) (95% CI) | 2.7 (2.4 to 2.9) | 4.0 (3.9 to 4.1) | 2.3 (2.0 to 2.5) | 3.6 (3.4 to 3.8) | -0.4 (-0.9 to 0.1) | -0.4 (-0.7 to -0.1) |
| mildCILE (MMSE 18-25) (95% CI) | 1.9 (1.4 to 2.3) | 2.7 (2.4 to 3.0) | 1.9 (1.5 to 2.4) | 2.8 (2.4 to 3.2) | 0.1 (-0.1 to 0.3) | 0.1 (0.0 to 0.3) |
| Proportion of life with mild cognitive impairment (95% CI) | 43.8 (33.2 to 54.4) | 50.7 (45.0 to 56.3) | 37.4 (29.0 to 45.9) | 42.1 (35.9 to 48.4) | -6.4 (-20.0 to 7.2) | -8.5 (-17.0 to -0.1) |
| sevCILE (MMSE 0-17) (95% CI) | 0.8 (0.3 to 0.5) | 1.3 (1.1 to 1.5) | 0.3 (0.3 to 0.5) | 0.8 (0.6 to 1.0) | -0.5 (-0.8 to -0.2) | -0.5 (-0.8 to -0.3) |
| Proportion of life with severe cognitive impairment (95% CI) | 19.8 (7.0 to 32.5) | 25.0 (18.0 to 31.9) | 6.4 (-4.8 to 17.6) | 11.9 (4.1 to 19.8) | -13.4 (-30.3 to 3.6) | -13.1 (-23.5 to -2.6) |

CILE=years with cognitive impairment, mildCILE=years with mild cognitive impairment, sevCILE=years with moderate-severe cognitive impairment

Supplementary Table 5: Life expectancy, healthy life expectancy (self-perceived health), and proportion of life healthy at age 85 in 1991 and 2011 and change between 1991 and 2011, by sex

| | 1991 | | 2011 | | Difference (2011-1991) | |
|---|------------------------|------------------------|------------------------|------------------------|------------------------|-------------------------|
| | Men | Women | Men | Women | Men | Women |
| Life expectancy (years) | 4.2 | 5.3 | 5.2 | 6.6 | 0.9 | 1.4 |
| HLE (95% CI) | 2.8 (2.6 to 3.1) | 3.5 (3.3 to 3.7) | 3.5 (3.3 to 3.8) | 4.6 (4.3 to 4.8) | 0.7 (0.3 to 1.1) | 1.1 (0.7 to 1.4) |
| Proportion of life spent healthy (95% CI) | 67.1 (60.2 to 73.9) | 66.9 (63.0 to 70.7) | 67.8 (62.8 to 72.8) | 69.3 (65.5 to 73.2) | 0.8 (-7.7 to 9.2) | 2.4 (-3.0 to 7.9) |
| unHLE (95% CI) | 1.4 (1.1 to 1.7) | 1.7 (1.5 to 1.9) | 1.7 (1.4 to 1.9) | 2.0 (1.8 to 2.3) | 0.3 (-0.1 to 0.7) | 0.3 (0.0 to 0.6) |
| fairHLE (95% CI) | 1.2 (0.6 to 1.7) | 1.4 (1.0 to 1.8) | 1.4 (0.9 to 1.8) | 1.7 (1.2 to 2.2) | 0.2 (-0.5 to 0.9) | 0.3 (-0.3 to 0.9) |
| Proportion of life with fair health (95% CI) | 27.3 (14.7 to 39.9) | 27.0 (19.8 to 34.2) | 26.2 (16.9 to 35.5) | 25.8 (18.6 to 33.0) | 1.1 (-14.5 to 16.7) | 1.2 (-9.0 to 11.4) |
| poorHLE (95% CI) | 0.2 (-0.4 to 0.8) | 0.3 (-0.1 to 0.7) | 0.3 (-0.2 to 0.8) | 0.3 (-0.2 to 0.9) | 0.1 (-0.7 to 0.9) | 0.0 (-0.7 to 0.7) |
| Proportion of life with poor health (95% CI) | 5.5 (-8.7 to 19.8) | 6.0 (-2.1 to 14.2) | 6.0 (-4.3 to 16.3) | 5.0 (-3.3 to 13.2) | 0.5 (-17.1 to 18.0) | -1.1 (-12.7 to 10.5) |

HLE=healthy life expectancy, unHLE=years with fair or poor health, fairHLE=years with fair health, poorHLE=years with poor health

Supplementary Table 6: Life expectancy, disability-free life expectancy (DFLE), and proportion of life free of disability at age 85 in 1991 and 2011 and change between 1991 and 2011, by sex

| | 1991 | | 2011 | | Difference (2011-1991) | |
|---|------------------------|------------------------|------------------------|------------------------|--------------------------|-------------------------|
| | Men | Women | Men | Women | Men | Women |
| Life expectancy (years) | 4.2 | 5.3 | 5.2 | 6.6 | 0.9 | 1.4 |
| DFLE (95% CI) | 1.5 (1.3 to 1.8) | 1.2 (1.0 to 1.3) | 2.1 (1.8 to 2.3) | 1.2 (1.1 to 1.5) | 0.5 (0.1 to 0.9) | 0.1 (-0.2 to 0.3) |
| Proportion of life disability free (95% CI) | 36.5 (29.7 to 43.3) | 22.4 (19.2 to 25.6) | 40.1 (34.9 to 45.4) | 18.7 (15.7 to 21.8) | 3.6 (-5.0 to 12.2) | -3.7 (-8.1 to 0.8) |
| DLE (95% CI) | 2.7 (2.4 to 3.0) | 4.1 (3.9 to 4.3) | 3.1 (2.8 to 3.4) | 5.4 (5.2 to 5.6) | 0.4 (0.0 to 0.8) | 1.3 (1.0 to 1.6) |
| mildDLE (95% CI) | 0.8 (0.5 to 1.0) | 1.2 (1.0 to 1.4) | 1.4 (1.1 to 1.6) | 2.3 (2.1 to 2.6) | 0.6 (0.3 to 1.0) | 1.1 (0.8 to 1.4) |
| Proportion of life with mild disability (95% CI) | 17.9 (12.7 to 23.0) | 22.6 (19.4 to 25.9) | 26.6 (21.8 to 31.4) | 35.0 (31.1 to 38.9) | 8.7 (1.7 to 15.8) | 12.4 (7.3 to 17.5) |
| sevDLE (95% CI) | 1.9 (1.6 to 2.2) | 2.9 (2.7 to 3.1) | 1.7 (1.5 to 2.0) | 3.1 (2.7 to 3.3) | -0.2 (-0.6 to 0.2) | 0.2 (-0.2 to 0.5) |
| Proportion of life with severe disability (95% CI) | 45.4 (38.5 to 52.3) | 54.8 (51.0 to 58.6) | 33.4 (28.3 to 38.5) | 46.3 (42.3 to 50.3) | -12.0 (-20.6 to -3.4) | -8.5 (-14.0 to -2.9) |

DLE=years with any disability, mildDLE=years with mild disability, sevDLE=years with moderate-severe disability

Supplementary Table 7: Worked example of reduction in age-specific prevalence of ill-health and increase in healthy life expectancy (HLE) but increases in overall prevalence of ill-health

| Age group | 1991 | | | 2011 | | | Age | Life expectancy ³ | | | HLE ³ | | |
|--------------|-------------------------|--|----------------------|-------------------------|--|----------------------|-----------|------------------------------|------|------------------|------------------|------|------------------|
| | Number (N) ¹ | Prevalence ill-health (%) ² | Number unhealthy (N) | Number (N) ¹ | Prevalence ill-health (%) ² | Number unhealthy (N) | | 1991 | 2011 | change 2011-1991 | 1991 | 2011 | change 2011-1991 |
| 65-69 | 16847 | 2 | 337 | 15093 | 1.6 | 241 | 65 | 16.7 | 20.3 | 3.6 | 15.0 | 18.3 | 3.2 |
| 70-74 | 14492 | 4 | 580 | 13353 | 3.2 | 427 | 70 | 13.2 | 16.4 | 3.2 | 11.5 | 14.3 | 2.8 |
| 75-79 | 12392 | 8 | 991 | 11866 | 6.4 | 759 | 75 | 10.1 | 12.6 | 2.5 | 8.3 | 10.5 | 2.2 |
| 80-84 | 8705 | 16 | 1393 | 10072 | 12.8 | 1289 | 80 | 7.3 | 9.4 | 2.0 | 5.4 | 7.3 | 1.8 |
| 85-89 | 4255 | 32 | 1362 | 6524 | 25.6 | 1670 | 85 | 5.3 | 6.6 | 1.4 | 3.3 | 4.6 | 1.3 |
| 90+ | 1545 | 50 | 773 | 3784 | 40 | 1514 | 90 | 4.4 | 4.2 | -0.1 | 2.2 | 2.5 | 0.4 |
| TOTAL | 58236 | 9.3 | 5435 | 60692 | 9.7 | 5901 | | | | | | | |

¹Actual numbers of women in the three CFAS centres in 1991 and 2011

²Hypothetical age-specific prevalence of ill-health in 1991 with 2% reduction for 2011.

³Life expectancy calculated from true mortality rates in the three CFAS centres in 1991 and 2011. HLE calculated by the Sullivan method as previously using the hypothetical prevalence of ill-health.

Supplementary Figure 1: Life expectancy by age and sex, 1991 and 2011, all centres combined

