

Posterior or Panuveitis Subtype	Number of eyes (percent of total)	Eyes with retinal NV (percent of total)	Prevalence (95% CI)
Vogt-Koyanagi-Harada Syndrome	275 (2.0%)	1 (0.94%)	0.40% (0.0% - 2.0%)
Sympathetic Ophthalmia	96 (0.70%)	0	0.0% (0.0% - 3.8%)
Primary or Secondary Panuveitis	1,314 (9.6%)	20 (1.9%)	1.5% (0.9% - 2.3%)
Retinochoroiditis	183 (1.3%)	1 (0.94%)	0.50% (0.0% - 3.0%)
Multifocal Choroiditis with Panuveitis	414 (3.0%)	2 (1.9%)	0.50% (0.1% - 1.7%)
Other Posterior Uveitis	409 (3.0%)	9 (8.5%)	2.2% (1.0% - 4.0%)
Birdshot Retinochorioretinopathy	345 (2.5%)	1 (0.94%)	0.30% (0.0% - 1.6%)
Retinal Vasculitis	647 (4.7%)	39 (37%)	5.7% (4.1% - 7.7%)

Table 3

Prevalence of retinal neovascularization by posterior or panuveitis subtype. The prevalence of retinal neovascularization was highest in patients with retinal vasculitis (5.7% of cases).