

Title: Inactivated poliovirus type 2 vaccine delivered to rat skin via high density microporation array elicits potent neutralising antibody responses

Authors names and affiliations:

David A. Muller^{1,2}, Frances E. Pearson¹, Germain J.P. Fernando^{1,2}, Christiana Agyei-Yeboah¹, Nick S. Owens¹, Simon R. Corrie^{1,2}, Michael L. Crichton¹, Jonathon C.J. Wei¹, William C. Weldon³, M. Steven Oberste³, Paul R. Young^{2,4}, Mark A. F. Kendall^{1,2*}.

¹Delivery of Drugs and Genes Group (D²G²), Australian Institute for Bioengineering and Nanotechnology, The University of Queensland, Brisbane, Queensland, Australia.

²Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane, Queensland, Australia.

³ Division of Viral Diseases, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, USA.

⁴School of Chemical and Molecular Biosciences, The University of Queensland, Brisbane, Queensland, Australia.

***Corresponding author:** Professor Mark Kendall

Address: The University of Queensland, Delivery of Drugs and Genes Group, Australian Institute for Bioengineering and Nanotechnology, Building 75, St Lucia, Queensland, 4072. Australia

Tel: +61 7 334 64203 Fax: +61 7 334 64197

Email: m.kendall@uq.edu.au

Supplementary data:




Figure S1: Neutralising antibody titres prior to immunisation. Seropositivity was defined as a titre $\geq 3.0 \log_2$ (the dotted line)