

Online supplement A: Systematic reviews, meta-analyses, and guidelines used in literature search.

Brenner H, Stock C, Hoffmeister M. Effect of screening sigmoidoscopy and screening colonoscopy on colorectal cancer incidence and mortality: systematic review and meta-analysis of randomised controlled trials and observational studies. *BMJ* 2014;348:g2467 doi: 10.1136/bmj.g2467 (Published 9 April 2014).

Holme Ø, Bretthauer M, Fretheim A, Odgaard-Jensen J, Hoff G. Flexible sigmoidoscopy versus faecal occult blood testing for colorectal cancer screening in asymptomatic individuals. *Cochrane Database of Systematic Reviews* 2013, Issue 9. Art. No.: CD009259. DOI: 10.1002/14651858.CD009259.pub2.

Screening for colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2008;149:627-37.

Levin B, Lieberman DA, McFarland B, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *Gastroenterology* 2008;134:1570-95.

Rex DK, Johnson DA, Anderson JC, Schoenfeld PS, Burke CA, Inadomi JM. American College of Gastroenterology guidelines for colorectal cancer screening 2009 [corrected]. *Am J Gastroenterol* 2009;104:739-50.

Qaseem A, Denberg TD, Hopkins RH, Jr., et al. Screening for colorectal cancer: a guidance statement from the American College of Physicians. *Ann Intern Med* 2012;156:378-86.

Littlejohn C, Hilton S, Macfarlane GJ, Phull P. Systematic review and meta-analysis of the evidence for flexible sigmoidoscopy as a screening method for the prevention of colorectal cancer. *Br J Surg* 2012;99:1488-500.

Elmunzer BJ, Hayward RA, Schoenfeld PS, Saini SD, Deshpande A, Waljee AK. Effect of flexible sigmoidoscopy-based screening on incidence and mortality of colorectal cancer: a systematic review and meta-analysis of randomized controlled trials. *PLoS Med* 2012;9:e1001352.

Burt RW, Barthel JS, Dunn KB, et al. NCCN clinical practice guidelines in oncology. Colorectal cancer screening. *J Natl Compr Canc Netw* 2010;8:8-61.

Chaparro M, Gisbert JP, Del Campo L, Cantero J, Mate J. Accuracy of computed tomographic colonography for the detection of polyps and colorectal tumors: a systematic review and meta-analysis. *Digestion* 2009;80:1-17.

- de Haan MC, van Gelder RE, Graser A, Bipat S, Stoker J. Diagnostic value of CT-colonography as compared to colonoscopy in an asymptomatic screening population: a meta-analysis. *Eur Radiol* 2011;21:1747-63.
- Haug U, Knudsen AB, Brenner H, Kuntz KM. Is fecal occult blood testing more sensitive for left- versus right-sided colorectal neoplasia? A systematic literature review. *Expert Rev Mol Diagn* 2011;11:605-16.
- Lee BI, Hong SP, Kim SE, et al. Korean guidelines for colorectal cancer screening and polyp detection. *Clin Endosc* 2012;45:25-43.
- Pickhardt PJ, Hassan C, Halligan S, Marmo R. Colorectal cancer: CT colonography and colonoscopy for detection--systematic review and meta-analysis. *Radiology* 2011;259:393-405.
- Winawer SJ, Krabshuis J, Lambert R, O'Brien M, Fried M. Cascade colorectal cancer screening guidelines: a global conceptual model. *J Clin Gastroenterol* 2011;45:297-300.
- Whitlock EP, Lin JS, Liles E, Beil TL, Fu R. Screening for colorectal cancer: a targeted, updated systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2008;149:638-58.
- McFarland EG, Levin B, Lieberman DA, et al. Revised colorectal screening guidelines: joint effort of the American Cancer Society, U.S. Multisociety Task Force on Colorectal Cancer, and American College of Radiology. *Radiology* 2008;248:717-20.
- Niv Y, Hazazi R, Levi Z, Fraser G. Screening colonoscopy for colorectal cancer in asymptomatic people: a meta-analysis. *Dig Dis Sci* 2008;53:3049-54.
- Hewitson P, Glasziou P, Watson E, Towler B, Irwig L. Cochrane systematic review of colorectal cancer screening using the fecal occult blood test (hemoccult): an update. *Am J Gastroenterol* 2008;103:1541-9.
- <http://www.cancer.gov/cancertopics/pdq/screening/colorectal/HealthProfessional/page1/AllPages/Print>. Accessed on 11 March 2013.

Online supplement B: Studies excluded after full manuscript review.

Hardcastle JD, Chamberlain JO, Robinson MH, et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. Lancet 1996;348:1472-7.

Singh H, Turner D, Xue L, Targownik LE, Bernstein CN. Risk of developing colorectal cancer following a negative colonoscopy examination: evidence for a 10-year interval between colonoscopies. JAMA 2006;295:2366-73.

Kronborg O, Jorgensen OD, Fenger C, Rasmussen M. Randomized study of biennial screening with a faecal occult blood test: results after nine screening rounds. Scand J Gastroenterol 2004;39:846-51.

Mandel JS, Bond JH, Church TR, et al. Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. N Engl J Med 1993;328:1365-71.

Brenner H, Chang-Claude J, Seiler CM, Sturmer T, Hoffmeister M. Potential for colorectal cancer prevention of sigmoidoscopy versus colonoscopy: population-based case control study. Cancer Epidemiol Biomarkers Prev 2007;16:494-9.

Selby JV, Friedman GD, Quesenberry CP, Jr., Weiss NS. A case-control study of screening sigmoidoscopy and mortality from colorectal cancer. N Engl J Med 1992;326:653-7.

Atkin WS, Morson BC, Cuzick J. Long-term risk of colorectal cancer after excision of rectosigmoid adenomas. N Engl J Med 1992;326:658-62.

Brenner H, Chang-Claude J, Seiler CM, Sturmer T, Hoffmeister M. Does a negative screening colonoscopy ever need to be repeated? Gut 2006;55:1145-50.

Brenner H, Arndt V, Sturmer T, Stegmaier C, Ziegler H, Dhom G. Long-lasting reduction of risk of colorectal cancer following screening endoscopy. Br J Cancer 2001;85:972-6.

Nishida HU, S. Effectiveness of repeated screening using the fecal occult blood test and its impact on reducing false-negative cancer cases. European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation 2011;20:184-9.

Winawer SJ, Flehinger BJ, Schottenfeld D, Miller DG. Screening for colorectal cancer with fecal occult blood testing and sigmoidoscopy. J Natl Cancer Inst 1993;85:1311-8.

Whynes DK, Mangham CM, Balfour TW, Scholefield JH. Analysis of deaths occurring within the Nottingham trial of faecal occult blood screening for colorectal cancer. *Gut* 2010;59:1088-93.

Hardcastle JDT, W. M.: Chamberlain, J.: Pye, G.: Sheffield, J.: James, P. D.: Balfour, T. W.: Amar, S. S.: Armitage, N. C.: Moss, S. M. Randomised, controlled trial of faecal occult blood screening for colorectal cancer. Results for first 107,349 subjects. *Lancet* 1989;1:1160-4.

Meagher AP, Stuart M. Does colonoscopic polypectomy reduce the incidence of colorectal carcinoma? *Aust N Z J Surg* 1994;64:400-4.

Nakama H. A study on the efficacy of a screening program for colorectal cancer in a small Japanese village. *Clin Investig* 1994;72:117-21.

Murakami R, Tsukuma H, Kanamori S, et al. Natural history of colorectal polyps and the effect of polypectomy on occurrence of subsequent cancer. *Int J Cancer* 1990;46:159-64.

Saito H, Soma Y, Nakajima M, et al. A case-control study evaluating occult blood screening for colorectal cancer with hemoccult test and an immunochemical hemagglutination test. *Oncol Rep* 2000;7:815-9.

Kewenter J, Brevinge H, Engaras B, Haglind E, Ahren C. Results of screening, rescreening, and follow-up in a prospective randomized study for detection of colorectal cancer by fecal occult blood testing. Results for 68,308 subjects. *Scand J Gastroenterol* 1994;29:468-73.

Mandel JS, Church TR, Ederer F, Bond JH. Colorectal cancer mortality: effectiveness of biennial screening for fecal occult blood. *J Natl Cancer Inst* 1999;91:434-7.

Slattery ML, Edwards SL, Ma KN, Friedman GD. Colon cancer screening, lifestyle, and risk of colon cancer. *Cancer Causes Control* 2000;11:555-63.

Bleiberg H, Autier P, Huet F, et al. Colorectal cancer (CRC) screening using sigmoidoscopy followed by colonoscopy: a feasibility and efficacy study on a cancer institute based population. *Ann Oncol* 2006;17:1328-32.

Blom J, Yin L, Liden A, et al. A 9-year follow-up study of participants and nonparticipants in sigmoidoscopy screening: importance of self-selection. *Cancer Epidemiol Biomarkers Prev* 2008;17:1163-8.

Rabeneck L, Lewis JD, Paszat LF, Saskin R, Stukel TA. Risk of proximal and distal colorectal cancer following flexible sigmoidoscopy: a population-based cohort study. *Am J Gastroenterol* 2008;103:2075-82.

Muller AD, Sonnenberg A. Prevention of colorectal cancer by flexible endoscopy and polypectomy. A case-control study of 32,702 veterans. Ann Intern Med 1995;123:904-10.

Newcomb PA, Storer BE, Morimoto LM, Templeton A, Potter JD. Long-term efficacy of sigmoidoscopy in the reduction of colorectal cancer incidence. J Natl Cancer Inst 2003;95:622-5.

Mulder SA, van Soest EM, Dieleman JP, et al. Exposure to colorectal examinations before a colorectal cancer diagnosis: a case-control study. Eur J Gastroenterol Hepatol 2010;22:437-43.

Winawer SJ, Zauber AG, Ho MN, et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workgroup. N Engl J Med 1993;329:1977-81.

Citarda F, Tomaselli G, Capocaccia R, Barcherini S, Crespi M. Efficacy in standard clinical practice of colonoscopic polypectomy in reducing colorectal cancer incidence. Gut 2001;48:812-5.

Brenner H, Chang-Claude J, Seiler CM, Rickert A, Hoffmeister M. Protection from colorectal cancer after colonoscopy: a population-based, case-control study. Ann Intern Med 2011;154:22-30.

Nishihara R, Wu K, Lochhead P, et al. Long-term colorectal-cancer incidence and mortality after lower endoscopy. N Engl J Med 2013 Sep;369:1095-105.

Online supplement C: Included studies.

Jorgensen OD, Kronborg O, Fenger C. A randomised study of screening for colorectal cancer using faecal occult blood testing: results after 13 years and seven biennial screening rounds. *Gut* 2002;50:29-32.

Lindholm E, Brevinge H, Haglind E. Survival benefit in a randomized clinical trial of faecal occult blood screening for colorectal cancer. *Br J Surg* 2008;95:1029-36.

Scholefield JH, Moss SM, Mangham CM, Whynes DK, Hardcastle JD. Nottingham trial of faecal occult blood testing for colorectal cancer: a 20-year follow-up. *Gut* 2012;61:1036-40.

Faivre J, Dancourt V, Lejeune C, et al. Reduction in colorectal cancer mortality by fecal occult blood screening in a French controlled study. *Gastroenterology* 2004;126:1674-80.

Lee KJ, Inoue M, Otani T, Iwasaki M, Sasazuki S, Tsugane S. Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: a prospective cohort study in Japan. *Cancer Detect Prev* 2007;31:3-11.

Malila N, Hakama M, Pukkala E. A 25-year follow-up of a population screened with faecal occult blood test in Finland. *Acta Oncol* 2007;46:1103-6.

Libby G, Brewster DH, McClements PL, et al. The impact of population-based faecal occult blood test screening on colorectal cancer mortality: a matched cohort study. *Br J Cancer* 2012;107:255-9.

Newcomb PA, Norfleet RG, Storer BE, Surawicz TS, Marcus PM. Screening sigmoidoscopy and colorectal cancer mortality. *J Natl Cancer Inst* 1992;84:1572-5.

Hiwatashi N, Morimoto T, Fukao A, et al. An evaluation of mass screening using fecal occult blood test for colorectal cancer in Japan: a case-control study. *Jpn J Cancer Res* 1993;84:1110-2.

Selby JV, Friedman GD, Quesenberry CP, Jr., Weiss NS. Effect of fecal occult blood testing on mortality from colorectal cancer. A case-control study. *Ann Intern Med* 1993;118:1-6.

Lazovich D, Weiss NS, Stevens NG, White E, McKnight B, Wagner EH. A case-control study to evaluate efficacy of screening for faecal occult blood. *J Med Screen* 1995;2:84-9.

Saito H, Soma Y, Koeda J, et al. Reduction in risk of mortality from colorectal cancer by fecal occult blood screening with immunochemical hemagglutination test. A case-control study. *Int J Cancer* 1995;61:465-9.

Zappa M, Castiglione G, Grazzini G, et al. Effect of faecal occult blood testing on colorectal mortality: results of a population-based case-control study in the district of Florence, Italy. *Int J Cancer* 1997;73:208-10.

Bertario L, Russo A, Crosignani P, et al. Reducing colorectal cancer mortality by repeated faecal occult blood test: a nested case-control study. *Eur J Cancer* 1999;35:973-7.

Faivre J, Tazi MA, El Mrini T, Lejeune C, Benhamiche AM, Dassonville F. Faecal occult blood screening and reduction of colorectal cancer mortality: a case-control study. *Br J Cancer* 1999;79:680-3.

Scheitel SM, Ahlquist DA, Wollan PC, Hagen PT, Silverstein MD. Colorectal cancer screening: a community case-control study of proctosigmoidoscopy, barium enema radiography, and fecal occult blood test efficacy. *Mayo Clin Proc* 1999;74:1207-13.

Hoff G, Grotmol T, Skovlund E, Bretthauer M. Risk of colorectal cancer seven years after flexible sigmoidoscopy screening: randomised controlled trial. *Bmj* 2009;338:b1846.

Atkin WS, Edwards R, Kralj-Hans I, et al. Once-only flexible sigmoidoscopy screening in prevention of colorectal cancer: a multicentre randomised controlled trial. *Lancet* 2010;375:1624-33.

Segnan N, Armaroli P, Bonelli L, et al. Once-only sigmoidoscopy in colorectal cancer screening: follow-up findings of the Italian Randomized Controlled Trial--SCORE. *J Natl Cancer Inst* 2011;103:1310-22.

Schoen RE, Pinsky PF, Weissfeld JL, et al. Colorectal-cancer incidence and mortality with screening flexible sigmoidoscopy. *N Engl J Med* 2012;366:2345-57.

Thiis-Evensen E, Hoff GS, Sauar J, Langmark F, Majak BM, Vatn MH. Population-based surveillance by colonoscopy: effect on the incidence of colorectal cancer. Telemark Polyp Study I. *Scand J Gastroenterol* 1999;34:414-20.

Kahi CJ, Imperiale TF, Julian BE, Rex DK. Effect of screening colonoscopy on colorectal cancer incidence and mortality. *Clin Gastroenterol Hepatol* 2009;7:770-5; quiz 11.

Singh H, Nugent Z, Demers AA, Kliewer EV, Mahmud SM, Bernstein CN. The reduction in colorectal cancer mortality after colonoscopy varies by site of the cancer. *Gastroenterology* 2010;139:1128-37.

Jacob BJ, Moineddin R, Sutradhar R, Baxter NN, Urbach DR. Effect of colonoscopy on colorectal cancer incidence and mortality: an instrumental variable analysis. *Gastrointest Endosc* 2012;76:355-64 e1.

Manser CN, Bachmann LM, Brunner J, Hunold F, Bauerfeind P, Marbet UA. Colonoscopy screening markedly reduces the occurrence of colon carcinomas and carcinoma-related death: a closed cohort study. *Gastrointest Endosc* 2012;76:110-7.

Zauber AG, Winawer SJ, O'Brien MJ, et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012;366:687-96.

Baxter NN, Goldwasser MA, Paszat LF, Saskin R, Urbach DR, Rabeneck L. Association of colonoscopy and death from colorectal cancer. *Ann Intern Med* 2009;150:1-8.

Baxter NN, Warren JL, Barrett MJ, Stukel TA, Doria-Rose VP. Association between colonoscopy and colorectal cancer mortality in a US cohort according to site of cancer and colonoscopist specialty. *J Clin Oncol* 2012;30:2664-9.

Muller AD, Sonnenberg A. Protection by endoscopy against death from colorectal cancer. A case-control study among veterans. *Arch Intern Med* 1995;155:1741-8.

Shaukat A, Mongin SJ, Geisser MS, et al. Long-term mortality after screening for colorectal cancer. *N Engl J Med* 2013;369:1106-14.