

Supplementary Materials: Chalcone-Induced Apoptosis through Caspase-Dependent Intrinsic Pathways in Human Hepatocellular Carcinoma Cells

Rodrigo Ramirez-Tagle, Carlos A. Escobar, Valentina Romero, Ignacio Montorfano, Ricardo Armisen, Vincenzo Borgna, Emanuel Jeldes, Luis Pizarro, Felipe Simon and Cesar Echeverria

Table S1. Primary and secondary antibodies used in Western blot experiments.

Primary Ab	Dilution	Source	Incubation Time	Incubation Temperature	Band
Bax	1:500	Rabbit	2 h	RT	Santa Cruz, CA, USA
Bcl-2	1:500	Rabbit	2 h	RT	Santa Cruz, CA, USA
Caspase-3	1:1000	Rabbit	2 h	RT	Cell Signaling, MA, USA
Caspase-8	1:1000	Mouse	2 h	RT	Santa Cruz, CA, USA
Caspase-9	1:1000	Mouse	2 h	RT	Cell Signaling, MA, USA
HSP70	1:5000	Mouse	2 h	RT	Origene, MD, USA
Secondary Ab	Dilution	Source	Incubation Time	Incubation Temperature	Band
Anti-Mouse HRP	1:5000	Goat	2 h	RT	Millipore, MA, USA
Anti-Rabbit HRP	1:5000	Goat	2 h	RT	Millipore, MA, USA

Ab: antibody; RT: room temperature; ON: overnight.

Table S2. Primary and secondary antibodies used in immunocytochemistry.

Primary Ab	Dilution	Source	Incubation Time	Incubation Temperature	Band
Bcl-2	1:100	Rabbit	ON	4 °C	Santa Cruz, CA, USA
Caspase-3	1:100	Rabbit	ON	4 °C	Cell Signaling, MA, USA
α -Tubulin	1:500	Mouse	ON	4 °C	Sigma-Aldrich, Mo, USA
Secondary Ab	Dilution	Source	Incubation Time	Incubation Temperature	Band
Anti-Rabbit Alexa-594	1:250	Goat	ON	4 °C	Invitrogen, CA, USA
Anti-Mouse Alexa-488	1:250	Goat	ON	4 °C	Invitrogen, CA, USA

Ab: antibody; RT: room temperature; ON: overnight.