

Web table A Genes associated with type 2 diabetes: Odds ratios from Whitehall II and the odds ratios from published meta-analysis (where possible), providing weighted odds ratios for the genetic risk function.

Gene	rs number	Genotype in WHII	OR (95% CI)‡ From WHII	Risk allele freq from meta-analysis	Odds ratio from meta-analysis	Comparison	Weight
<i>ADAMTS9</i>	rs4607103	GG GA AA	0.96 (0.79-1.18) P=0.70	0.76	1.09 (1.06-1.12) ¹	Per allele	0.09
<i>BCL11A</i>	rs10490072	AA AG GG	1.12 (0.92-1.37) P=0.25	0.72	1.05 (1.03-1.08) ¹	Per allele	0.049
<i>CALPN10</i>	rs3792267	GG GA AA	0.90 (0.71-1.15) P=0.42	0.27	1.18 (1.05-1.32) ²	GG vs. GA+AA	0.166
<i>CDC123/ CAMK1D</i>	rs12779790	AA AG GG	1.06 (0.85-1.33) P=0.61	0.18	1.11 (1.07-1.14) ¹	Per allele	0.10
<i>CDKAL1</i>	rs17036101	GG GA AA	1.08 (0.74-1.58) P=0.68	0.93	1.15 (1.10-1.21) ³	Per allele	0.140
<i>CDKN2A/2B</i>	rs10811661	AA AG GG	1.22 (0.95-1.56) P=0.13	0.83	1.20 (1.14-1.25) ³	Per allele	0.18
<i>FTO</i>	rs8050136 *	TT TC CC	1.04 (0.88-1.23) 0.67	0.47	1.17 (1.12-1.22) ⁴	Per allele	0.157
<i>HHEX</i>	rs1111875	GG	0.94 (0.79-1.13)	0.65	1.15 (1.10-1.19) ³	Per allele	0.12

		GA AA	P=0.52				
<i>HNF1A</i>	rs1800574	CC CT TT	1.29 (0.79-2.09) P=0.31	0.03	1.31 (1.08-1.57) ⁵	AV+VV vs. AA	0.270
<i>IGF2BP2</i>	rs4402960	CC AC AA	1.16 (0.97-1.40) P=0.11	0.32	1.14 (1.11-1.18) ¹	Per allele	0.13
<i>JAZF1</i>	rs864745	AA AG GG	1.23 (1.03-1.46) P=0.02	0.50	1.10(1.07-1.13) ¹	Per allele	0.095
<i>KCNJ11</i>	rs5215	CC CT TT	1.19 (1.00-1.41) P=0.05	0.56	1.14 (1.10-1.19) ³	Per allele	0.131
<i>NOTCH2</i>	rs10923931	CC AC AA	1.14 (0.87-1.49) P=0.34	0.11	1.13(1.08-1.17) ³	Per allele	0.12
<i>PPARG</i>	rs1801282	CC CG GG	0.90 (0.70-1.15) P=0.39	0.08	1.14 (1.08-1.20) ³	Per allele	0.131
<i>SLC30A8</i>	rs13266634	GG GA AA	0.98 (0.81-1.19) P=0.84	0.69	1.12(1.12-1.19) ³	Per allele	0.11
<i>TCF2 / HNF1B</i>	rs4430796	AA AG GG	1.04 (0.87-1.24) P=0.65	0.47	1.10 (1.07-1.14) ³	Per allele	0.095
<i>TCF7L2</i>	rs7901695	GG GT TT	1.25 (1.04-1.50) P=0.02	0.18	1.37 (1.31-1.43) ¹	Per allele	0.315
<i>THADA</i>	rs7578597	AA	0.99 (0.75-1.29)	0.90	1.15 (1.10-1.20) ³	Per allele	0.14

		AG GG	P=0.91				
<i>TSPAN8/</i> <i>LGR5</i>	rs7961581	AA AG GG	0.94 (0.77-1.14) P=0.54	0.27	1.09(1.06-1.12) [†]	Per allele	0.09
<i>VEGFA</i>	rs9472138	GG GA AA	0.996 (0.82-1.21) P=0.96	0.28	1.06 (1.04-1.09) [†]	Per allele	0.058

[†]from logistic regression adjusted for age and gender.

*LD between rs8050136 and rs9930506 = 0.93

References

- (1) Zeggini E, Scott LJ, Saxena R, Voight BF, Marchini JL, Hu T et al. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat Genet* 2008; 40:638-645.
- (2) Song Y, Niu T, Manson JE, Kwiatkowski DJ, Liu S. Are variants in the CAPN10 gene related to risk of type 2 diabetes? A quantitative assessment of population and family-based association studies. *Am J Hum Genet* 2004; 74:208-222.
- (3) Perry JR, Frayling TM. New gene variants alter type 2 diabetes risk predominantly through reduced beta-cell function. *Curr Opin Clin Nutr Metab Care* 2008; 11:371-377.
- (4) Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H et al. Replication of Genome-Wide Association Signals in U.K. Samples Reveals Risk Loci for Type 2 Diabetes. *Science* 2007; 316:1336-1341.
- (5) Weedon MN, Owen KR, Shields B, Hitman G, Walker M, McCarthy MI et al. A large-scale association analysis of common variation of the HNF1alpha gene with type 2 diabetes in the U.K. Caucasian population. *Diab* 2005; 54:2487-2491.