

### Appendix 3 Characteristics of couples at randomisation by trial [posted as supplied by author]

	Davies 2003 (N=27)	Bhattacharya 2006 (N=24)	Gerris 1999 (N=53)	Lukassen 2005 (N=108)	Martikainen 2001 (N=144)	Thurin 2004 (N=661)	Thurin 2005 (N=42)	van Montfoort 2006 (N=308)
Female age in years, median (IQR)	30.9 (29.0, 33.1)	31.2 (29.2, 34.4)	28.2 (26.0, 30.9)	31.3 (29.3, 32.9)	31.1 (28.3, 33.9)	31.0 (28.7, 33.5)	38.0 (36.8, 38.5)	32.5 (30.4, 34.7)
Missing n (%)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	4 (9.5)	0 (0.0)
Female BMI in kg/m <sup>2</sup> , median (IQR)	24.3 (22.7, 28.5)	26.5 (23.5, 29.5)	23.2 (22.5, 23.9)	22.2 (20.5, 25.1)	22.0 (20.0, 24.5)	23.4 (21.5, 26.4)	24.1 (21.4, 26.2)	23.8 (21.7, 26.2)
Missing n (%)	4 (14.8)	0 (0.0)	50 (94.3)	25 (23.1)	0 (0.0)	3 (0.5)	1 (2.4)	142 (46.1)
Type of infertility, n (%)								
Primary	7 (25.9)	4 (16.7)	33 (62.3)	78 (72.2)	120 (83.3)	480 (72.6)	25 (59.5)	249 (80.8)
Secondary	14 (51.9)	20 (83.3)	9 (17.0)	30 (27.8)	15 (10.4)	181 (27.4)	15 (35.7)	59 (19.2)
Missing	6 (22.2)	0 (0.0)	11 (20.8)	0 (0.0)	9 (6.3)	0 (0.0)	2 (4.8)	0 (0.0)
Duration of infertility in years, median (IQR)	5.3 (2.3, 7.5)	4.0 (3.1, 4.5)	1.5 (1.0, 2.5)	3.2 (2.2, 4.2)	3.0 (2.0, 4.4)	3.0 (3.0, 4.0)	4.0 (3.0, 5.0)	3.0 (2.1, 4.2)
Missing n (%)	6 (22.2)	0 (0.0)	14 (26.4)	0 (0.0)	6 (4.2)	5 (0.8)	0 (0.0)	2 (0.6)
Primary cause of infertility, n (%)								
Male	16 (59.3)	5 (20.8)	27 (50.9)	58 (53.7)	50 (34.7)	215 (32.5)	9 (21.4)	176 (57.1)
Female	7 (25.9)	4 (16.7)	5 (9.4)	20 (18.5)	52 (36.1)	183 (27.7)	15 (35.7)	61 (19.8)
Unexplained	2 (7.4)	6 (25.0)	8 (15.1)	20 (18.5)	21 (14.6)	111 (16.8)	11 (26.2)	71 (23.1)
Mixed/Other	0 (0.0)	9 (37.5)	13 (24.5)	10 (9.3)	21 (14.6)	151 (22.8)	6 (14.3)	0 (0.0)
Missing	2 (7.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.2)	1 (2.4)	0 (0.0)

BMI Body Mass Index

IQR Interquartile range