

1 **SUPPLEMENTAL TABLE**

2 **Table 1.** Factors associated with treatment including calcineurin inhibitors (vs.

3 glucocorticoids alone)

Variables	Univariate		Multivariate	
	OR	95% CI	OR	95% CI
Age at biopsy (per 10 year higher)	<b>0.74</b>	0.66-0.90	0.82	0.74-1.00
Male sex	0.67	0.40-1.12	1.05	0.55-2.04
Black race	<b>1.77</b>	1.06-2.97	1.14	0.59-2.19
Variant				
NOS	1.00		1	
Tip	<b>0.18</b>	0.07-0.47	<b>0.17</b>	0.05-0.53
Collapsing	1.85	0.92-3.70	1.73	0.74-4.02
Baseline eGFR <30 mL/min	0.86	0.40-1.84	1.28	0.51-3.24
Baseline proteinuria ≥3.5 g/d	0.69	0.35-1.36	0.58	0.25-1.36
Serum albumin at biopsy (per g/dL lower)	<b>1.39</b>	1.03-1.85	1.43	0.95-2.17
Edema at baseline	1.08	0.61-1.94	1.25	0.59-2.65
Hypertension at baseline	<b>0.53</b>	0.30-0.94	0.66	0.34-1.28

4 OR, odds ratio; CI, confidence interval; FSGS, focal segmental glomerulosclerosis; NOS,

5 not otherwise specified; eGFR, estimated glomerular filtration rate.

6

7