

**P2X7 Receptor Suppression Preserves Blood-Brain Barrier
through Inhibiting RhoA Activation after Experimental
Intracerebral Hemorrhage in Rats**

Hengli Zhao¹, Xuan Zhang¹, Zhiqiang Dai¹, Yang Feng¹, Qiang Li¹,

John H. Zhang², Xin Liu¹, Yujie Chen^{1, *}, Hua Feng^{1, *}

1 Department of Neurosurgery, Southwest Hospital, Third Military Medical University, Chongqing, China;

2 Department of Anesthesiology, Neurosurgery and Physiology, Loma Linda University, California, USA.

* Corresponding author:

Hua Feng, MD, PhD

Department of Neurosurgery, Southwest Hospital, Third Military Medical University. 29 Gaotanyan Street, Shapingba District, Chongqing, 400038, China.

E-mail: fenghua8888@vip.163.com

Phone: (86)023-68754153, Fax: (86)023-68763954

and


Yujie Chen, MD, PhD

Department of Neurosurgery, Southwest Hospital, Third Military Medical University, Chongqing, China. 29 Gaotanyan Street, Shapingba District,

Chongqing, 400038, China.

E-mail: yujiechen6886@foxmail.com

Phone: (86)023-68765265, Fax: (86)023-68765265



Supplemental Figure 1. Experimental design and animal groups' classification.

ICH: Intracerebral hemorrhage; BBB: Blood brain barrier; P2X7: P2X7 receptor;

EJ: endothelial junction; WB: Western blot; EB: Evans blue extravasation and

fluorescence; FJC: Fluoro-Jade C staining; MRI: Magnetic Resonance Imaging;

IHC: immunohistochemistry; ICV: intracerebralventricular; I.P.:

intraperitoneal; Hematoma: Hematoma volume.