

Ndfip2 is a potential regulator of the iron transporter DMT1 in the liver

Natalie J. Foot, Kelly M. Gembus, Kimberly Mackenzie and Sharad Kumar*

Centre for Cancer Biology, University of South Australia, c/o SA Pathology, PO Box 14,
Rundle Mall, Adelaide, SA 5001, Australia

* Corresponding author: sharad.kumar@unisa.edu.au

Supplementary Material

Supplementary Table 1. Primers used for genotyping and qPCR. All primers were designed to amplify the mouse sequences.

Gene	Primer sequence (5'-3')
<i>Ndfip2</i> genotyping primer "a"	CAGCATTGATGCATGGTCAGC
<i>Ndfip2</i> genotyping primer "b"	GATTACAAATGCTCCTGCAGG
<i>Ndfip2</i> genotyping primer "c"	CAACGGGTTCTTCTGTAGTCC
<i>TBP</i>	F: CAAACCCAGAATTGTTCTCCTT R: ATGTGGTCTTCCTGAATCCCT
<i>Fpn</i>	F: TAAAGTGGCCCAGACGTCAC R: AGCAGACAGTAAGGACCCATC
<i>HAMP</i>	F: ACATTGCGATACCAATGCAGAA R: GCAACAGATACCACACTGGGA