

**Supplementary Table 2 Resource consumption before treatment and after 12 months follow-up. Median (IQR)**

<b>Resources</b>	<b>HA + CS (N=181)</b>	<b>Standard Care (N=95)</b>	<b>P Value*</b>
N. GP visits <i>Before treatment</i>	5 (3-6)	3 (3-4)	0.964
<i>Follow-up</i>	2 (1.5-3)	0 (0-1)	
N. Specialist visits	6 (2-10) 2 (1-5)	6 (4-8) 4 (4-4)	0.644
N. Urine microscopic analysis	5 (3-7) 2 (0-3)	6 (6-8) 4 (1-4)	0.219
N. Urine culture	4.5 (3-7) 2 (1-4)	6 (5-8) 1 (0-3)	<0.001
N. Urine cytology	2 (1-3) 1 (1-1)	3 (1-3) 0 (0-1)	0.003
N. Vaginal swab test	3 (1-3) 0 (0-0)	3 (1-3) 0 (0-0)	0.809
N. Transabdominal ultrasound	2 (1-2) 1 (0-1)	2 (1-2) 0 (0-1)	0.091
N. Transvaginal ultrasound	1 (1-2) 0 (0-0)	2 (1-2) 0 (0-0)	0.938
N. Uroflowmetry	1 (0-1) 0 (0-1)	0 (0-1) 0 (0-1)	0.933
N. Invasive urodynamic test	1 (1-1) 1 (1-1)	0 (0-1) 0 (0-0)	0.777
N. Cystoscopy with bladder biopsy	0 (0-1) 0 (0-0)	0 (0-0) 0 (0-0)	0.234
N. Cystoscopy	1 (1-3) 1 (0-1)	1 (0-1) 0 (0-1)	0.047
N. Antibiotic prescriptions	- 0 (0-1)^	- 1 (0-1)	0.001

IQR = Interquartile range

\* Wilcoxon-Mann-Whitney test on the differences in the before treatment/ follow-up resource consumption between the two groups

^ mean (SD), 0.99 (1.85) in HA + CS vs 1.31 (2.47) in Standard care