

Quercetin Inhibits Fibroblast Activation and Kidney Fibrosis Involving the
Suppression of Mammalian Target of Rapamycin and β -catenin Signaling

Jiafa Ren¹, Jianzhong Li¹, Xin Liu¹, Ye Feng¹, Yuan Gui¹, Junwei Yang¹,
Weichun He¹ and Chunsun Dai^{1,2¶}

¹ Center for Kidney Disease, 2nd Affiliated Hospital, Nanjing Medical University,
262 North Zhongshan Road, Nanjing, Jiangsu, China

² State Key Laboratory of Reproductive Medicine, Institute of Toxicology,
Nanjing Medical University, Nanjing, Jiangsu, China

¶To whom correspondence should be addressed:

Chunsun Dai, MD/PhD, Center for Kidney Diseases, 2nd Affiliated Hospital,
Nanjing Medical University, 262 North Zhongshan Road, Nanjing, Jiangsu, P.R.
China. 210003 (86-25-52636077, E-mail: daichunsun@njmu.edu.cn to Dai C).

Supplemental Figure 1: The general parameters of the mice at 2 weeks after UUO surgery w/o Quercetin treatment. The mice were treated with vehicle or quercetin (25mg/kg.day, ip). At 2 weeks after UUO surgery, the blood from each mouse was harvested. The body weight and the biochemical parameters were examined. (A) Body weight, (B) Serum albumin level, (C) ALT; (D) AST; (E) TB, Total bilirubin; (F) DB, Direct bilirubin. ($n=4$).

