

SUPPLEMENTAL TABLE. Association of migraine and suicidal ideation stratified by depression status

Participants Without Depression	No suicidal ideation (N = 2137)		Suicidal ideation (N = 295)			
	n	%	n	%	Unadjusted OR (95% CI)	Adjusted OR (95% CI) ^a
No migraine	1544	72.3	168	56.9	Reference	Reference
Any migraine	593	27.7	127	43.1	1.97 (1.53-2.53)	1.80 (1.40-2.32)
Types of migraine						
No migraine	1544	72.3	168	56.9	Reference	Reference
Probable migraine	408	19.1	90	30.5	2.03 (1.54-2.68)	1.86 (1.40-2.47)
Migraine	185	8.7	37	12.5	1.84 (1.25-2.71)	1.67 (1.13-2.47)
Participants With Depression	No suicidal ideation (N = 636)		Suicidal ideation (N = 229)			
	n	%	n	%	Unadjusted OR (95% CI)	Adjusted OR (95% CI) ^a
No migraine	373	58.6	99	43.2	Reference	Reference
Any migraine	263	41.4	130	56.8	1.86 (1.37-2.53)	1.75 (1.28-2.39)
Types of migraine						
No migraine	373	58.6	99	43.2	Reference	Reference
Probable migraine	149	23.4	64	27.9	1.62 (1.12-2.34)	1.56 (1.08-2.27)
Migraine	114	17.9	66	28.8	2.18 (1.50-3.18)	2.00 (1.36-2.93)

Abbreviations: OR, odds ratio; CI, confidence interval

a. Adjusted for age, marital status, difficulty paying for the very basics, difficulty paying for medical care