

Lifecourse socioeconomic status and type 2 diabetes: the role of chronic inflammation in the English Longitudinal Study of Ageing

Silvia Stringhini¹, Paola Zaninotto², Meena Kumari³, Mika Kivimaki², G. David Batty²

¹ Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital, Lausanne, Switzerland

² University College London, Department of Epidemiology and Public Health, London, United Kingdom

³ Institute for Social and Economic Research, University of Essex, Essex

Running head: Socioeconomic status and type 2 diabetes at older ages

SUPPLEMENTARY FILES

Table S1. Association of lifecourse socioeconomic score with diabetes incidence ANALYSIS STRATIFIED BY GENDER

	MEN		WOMEN	
	HR (95%CI)	%Δ	HR (95%CI)	%Δ
Model 1: Adjusted for age, sex, ethnicity, prevalent conditions	2.89 (1.76;4.75)		2.29 (1.36;3.87)	
Model 2: Model 1 + lifestyle factors	1.63 (0.96;2.76)	-54	1.51 (0.86;2.62)	-51
Model 3: Model 1 + inflammation	2.23 (1.34;3.69)	-25	1.97 (1.16; 3.36)	-18
Model 4: Model 1 + all factors	1.58 (0.93;2.68)	-57	1.45 (0.83; 2.55)	-55

Δ: Attenuation; BMI: Body Mass Index; CI: Confidence Interval; CRP: C-reactiv protein; HR: Hazard ratio
Lifestyle factors include smoking, alcohol intake, physical activity, BMI. Inflammation include fibrinogen and CRP.