

SUPPLEMENTAL MATERIAL

Supplemental Table

P 2

Table S1. Bleeding risk score calculation in the PRODIGY trial.

CRUSADE SCORE (N=1948)	
Baseline hematocrit, %	
<31	102 (5.2%)
31–33.9	131 (6.7%)
34–36.9	230 (11.8%)
37–39.9	426 (21.9%)
≥40	1059 (54.4%)
Creatinine clearance,* mL/min	
≤15	26 (1.3%)
>15-30	48 (2.5%)
>30-60	537 (5.2%)
>60-90	649 (33.3%)
>90-120	541 (27.8%)
>120	180 (9.2%)
Heart rate (bpm)	
≤70	1231 (63.2%)
71-80	378 (19.4%)
81-90	212 (10.9%)
91-100	97 (5.0%)
101-110	18 (0.9%)
111-120	5 (0.2%)
>120	7 (0.3%)
Sex	
Male	1495 (76.7%)

Female	453 (23.3%)
--------	-------------

Signs of CHF at presentation

No	1836 (94.2%)
Yes	112 (5.8%)

Prior vascular disease

No	1696 (87.1%)
Yes	252 (12.9%)

Diabetes Mellitus

No	1475 (75.7%)
Yes	473 (24.3%)

Systolic blood pressure, mm Hg

≤90	43 (2.2%)
91-100	69 (3.5%)
101-120	225 (11.6%)
121-180	1472 (75.6%)
181-200	85 (4.4%)
>200	54 (2.8%)

ACUITY SCORE (N=1970)

Gender

Male	1517 (77.0%)
Female	453 (23.0%)

Age

<50	151 (7.7%)
-----	------------

50-59	336 (17.1%)
60-69	565 (27.5%)
70-79	649 (32.9%)
≥80	269 (13.7%)

Serum Creatinine

≤1	916 (46.5%)
1-1.2	565 (28.7%)
1.2-1.4	224 (11.4%)
1.4-1.6	136 (6.9%)
1.6-1.8	41 (2.1%)
1.8-2.0	25 (1.3%)
≥2.0	63 (3.2%)

White Blood Cell Count (giga/L)

<10	1475 (74.9%)
10-11	275 (14.0%)
12-13	116 (5.9%)
14-15	56 (2.8%)
16-17	30 (1.5%)
18-19	9 (0.5%)
≥20	9 (0.5%)

Previous Anemia

STEMI	670 (34.0%)
NSTEMI	460 (23.4%)
Normal Biomarkers	840 (42.6%)

Antithrombotic medication Not Applicable

HAS BLED SCORE (N=1968)

Hypertension (sBP >160 mmHg) 1445 (73.4%)

Abnormal renal function (ClCr< 50ml/min) 331 (16.8%)

Abnormal liver function (2 (0.1%)

Elderly (Age> 65y) 1215 (61.7%)

Previous Stroke 14 (0.7%)

Previous Bleeding 12 (0.6%)

Labile INR Not Applicable

Use of drugs 0 (0.0%)

Use of alcohol 1 (0.1%)
